



PHD

## Host response to polymicrobial infections in the lung

Ji, Yuan

*Award date:*  
2019

*Awarding institution:*  
University of Bath

[Link to publication](#)

## Alternative formats

If you require this document in an alternative format, please contact:  
[openaccess@bath.ac.uk](mailto:openaccess@bath.ac.uk)

Copyright of this thesis rests with the author. Access is subject to the above licence, if given. If no licence is specified above, original content in this thesis is licensed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International (CC BY-NC-ND 4.0) Licence (<https://creativecommons.org/licenses/by-nc-nd/4.0/>). Any third-party copyright material present remains the property of its respective owner(s) and is licensed under its existing terms.

### Take down policy

If you consider content within Bath's Research Portal to be in breach of UK law, please contact: [openaccess@bath.ac.uk](mailto:openaccess@bath.ac.uk) with the details. Your claim will be investigated and, where appropriate, the item will be removed from public view as soon as possible.

*Citation for published version:*

Ji, Y 2018, 'Host response to polymicrobial infections in the lung', Ph.D., University of Bath.

*Publication date:*

2018

*Document Version*

Publisher's PDF, also known as Version of record

[Link to publication](#)

## University of Bath

**General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# **Host Response to Polymicrobial Infections in The Lung**

Yuan Ji

A thesis submitted for the degree of Doctor of Philosophy

University of Bath

Department of Pharmacy and Pharmacology

September 2018

## **COPYRIGHT**

Attention is drawn to the fact that copyright of this thesis rests with the author. A copy of this thesis has been supplied on condition that anyone who consults it is understood to recognise that its copyright rests with the author and that they must not copy it or use material from it except as permitted by law or with the consent of the author.

This thesis may be made available for consultation within the University Library and may be photocopied or lent to other libraries for the purposes of consultation.

## Table of Contents

<b>List of figures</b>	<b>5</b>
<b>List of tables</b>	<b>7</b>
<b>Acknowledgement</b>	<b>8</b>
<b>Abstract</b>	<b>9</b>
<b>List of abbreviations</b>	<b>10</b>
<b>Chapter 1 Introduction</b>	<b>13</b>
1.1 <i>Microbial infections of the lower respiratory tract</i>	14
1.1.1 Overview of lower respiratory tract infection	14
1.1.2 Dilemmas of antibiotic resistance	15
1.1.3 Polymicrobial biofilm communities in LRTI	16
1.1.4 Pathogenicity of <i>Staphylococcus aureus</i>	19
1.1.5 Pathogenesis of Bcc	25
1.2 <i>Microbial recognition by the innate immune system and host response</i>	29
1.2.1 Cells of the innate immune system in the airway	29
1.2.2 TLR-mediated pathogen recognition and host defence	35
1.3 <i>Hypothesis and aims</i>	44
<b>Chapter 2 Material and Methods</b>	<b>45</b>
2.1 <i>Human Airway Epithelial Cell Lines</i>	46
2.2 <i>Bacteria Strains</i>	46
2.3 <i>Biofilm Assay</i>	49
2.4 <i>Measurement of Epithelial Barrier Integrity</i>	49
2.4.1 Measurement of Transepithelial Electric Resistance (TEER)	49
2.4.2 Flux of Fluorescein Isothiocyanate (FITC) Labelled-Dextran	50
2.5 <i>Invasion Assay</i>	50
2.6 <i>Western Blot Analysis</i>	51
2.7 <i>Detection of Cytokine Production</i>	52
2.8 <i>Cell Viability Assay</i>	53
2.9 <i>Statistical Analysis</i>	53

2.10	<i>Reagents and Consumables</i> .....	54
<b>Chapter 3 Investigating Host Response to Bacterial Infections in Human Airway Epithelial Cells</b> .....		
		<b>57</b>
3.1	<i>Introduction</i> .....	58
3.2	<i>Airway Barrier Function Is Compromised by Bacterial Pathogens</i> .....	60
3.3	<i>Evaluation of Biofilm Formation by Bcc and S. aureus sp.</i> .....	62
3.4	<i>Modification of IL-8 Production by B. cenocepacia and S. aureus sp. in 16HBE Cells.</i> .....	63
3.5	<i>PI3K is required for the proinflammatory response to B. cenocepacia J2315</i> ..	65
3.6	<i>Discussion</i> .....	66
3.7	<i>Conclusions</i> .....	69
<b>Chapter 4 Investigating the Role of PI3K/Akt, MAPK and NF-κB Signalling in Bacterial Infection of Human Airway Epithelial Cells</b> .....		
		<b>70</b>
4.1	<i>Introduction</i> .....	71
4.2	<i>PI3K activity regulates S. aureus invasion of airway epithelial cells</i> .....	72
4.3	<i>Modulation of PI3K/Akt signalling by S. aureus isolates</i> .....	75
4.4	<i>B. cenocepacia J2315 and S. aureus supernatants modulating PI3K/Akt, MAPK and NF-κB signalling in human airway epithelial cells</i> .....	77
4.5	<i>Discussion</i> .....	84
4.6	<i>Conclusion</i> .....	86
<b>Chapter 5 Investigating Host Response to Polymicrobial Infections in The Airway</b> .....		
		<b>88</b>
5.1	<i>Introduction</i> .....	89
5.2	<i>Suppression of B. cenocepacia-induced IL-8 production by S. aureus supernatants</i> .....	91
5.3	<i>Characterisation of anti-inflammatory factors in S. aureus supernatants</i> .....	97
5.4	<i>16HBE cells are hypo-responsive to Pam<sub>3</sub>CSK4, LPS but not flagellin</i> .....	99
5.5	<i>Discussion</i> .....	104
5.6	<i>Conclusions</i> .....	108

<b>Chapter 6</b>	<b>General discussion</b>	<b>109</b>
6.1	<i>Host responses to monomicrobial infections</i>	110
6.2	<i>Host response to polymicrobial infections and the comparison with monomicrobial infections</i>	112
6.3	<i>Interpretation of the host receptor for S. aureus anti-inflammatory factor(s)</i>	113
6.4	<i>Characterisation of S. aureus immunomodulatory factor(s)</i>	115
6.5	<i>Limitations</i>	115
6.6	<i>Final conclusions</i>	116

## List of figures

Figure 1.1 SpA targets and downstream host signalling.....	22
Figure 1.2 Staphylococcal inhibition of neutrophil activation and chemotaxis.....	23
Figure 1.3 Schematic of the conducting airway and alveolus.....	31
Figure 1.4 MyD88-dependent and MyD88-independent TLR signalling pathways.....	39
Figure 1.5 Negative regulators of MyD88-dependent TLR signalling pathways.....	43
Figure 2.1 Schematic of measuring TEER for airway epithelial cell monolayer.....	50
Figure 3.1 Compromised Calu-3 barrier by MRSA252 or <i>B. multivorans</i> 13243 invasion.....	61
Figure 3.2 Biofilm formation by Bcc and <i>S. aureus</i> strains.....	62
Figure 3.3 IL-8 production and cytotoxicity caused by <i>B. cenocepacia</i> J2315 and <i>S. aureus</i> sp. in 16HBE cells.....	64
Figure 3.4 IL-8 response to <i>S. aureus</i> and Bcc with the presence or absence of PI3K inhibitors in 16HBE cells.....	65
Figure 4.1 Pilot study of adherence to and invasion of airway epithelial 16HBE cells.....	73
Figure 4.2 Inhibition of bacterial adherence and internalisation by pharmacological inhibitors of PI3K.....	74
Figure 4.3 Planktonic MRSA252 or MSSA NCTC 6571 do not affect PI3K activity.....	75
Figure 4.4 Biofilm MRSA252 or MSSA NCTC 6571 do not affect PI3K activity.....	76
Figure 4.5 <i>B. cenocepacia</i> J2315 has no effect on MAPK or NF- $\kappa$ B signalling.....	78
Figure 4.6 Activation of Erk and p38 by MSSA NCTC 6571 supernatant.....	79
Figure 4.7 MAPK activation in response to <i>B. cenocepacia</i> J2315 live bacteria or MSSA NCTC 6571 supernatant.....	80
Figure 4.8 Host signalling modulation by unwashed <i>B. cenocepacia</i> J2315 culture.....	82
Figure 4.9 Comparison of washed and unwashed <i>B. cenocepacia</i> J2315 culture on stimulating 16HBE signalling.....	83
Figure 5.1 The anti-inflammatory effect of <i>S. aureus</i> on <i>B. cenocepacia</i> J2315-induced IL-8.....	92
Figure 5.2 IL-8 response to co-stimulation with <i>B. cenocepacia</i> J2315 and different <i>S. aureus</i> supernatants or heat-killed <i>S. aureus</i> .....	93

Figure 5.3 MAPK response to co-stimulation with <i>B. cenocepacia</i> J2315 and MSSA NCTC 6571 supernatant.....	95
Figure 5.4 PI3K, MAPK and NF- $\kappa$ B response to co-stimulation with <i>B. cenocepacia</i> J2315 and MRSA252 or MSSA NCTC 6571 supernatant.....	96
Figure 5.5 Characterisation of anti-inflammatory effectors in <i>S. aureus</i> supernatants.....	98
Figure 5.6 Pro-inflammatory response to TLR2, 4, or 5 agonists and <i>S. aureus</i> supernatant.....	99
Figure 5.7 Time course of Pam3CSK4 on PI3K, MAPK and NF- $\kappa$ B activation.....	101
Figure 5.8 Time course of flagellin on PI3K, MAPK and NF- $\kappa$ B activation.....	102
Figure 5.9 TLR4 antagonist LPS-RS inhibits <i>B. cenocepacia</i> J2315-induced IL-8 production but not TLR5 antagonist hTLR5 fc.....	103



## List of tables

Table 1.1 Key immune evasion factors of <i>S. aureus</i> .....	20
Table 1.2 Key virulence factors of Bcc.....	28
Table 2.1 Bacterial strains used in this study.....	47
Table 2.2 Antibiotics treatment for bacterial strains.....	51
Table 2.3 Reagents and consumables for mammalian cell and bacteria culture. ....	54
Table 2.4 Drugs for <i>in vitro</i> treatment.....	55
Table 2.5 Reagents for western blot.....	55
Table 2.6 List of primary antibodies.....	56
Table 2.7 List of secondary antibodies.....	56
Table 2.8 Assay kits.....	56
Table 3.1 Translocation in airway epithelial monolayer.....	67

## **Acknowledgement**

I would like to say thank you to many people who have contributed enormously to the work presented in this thesis. Without them, the experience of a PhD would not have been as delightful as it has been, let alone possible.

Firstly, I would like to thank the University of Bath for offering a Fee Waiver. I gratefully thank my dedicated supervisors, Malcolm Watson and Albert Bolhuis, for their tremendous academic support, for giving me so many wonderful opportunities and for encouraging me to step outside of my comfort zone.

Similarly, I would like to express immense gratitude to staff working in the Department of Pharmacy and Pharmacology (past and present): Jo Carter, Kevin Smith, Steve Philips, Ben Groombridge, Sam Goodwin, Pascal Loizeau, and Savannah Britton. Thank you for going far beyond the call of duty, especially with that liquid nitrogen incident. A special thank you to Amel Badoume, Rami Alnajadat, Emma Robson, Robin Wickens, Jo Palandri, Marina Hadjicharalambous, Benoit Roux, Alex Disney, Nour Alhusein and many others who shared office and lab with me, for generously offering their friendship and unwavering cheering. Thank you to my good friends Layla Malt and Chris Danson, not only for always being there for me through the ups and downs, but for letting me be a bridesmaid a month before my submission. A special thank you to Dr Ruth Skrine, who usually makes me a cup of tea and shares vegetables and flowers picked from her spectacular garden. To Shang Chen and Lei Chu, my old friends who are undertaking their PhDs in China, for still being good friends even though they are so far away.

Last, but by no means the least, a huge thank you to my Mum and Dad, who always encourage me to live life to the fullest. How exceptionally lucky I am to have their unconditional love and support - I dedicate this thesis to them.

## Abstract

Lower respiratory tract infection associated with bacterial pathogens is a major cause of morbidity and mortality globally. As a result of the large volume of inhaled air, inevitably the lower respiratory tract is exposed to foreign substances, such as pathogens and allergens. Hence, to maintain homeostasis and sterility, the lower respiratory tract is equipped with multiple barriers and cells with distinct functions. Tightly connected ciliated epithelial cells are the first lining of the conducting airway, covered by a thin layer of mucus. Serving as physical barriers, mucus and the epithelium regulate mucociliary clearance and separate harmful particles from professional immune cells. On the other hand, the chemical barrier consists of secreted products, such as mucin, lytic enzymes, and antimicrobial peptides. In addition to epithelial cells, dendritic cells and macrophages are found in the lower respiratory tract. When epithelial cells are confronted with invading pathogens, bacterial ligands are recognised by receptors such as Toll-like receptors (TLR), which leads to sequential events, comprising activation of signalling cascades, expression and secretion of pro-inflammatory cytokines and chemokines, and recruitment and activation of professional immune cells. To bypass the immune response and facilitate successful infection, pathogens have evolved a wide range of virulence factors that subvert the host response at multiple steps.

The aims of this thesis were to investigate the differences in host responses to mono- and co-infection by common pulmonary pathogens *Staphylococcus aureus* and *Burkholderia cepacia* complex, to interpret the host receptor and signalling cascades targeted by *S. aureus*, and to characterise *S. aureus* immunomodulatory factors. Disruption of the intact airway epithelial barrier and cellular internalisation were observed with both species. *B. cenocepacia* J2315 activated TLR-mediated MAPK and NF- $\kappa$ B signalling pathways, subsequently eliciting a robust interleukin (IL)-8 production. However, when airway epithelial cells were co-treated with *B. cenocepacia* live bacteria and *S. aureus* supernatants, the pro-inflammatory response was subverted by Staphylococcal effectors. This anti-inflammatory effect was widely exhibited in *S. aureus* isolates tested, and was mediated via TLRs but not via the IL-1 receptor or the tumour necrosis factor receptor. TLR4 but not TLR5 was partially blocked by *S. aureus* secreted products, though the TLR4 agonist lipopolysaccharide did not affect IL-8 expression at a high concentration. The Staphylococcal effectors were preliminarily characterised as small, heat-stable, non-proteinaceous, and not cell wall-related factors.

This thesis demonstrates the complex nature of host response in a co-infection model and provides insight into a novel *S. aureus* immune evasion mechanism, as well as a therapeutic intervention of abnormal chronic inflammation.

## List of abbreviations

AP1	Activator protein-1
BCAP	B-cell adaptor for PI3K
Bcc	<i>Burkholderia cepacia</i> complex
CCR	C-C chemokine receptor
CD	Cluster of Differentiation
CF	Cystic fibrosis
CFU	Colony-forming unit
COPD	Chronic obstructive pulmonary disease
CXCR	C-X-C chemokine receptor
CYLD	Cylindromatosis D
DC	Dendritic cell
DUBA	Deubiquitinating enzyme A
EGFR	Epidermal growth factor receptor
Erk	Extracellular signal-regulated kinase
Fab	Fragment antigen-binding
FAK	Focal adhesion kinases
FITC	Fluorescein Isothiocyanate
FLIPr	FPR2-inhibitory protein
FLIPrL	FPR2-inhibitory protein-like
FnBP	Fibronectin-binding proteins
Foxo1	Forkhead box protein O 1
FPR	Formyl peptide receptor
GSK	Glycogen synthase kinase
hBD	Human $\beta$ -defensin
HBSS	Hanks' Balanced Salt Solution
HIg	Haemolysin
HQNO	2-n-heptyl-4-hydroxyquinoline N-oxide
IFN	Interferon
IKK	Inhibitor of NF- $\kappa$ B kinase
IL	Interleukin
IM	Interstitial macrophage
IRAK	IL-1 receptor-associated kinase
IRF	IFN regulatory factor

I $\kappa$ B $\alpha$	Nuclear factor of $\kappa$ light polypeptide gene enhancer in B-cells inhibitor $\alpha$
JAK/SOCS3	Janus kinase 1/Suppressor of cytokine signalling 3
JNK	c-Jun N-terminal kinase
LB	Luria-Bertani
LDH	Lactate dehydrogenase
LPS	Lipopolysaccharide
LRTI	Lower respiratory tract infection
Luk	Leucocidin
Mal	MyD88 adaptor-like
MAPK	Mitogen-activated protein kinase
MCC	Mucociliary clearance
MOI	Multiplicity of infection
MRSA	Methicillin-resistant <i>Staphylococcus aureus</i>
MSSA	Methicillin-sensitive <i>Staphylococcus aureus</i>
MWCO	Molecular weight cut-off
MyD88	Myeloid differentiation primary response 88
NADPH	Nicotinamide adenine dinucleotide phosphate
NF- $\kappa$ B	Nuclear factor $\kappa$ -light-chain-enhancer of activated B cells
NLR	Nucleotide-binding oligomerization domain-like receptor
OD	Optical density
PAMP	Pathogen-associated molecular pattern
PBMC	Peripheral blood mononuclear cells
PI3K/Akt	Phosphoinositide-3-kinase/protein kinase B
PRR	Pattern recognition receptor
PSGL-1	P-selectin glycoprotein ligand-1
PSM	phenol-soluble modulin
PVL	Panton-Valentine Leucocidin
Rab7	Ras-related protein-7
RhoA/ROCK/MLC	RhoA/Rho-associated protein kinase/ myosin light chain
RLR	Retinoid acid-inducible gene-I receptor
RNS	Reactive nitrogen species
ROS	Reactive oxygen species
SCC $mec$	Staphylococcal cassette chromosomal element
SCV	Small-colony variants
SIGIRR	Single immunoglobulin IL-1R-related molecule

SpA	Staphylococcal protein A
SPLI	Secretory leucocyte protease inhibitor
SSL	Staphylococcal superantigen-like proteins
T6SS	Type VI secretion system
TAB	TGF- $\beta$ -activated kinase
TAK	TAK1-binding
TEER	Transepithelial Electric Resistance
TGF	Transforming growth factor
TIR	Toll/IL-1 receptor
TLR	Toll-like receptor
TNFR	Tumour necrosis factor receptor
TRAF	TNF receptor associated factor
TRAM	TRIF-related adaptor molecule
TRIF	TIR domain-containing adapter-inducing IFN- $\beta$

# **Chapter 1 Introduction**

## 1.1 Microbial infections of the lower respiratory tract

### 1.1.1 Overview of lower respiratory tract infection

Lower respiratory tract infection (LRTI) is infection and inflammation caused by bacterial, fungal, parasitic or viral microorganisms in the trachea, bronchi and lungs. Depending on the rate at which it is spontaneously resolved by antibiotic treatments, it is defined as an acute or chronic illness, with cough and other symptoms including sputum production, dyspnoea, wheeze or chest discomfort/pain, with no alternative explanation (Woodhead et al., 2011). In contrast to relative short recovery period and less residual impact on the lower respiratory tract in the case of acute infections, insufficient but unrelenting immune responses and inefficient antibiotic therapies cause persistent chronic infections and exacerbations of pulmonary dysfunction. Chronic LRTI occurs often in patients with chronic obstructive pulmonary disease (COPD), cystic fibrosis (CF) and non-CF bronchiectasis (Pragman et al., 2016). Children younger than 5 years and immunocompromised people are prone to LRTI and have severe illness. LRTI is the leading infectious cause and the fifth-leading cause of morbidity and mortality globally (Troeger et al., 2017). In 2015, mortality due to LRTI was 2.74 million in all ages and 0.7 million deaths in children younger than 5 years. Childhood undernutrition and air pollution are the major risk factors. According to the Global Burden of Disease Study, four aetiologies associated with LRTI are *Haemophilus influenzae* type B, *Streptococcus pneumoniae*, influenza and respiratory syncytial virus infection (Troeger et al., 2017). Although viral causes are the main causes for mild to moderate LRTI, bacterial pneumonia results in higher mortality compared to viral causes. *S. pneumoniae* leads to 55% of LRTI deaths in all ages as the most common cause and *Haemophilus influenzae* type B is mainly attributed to mortality in children younger than 5 years. Fortunately, vaccines against *Haemophilus influenzae* type B and *S. pneumoniae* are available and thus vaccination might reduce the occurrence of bacterial LRTI, in addition to antibiotics. Apart from the aetiologies mentioned above, the opportunistic pathogen *S. aureus* is another common causative agent in LRTI, particularly in hospital- and community-acquired settings, owing to the emergence of multi-drug resistance (Jones, 2010; Udo, 2013). Other common bacterial organisms are *Pseudomonas aeruginosa*, *Klebsiella pneumoniae*, *Escherichia coli*, and *Burkholderia cepacia* complex (Bcc) (Uzoamaka et al., 2017).



### 1.1.2 Dilemmas of antibiotic resistance

Antibiotics are the most commonly used treatment for LRTI. However, there are many limitations and adverse effects, including no effect on viral infections, rapid development of resistance, and side effects such as gastrointestinal disturbances. With the emergence of multi-antibiotic resistant isolates, effective treatments become more complicated, resulting in higher medical cost. The definition of antibiotic resistance is the ability of microbes to grow in the presence of antimicrobial agents at a concentration that would normally inhibit the growth (Brauner et al., 2016). Resistance is usually determined by measuring the minimum inhibitory concentration of planktonic (free floating) microorganisms, which acquire resistance either by chromosomal mutations or horizontal gene transfer of mobile genetic elements (Foster, 2017). Although antibiotic resistance evolves naturally, misuse or overuse of antibiotics accelerates this process. Therefore, it is critical to control and prevent infections, coupled with better stewardship of antibiotics.

*S. aureus* is a Gram-positive, opportunistic pathogen and readily acquires mobile genetic elements to develop antibiotic resistance using a variety of mechanisms (Haaber et al., 2017). Firstly, *S. aureus* acquires resistance to broad-spectrum  $\beta$ -lactams such as penicillin by acquiring plasmids that expresses  $\beta$ -lactamase. Soon after the clinical use of  $\beta$ -lactamase-stable derivatives such as methicillin, *S. aureus* adopted Staphylococcal cassette chromosomal element (SCCmec, encoding penicillin binding protein 2a) which confers resistance to  $\beta$ -lactams. These isolates are called methicillin-resistant *S. aureus* (MRSA) (Foster, 2017; Jevons, 1961). Alternative antibiotics choices include vancomycin (glycopeptide), tetracycline (tetracycline), fusidic acid (steroid-like antibiotic), and linezolid (oxazolidinone). However, resistant phenotypes towards these alternatives have emerged, posing significant challenges on *S. aureus* treatment (Gardete and Tomasz, 2014; Gu et al., 2012; McLaws et al., 2011; Schmitz, 2001).

Another opportunistic pathogen, Bcc is a group of at least 20 closely related Gram-negative, motile bacteria and a causative agent of LRTI, particularly in immunocompromised patients such as those with CF. *B. cenocepacia* and *B. multivorans* are the most common species isolated from CF patients, whereas *B. cepacia* is the most prevalent isolate in non-CF patients (Hauser et al., 2011; Kenna et al., 2017). Due to intrinsic or acquired antibiotic resistance, Bcc is very difficult to eradicate (Rhodes and Schweizer, 2016). Varied antibiotic susceptibility is seen between different species, within the same species, and between CF and non-CF isolates (Kenna et al., 2017). Factors such as efflux pumps, cell wall permeability and remodelling, expression of  $\beta$ -lactamases, and altered penicillin binding proteins all contribute to resistance to a large number of

antibiotics (Everaert and Coenye, 2016; Nitin Kumar et al., 2017; Malott et al., 2012; Podnecky et al., 2015). Antibiotics in clinical use against *Burkholderia* species infections include tobramycin (aminoglycoside), aztreonam and doripenem ( $\beta$ -lactams) (Scoffone et al., 2017; Waters et al., 2017). Combination treatments have also been proposed, such as moxifloxacin-ceftazidime-colistin, or ceftolozane-tazobactam (El-Halfawy et al., 2017; Mazer et al., 2017).

### 1.1.3 Polymicrobial biofilm communities in LRTI

In the natural environment or infection sites, microorganisms rarely exist in monospecies communities (Elias and Banin, 2012; Wolcott et al., 2013). There is a growing appreciation that bacteria, fungi and viruses co-exist in a polymicrobial biofilm community which is defined as a coherent community of multiple microorganisms embedded in a self-produced matrix adhering on a surface (Hall and Mah, 2017). Abundant evidence has shown that pathogens exist within biofilm colonies in the lower respiratory tract of patients with COPD, CF and non-CF bronchiectasis (Hassett et al., 2014; Marsh et al., 2014). Chronic infections caused by polymicrobial biofilms may eventually lead to the loss of pulmonary functions due to recurrent inflammation (LiPuma, 2010; Lynch and Bruce, 2013; Surette, 2014).

To switch from a free-floating mode of growth to a biofilm lifestyle requires environmental and genetic triggers as well as precise gene regulation. For example, subinhibitory concentrations of clindamycin contribute to upregulation of genes related to the stress response, attachment, and extracellular DNA production in *S. aureus*, and consequently lead to increased biofilm formation (Schilcher et al., 2016). Extracellular DNA is extended from the surface from active secretion or controlled cell lysis, and plays a critical role in attachment, maintaining integrity and protection (Okshevsky and Meyer, 2015). After initial attachment, cells in the colony produce an extracellular matrix containing exopolysaccharides, extracellular DNA, proteins and lipids (Mann and Wozniak, 2012). Subsequently, microbes multiply and mature into a more sophisticated community. Within the highly differentiated structure, spatial physiological heterogeneity occurs due to the steep gradient of oxygen, nutrients and waste. Compared to the cells in the outer layer, cells in the biofilm core are slow growing or even dormant (Wood et al., 2013). Finally, under shear stress or environmental factors (for example, antibiotics and immune response), microbes escape from the colony and disperse into the environment to find new optimal living space, which benefits microbial survival and dissemination (Kaplan, 2010).

Microbes dwelling inside biofilms are phenotypically distinct from their planktonic counterparts, and thus they differ in transcriptomes and proteomes (Freiberg et al., 2016; Resch et al., 2006). Among them, increased antibiotic resistance in biofilm phenotype may be the most important challenge (de Oliveira et al., 2016; Singla et al., 2012). Multiple factors contribute to modulating biofilm susceptibility to antibiotics, depending on the microbial strains and species, growing conditions, and developmental stage of the biofilm. The extracellular matrix confers the first layer of protection. A penetration barrier occurs where certain antibiotics such as vancomycin and chloramphenicol have limited penetration of Staphylococcal biofilms (Singh et al., 2016). The matrix also contains antibiotic-modifying enzymes such as  $\beta$ -lactamases, responsible for degrading  $\beta$ -lactam antibiotics (Bowler et al., 2012). Microorganisms are also protected against host response by molecules in extracellular matrix. For instance, alginate, the exopolysaccharide present in the matrix of *P. aeruginosa* biofilms, plays an important role in protecting biofilm cells from phagocytosis, even in the presence of interferon (IFN)- $\gamma$  (Leid et al., 2005). Molecules responsible for adherence and aggregation in *S. epidermidis* biofilm matrix exhibit an immunomodulatory effect on murine macrophage, through inhibiting NF- $\kappa$ B activation and pro-inflammatory cytokine production (Schommer et al., 2011). Of note, dispersal of the biofilm structure by gentle ultrasonication without affecting cell viability restores phagocytosis and the pro-inflammatory response to *S. epidermidis*, even though dispersed bacteria are still coated with these molecules. This indicates that the secreted products in the matrix, as well as intact biofilm architecture, contribute to immune evasion. Overall, the matrix confers protection against antibiotic treatment and host defence to all the dwellers.

In addition to the protection by matrix, dormant cells in the core of biofilm remain at low growth and metabolic rate as a result of low oxygen, nutrient starvation and high levels of waste. Consequently, these cells are less susceptible to antibiotics that target fast-growing cells (Vandecandelaere et al., 2017; Walters et al., 2003). Additionally, the close proximity of cells to each other favours horizontal gene transfer via extracellular DNA or conjugation, which facilitates the exchange of multiple mobile genetic elements that may contain antibiotic resistance genes (Wintersdorff et al., 2016). In fact, biofilms confer a favourable environment as *S. aureus* biofilms exhibit up to ~16,000-fold higher frequency of conjugation compared to planktonic counterparts (Savage et al., 2013). The interactions are in a synergistic or antagonistic manner, and have impact on antibiotic resistance (Weigel et al., 2007). For example, in a dual-microbial biofilm model, *Moraxella catarrhalis* confers protection to *S. pneumoniae* by secreting  $\beta$ -lactamases, and in return, *S. pneumoniae* protects *M. catarrhalis* in a quorum sensing-independent pathway (Perez

et al., 2014). In contrast, antagonistic microbe-microbe interactions also promote resistance. *P. aeruginosa* secretes 2-n-heptyl-4-hydroxyquinoline N-oxide (HQNO), which suppresses the growth of *S. aureus* and ultimately reduces *S. aureus* viability (Orazi and O'Toole, 2017). HQNO leads to *S. aureus* shifting to small-colony variants (SCV) phenotype that is commonly associated with slow growing and persistence, whereby it promotes *S. aureus* biofilm resistance to vancomycin and other antibiotics that target biosynthesis (Proctor et al., 2006). Thus, polymicrobial interactions within biofilms play a critical role in regulating resistance and persistence.

Apart from the effect on antimicrobial resistance, an antagonistic interaction was observed in viability and biofilm formation in a *P. aeruginosa* and *S. aureus* co-culture model (Baldan et al., 2014). Long-term antistaphylococcal cephalixin treatment reduced *S. aureus* colonisation in infants and young children, whilst accelerated *P. aeruginosa* infection without improving clinical outcomes (Stutman et al., 2002). Notably, *S. aureus* is positively associated with 5-year survival in CF, while *P. aeruginosa* and Bcc associated with increased mortality and morbidity in children younger than 7 years old with CF (Liou et al., 2001; Nixon et al., 2001). In addition, *P. aeruginosa* and Bcc limit the microbiome diversity by suppressing the acquisition and persistence of other microorganisms, which correlates with worsen pulmonary function (Granchelli et al, 2018; Coburn et al., 2015).

Polymicrobial infections are also highly resistant to host immune systems (Bjarnsholt et al., 2009). However, due to the complex nature of microbe-microbe and host-microbe interactions, the features of host response to polymicrobial infections remain largely unclear. Different species seem to work in a cooperative manner inside the host. For example, an avirulent oral cavity commensal *Streptococcus* produces hydrogen peroxide (Ramsey and Whiteley, 2009). Once it is sensed by the oral pathogen *Aggregatibacter actinomycetemcomitans*, a genetic response is induced which contributes to the production of the complement resistance protein ApiA. As a result, this interaction facilitates the survival of both species from the killing by human serum. In addition, polymicrobial infections delay wound healing. For example, Dalton et al. (2011) transplanted mature biofilms composed of *P. aeruginosa*, *S. aureus*, *Enterococcus faecalis* and *Finnegoldia magna* onto a murine chronically-wounded model, and observed significantly delayed wound healing at day 8 and increased antimicrobial resistance compared to wounds infected with *P. aeruginosa*.

Taken together, polymicrobial biofilm infections exhibit increased resistance to antibiotic and host defence, and delay wound healing, the net impact of which may lead to a worse outcome of patients. Therefore, understanding the mechanisms of host response to

polymicrobial infections may shed light on the persistent infections and inefficiency of antibiotics and immune clearance, as well as suggest novel clinical treatments.

#### 1.1.4 Pathogenicity of *Staphylococcus aureus*

*S. aureus* is a commensal yet opportunistic human pathogen, harboured by about 30% of individuals (Wertheim et al., 2005). It is frequently isolated from the mucosal surfaces (such as nasopharynx, respiratory tract, urinary tract and gastrointestinal tract) and skin (van Belkum, 2016). While colonisation has different impacts to the host, it is a known risk factor of infections, as about 20% individuals harbor *S. aureus* persistently and a large proportion (60%) of individuals are colonised intermittently (Albrecht et al., 2015; Neha Kumar et al., 2015; Williams, 1963). Host symptoms may vary from asymptomatic carriage to severe infection, which is largely determined by the host response (Greenberg et al., 2018; Montgomery et al., 2014). Acquired in hospital and community settings, *S. aureus* is involved in life-threatening infections, including bacteraemia, sepsis, pneumonia, wound infections and indwelling device-associated infections (Tong et al., 2015).


In addition to the rapid evolvement of antibiotic resistance, *S. aureus* expresses a large arsenal of cell wall-anchored and secreted virulence factors to enhance the potential to cause infectious diseases (key factors are listed in Table 1.1). Two systematic reviews give detailed information on *S. aureus* virulence factors (Foster et al., 2014; Thammavongsa et al., 2015).

Adherence and internalisation are the prerequisite steps to establish infection. They rely on cell wall-anchored virulence factors such as clumping factor A and B (Elkhatib et al., 2015; Hair et al., 2010; Palmqvist et al., 2004), fibronectin-binding proteins A and B (FnBPA and FnBPB) (Foster, 2016), collagen adhesin (Zong et al., 2005), and SpA (Falugi et al., 2013; Jung et al., 2001; Pauli et al., 2014), through binding to extracellular matrix such as fibrinogen, fibronectin and elastin (Jonsson et al., 1991; Roche et al., 2004; Wann et al., 2000). Fibronectin serves as a bridge between  $\alpha_v\beta_1$ -integrin on the host side and FnBPs on *S. aureus* surface, and thereby *S. aureus* enters into non-professional phagocytes by endocytosis (Sinha et al., 1999). Mutations in *fnbA* and *fnbB* significantly reduce renal *S. aureus* burden and mortality of mice, indicating that both FnBPs are crucial for Staphylococcal invasion (Shinji et al., 2011).

**Table 1.1 Key immune evasion factors of *S. aureus*.**

<b>Name</b>	<b>Proposed Function</b>	<b>Target</b>	<b>Reference</b>
Aureolysin	Immune evasion	C3	Laarman et al., 2011
Clumping factor A	Phagocytosis inhibition, adhesion	Fibrinogen, complement factor I, serum apolipoprotein E	Elkhatib et al., 2015; Hair et al., 2010; Palmqvist et al., 2004
Coagulase	Phagocytosis inhibition	Fibrinogen	Cheng et al., 2010
Extracellular complement-binding protein	Complement inhibition	C3d	Hammel et al., 2007a
Extracellular fibrinogen-binding protein	Complement inhibition	C3d	Ricklin et al., 2008
PSMs	Neutrophil chemotaxis, neutrophil lysis	FPR2	Geiger et al., 2012; Kretschmer et al., 2010; Surewaard et al., 2013
PVL	Neutrophil lysis	C5aR	Genestier et al., 2005; Spaan et al., 2013a
SpA	Complement and phagocytosis inhibition, inflammation, B cell superantigen, adaptive immune responses disruption	Immunoglobulin Fcγ and Fab, TNFR1, von Willebrand factor, EGFR	Dossett et al., 1969; Forsgren and Sjoquist, 1966; Gomez et al., 2004; 2007; Hartleib et al., 2000; Sasso et al., 1989
SSL10	Complement and phagocytosis inhibition	IgG, prothrombin, factor Xa, fibrinogen	Itoh et al., 2010; 2013; Patel et al., 2010
SSL3	TLR2 signalling inhibition	TLR2	Bardoel et al., 2012; Yokoyama et al., 2012
Staphylococcal binder of immunoglobulin	Complement inhibition	Immunoglobulin, C3 and factor H	Haupt et al., 2008
Staphylococcal complement inhibitor	Complement inhibition	C3 convertase (C3bBb)	Rooijackers et al., 2005

The key virulence factor SpA (Figure 1.1) anchors on the cell wall and can be released into the extracellular milieu (O'Halloran et al., 2015). While SpA is responsible for immune evasion by binding to Fc $\gamma$  and Fab domains of immunoglobulin (Deisenhofer, 1981; Graille et al., 2000), it contributes to pneumonia and inflammation in a murine model by engaging with TNFR1 and epidermal growth factor receptor (EGFR) (Gomez et al., 2004; Soong et al., 2011). On recognising TNFR1 and EGFR, SpA leads to activation of RhoA/Rho-associated protein kinase/ myosin light chain (RhoA/ROCK/MLC), followed by MAPK signalling and calpain activity in human airway epithelial cells (Soong et al., 2011). The cysteine protease calpains cleave junctional proteins occludin and E-cadherin (Chun and Prince, 2009), and thereby allow *S. aureus* and neutrophil penetration across airway epithelial cell monolayers through the paracellular junctions. However, in mice infected with *S. aureus* intranasally, calpain inhibitor but not EGFR inhibitor, increases Staphylococcal load, which indicates that SpA requires activation of EGFR and TNFR1 as well as proteolytic activity to invade airway epithelial barriers *in vivo*. Of interest, a later paper published shows that the *S. aureus* load is the same in *Tnfr1*<sup>-/-</sup> mice as in wild type mice (Ahn et al., 2014). An experiment that blocks both TNFR1 and EGFR should be done to confirm both receptors are critical for Staphylococcal invasion. Moreover, SpA seems to rely on TNFR1, EGFR and Fas ligands to induce IL-16 secretion in cluster of differentiation (CD)-4<sup>+</sup> cells, which is mediated by Ca<sup>2+</sup> flux and calpain activity (Ahn et al., 2014). Of note, neutralising IL-16 augments phagocytosis and alleviates tissue damage without affecting phagocyte influx, and thus IL-16 promotes pneumonia and excessive inflammation in SpA-mediated infections. Overall, due to a wide range of virulence factors, *S. aureus* is able to shift from innocuous colonisation to pathogenic infections; therefore, anti-adhesion therapies may be applied to prevent Staphylococcal infections (Arya et al., 2015).



**Figure 1.1 SpA targets and downstream host signalling.**

**(A)** SpA can be recognised by multiple host receptors and modulate host signalling cascades in airway epithelial cells and CD4<sup>+</sup> cells, as described in Gomez et al., 2004; Soong et al., 2011; Ahn et al., 2014.


**(B)** Known signalling cascades in airway epithelial cells (Gomez and Prince, 2008).

Own original artwork.

Neutrophils are a major component in host defence against *S. aureus* and one of the first responders recruited to the site of infection. The migration of neutrophils, namely chemotaxis, relies on sensing the gradient of chemoattractant (Li Jeon et al., 2002) and adhesive interactions between endothelium receptors and neutrophil ligands (Choi et al., 2009). *S. aureus* possesses a wide range of ligands that bind to neutrophil surface receptors and disturb cellular homeostasis (Figure 1.2). For example, highly virulent community acquired-MRSA isolates express amphipathic phenol-soluble modulins (PSMs, Table 1.1) which recognise formyl peptide receptor (FPR)-2 on neutrophils (Kretschmer et al., 2010). PSMs elicit a robust pro-inflammatory response as well as neutrophil chemotaxis and subsequently lyse neutrophils. In contrast, *S. aureus* interferes with FPR in the opposite way. Staphylococcal FPR2-inhibitory protein (FLIPr) and its homologue FLIPr-like (FLIPrL) block FPRs and therefore subvert the neutrophil response to FPR agonists (Prat et al., 2006; 2009). Additionally, chemotaxis inhibitory protein of *S. aureus* blocks FPR1 and neutrophil complement receptor C5aR (de Haas et al., 2004). To hinder chemotaxis, Staphylococcal superantigen-like proteins 5 and 11 (SSL5 and



SSL11) recognise P-selectin glycoprotein ligand (PSGL-1) on neutrophils and consequently, disturb the interaction between PSGL-1 and P-selectin on endothelial cells (Bestebroer et al., 2007; Chung et al., 2007). Additionally, SSL3 (Table 1.1) binds to Toll-like receptor (TLR)-2 heterodimers and thereby inhibits TLR2-mediated pro-inflammatory response induced by Staphylococcal lipoteichoic acid and lipopeptides (Bardoel et al., 2012; Yokoyama et al., 2012). Staphopain and SSL10 (Table 1.1) target C-X-C chemokine receptor 2 and 4 (CXCR2 and CXCR4) respectively (Laarman et al., 2012; Walenkamp et al., 2009), which are critical for neutrophil activation and chemotaxis from the bone marrow (C. Martin et al., 2003). Furthermore, *S. aureus* promotes neutrophil lysis through secreting bicomponent pore-forming toxins which bind to the leucocyte membrane and form  $\beta$ -barrel pores that span the lipid bilayer (Alonzo and Torres, 2014; Spaan et al., 2017). Leucocidins consist of LukAB/HG, LukED,  $\gamma$ -haemolysin AB (HlgAB),  $\gamma$ -haemolysin CB (HlgCB) and Panton-Valentine Leucocidin (PVL, Table 1.1). LukAB/HG binds to CD11b subunit of the  $\alpha$ M $\beta$ 2 integrin on neutrophils (DuMont et al., 2013), and LukED binds to CXCR1 and CXCR2 on neutrophils and C-C chemokine receptor type 5 (CCR5) on T cells, macrophages and DCs (Alonzo et al., 2013; Reyes-Robles et al., 2013). HlgAB targets chemokine receptors including CXCR1, CXCR2 and CCR2, whereas HlgCB targets complement receptors C5aR and C5L2 (Spaan et al., 2014). Lastly, frequently exhibited in community acquired-MRSA isolates, PVL recognises C5aR and C5L2 (Spaan et al., 2013a). In addition to inhibiting neutrophil chemotaxis and complement activation, leucocidins (LukED and HlgAB) disrupt the erythrocyte membrane to sequester iron, and thereby acquire essential nutrients for proliferation inside the host (Spaan et al., 2015).



**Figure 1.2 Staphylococcal inhibition of neutrophil activation and chemotaxis.**

*S. aureus* subverts neutrophil extravasation and chemotaxis with multiple antagonists (yellow boxes) targeting neutrophil receptors (own original artwork).

Apart from SpA, *S. aureus* evades complement attack using other virulence factors. First, Gram-positive bacteria are protected by a much thicker peptidoglycan layer (20 nm) compared to Gram-negative bacteria (2.5 nm), which confers protection from membrane attack complex-mediated killing (Joiner et al., 1984). Capsular polysaccharides are reported to play an important role in evading opsonophagocytosis (Luong and Lee, 2002; Thakker et al., 1998; Nanra et al., 2013). Staphylococcal complement inhibitor recognises and blocks C3 convertases, which regulate all three complement activation pathways (Rooijackers et al., 2005). Its homologues extracellular fibrinogen-binding protein and extracellular complement-binding protein (Table 1.1) bind to C3d and consequently inhibit C3 and C5 convertases (Hammel et al., 2007b; Jongerius et al., 2010a; Jongerius et al., 2007).

The fates of intracellular *S. aureus* and the infected host cells are determined by the host cell types, the isolates and the ratio of *S. aureus* to host cells (multiplicity of infection, MOI) (Kubica et al., 2008; Strobel et al., 2016). Following internalisation, *S. aureus* are rapidly engulfed into endosomes (in non-professional phagocytes such as epithelial cells, endothelial cells and fibroblasts) or phagosomes (in professional phagocytes such as neutrophils, macrophages and monocytes), which are intracellular compartments derived from the plasma membrane (Chi et al., 2014). Endosomes and phagosomes fuse with vacuoles and initiate the degradation of intracellular bacteria with proteases (Pillay et al., 2002), antimicrobial peptides (Rosenberger et al., 2004), reactive oxygen species (ROS) and reactive nitric species (RNS) (Fang, 2004), and are highly acidic (Hamaguchi et al., 2014). Nevertheless, *S. aureus* has the ability to survive in the hostile environment, escape into the cytoplasm and manipulate host cell programmed cell death.

To subvert lysozyme degradation of the cell wall peptidoglycan, *S. aureus* produces neutrophil serine protease inhibitors (Stapels et al., 2014). To reduce susceptibility to antimicrobial peptides, *S. aureus* exploits two distinct mechanisms. Due to the commonly cationic nature of antimicrobial peptides, the first tactic is to modify membrane lipids, which results in reduced negative membrane surface charge (Peschel et al., 2001; 1999). In addition, *S. aureus* secretes proteases including aureolysin and staphylokinase, which cleaves cathelicidin LL-37 and neutralises  $\alpha$ -defensin, respectively (Jin et al., 2004; Sieprawska-Lupa et al., 2004).

*S. aureus* secretes antioxidants such as the carotenoid pigment staphyloxanthin, superoxide dismutase, Staphylococcal peroxidase inhibitor, catalase, and L-lactate dehydrogenase to reduce oxidative stress or nitrosative stress (Clauditz et al., 2006; de Jong et al., 2017; Gaupp et al., 2012; Mandell, 1975; Richardson et al., 2008).  $\alpha$  toxin contributes to resistance towards oxidative burst through disengaging the close proximity

of internalised *S. aureus* to mitochondria in macrophages, and thereby protects themselves from mitochondrial ROS, caspase-1 and the antimicrobial metabolite fumarate (Cohen et al., 2018).

Despite phagosome pH being as low as 4 (Cachat et al., 2015), the pH lethal for *S. aureus* is 2 *in vitro* (Chan et al., 1998). In contrast to Gram-negative bacteria such as *E. coli*, phagocytosis and phagosome acidification play critical role in TLR-mediated responses by *S. aureus* (Ip et al., 2010). Unlike other MRSA and methicillin-sensitive *S. aureus* (MSSA) isolates, the highly virulent MRSA USA300 is able to survive in macrophage phagosome at pH 5 (Tranchemontagne et al., 2016). The increased virulence of USA300 is correlated with increased expression of regulatory elements including *agr*, *sarA*, and *saeRS* (Montgomery et al., 2008; Wang et al., 2007). The resistance of USA300 to phagocytosis appears to be triggered by acidification and other environmental cues in the phagosome, which leads to upregulation of *agr* gene locus expression (Tranchemontagne et al., 2016).

As part of the antimicrobial response, host cells initiate apoptosis once infected with bacteria and are subsequently engulfed by macrophages. Compared to other modes of cell death, apoptosis is preferable and non-inflammatory through multiple mechanisms (Szondy et al., 2017). However, *S. aureus* either evades apoptosis through EsxA, a substrate of ESAT-6-like secretion system or switches to necrosis in an *agr*-dependent manner (Korea et al., 2014; van Krüchten et al., 2018). In contrast to apoptosis, necrosis involves the disruption of the plasma membrane and release of cellular contents. Thus, it is highly pro-inflammatory and detrimental to the host, leading to tissue damage (Edinger and Thompson, 2004).

Taken together, the opportunistic pathogen *S. aureus* is capable of shifting from commensal to pathogenic and fully equipped with a plethora of virulence factors to survive and persist in the host.

#### 1.1.5 Pathogenesis of Bcc

As mentioned above, Bcc species are opportunistic pathogens that mainly affect immunocompromised groups such as CF patients, and one of the major challenges is its intrinsic antibiotic resistance. The clinical manifestations of *Burkholderia* colonisation can vary greatly and is unpredictable. Whilst some patients exhibit asymptomatic chronic infection, about 20% of patients have a life-threatening exacerbation of respiratory function among CF and non-CF patients (Isles et al., 1984; Hauser and Orsini, 2015). The

'cepacia syndrome' is characterised by pyrexia, rapidly progressive pulmonary function decline, sepsis and bacteraemia.

Although only a small proportion of CF individuals (2.7%) are colonised by Bcc, the acquisition is a major concern among CF patients, their caregivers and families. It has been acknowledged since the 1980s that certain epidemic strains (for example the ET12 clone and PHDC strain of *B. cenocepacia*) are highly transmissible among patients and even from patients to non-CF individuals (LiPuma et al., 1988; Govan et al., 1993; Holmes et al., 1999). Thus, a strict cohort segregation for CF patients harbouring Bcc was implemented worldwide; however, it caused fear and loneliness to worsen the psychological conditions of these individuals (Duff, 2002). Due to the strict infection control measures, the new acquisition of *B. cenocepacia* dropped dramatically after 1995 and *B. cenocepacia* became less predominant among CF individuals (LiPuma, 2010; Zlosnik et al., 2015). Survival analysis reveals that patients colonised with *B. cenocepacia* have a significantly worse outcome compared to *B. multivorans* (Zlosnik et al., 2015).

Nevertheless, new acquisition of Bcc still occurs in spite of segregation practice. In Queensland, Australia, 75% new cases of the total incidence of Bcc colonisation occurred during in the period 2001-2011. The isolates reveal high genetic diversity and the majority of patients (85%) are associated with unique strains (Ramsay et al., 2013). This indicates the sporadic acquisition of Bcc from independent sources in the post-segregation period, most likely from the environment and can be affected by environmental factors such as the amount of rainfall (Peddayelachagiri et al., 2016; Ramsay et al., 2013).

Bcc species are known to breach the epithelial barrier, leading to bacteraemia, as well as trigger an exacerbated proinflammatory response (Downey et al., 2007; McClean and Callaghan, 2009; Saldias and Valvano, 2009). Moreover, Bcc is capable of survival within bronchial epithelial cells and professional phagocytes or mucus extracellularly in the lumen (Schwab et al., 2014). Intracellular *B. cenocepacia* J2315 (ET12 clone) is sequestered in *B. cenocepacia*-containing vacuoles that are fused with early endosomes. In contrast to heat-inactivated bacteria, which are rapidly degraded in the phagolysosome, live bacteria subvert the phagocytic pathway by delaying phagosomal maturation and fusion with the lysosome by up to 6 h, and thereby maintaining the vacuole at pH 6.4 (Lamothe et al., 2007).

Furthermore, the assembly of Nicotinamide adenine dinucleotide phosphate (NADPH) oxidase complex on live *B. cenocepacia*-containing vacuoles is delayed compared to heat-inactivated *B. cenocepacia* (Keith et al., 2009). NADPH oxidase contributes to ROS by producing the precursor superoxide. This results from the manipulation of Rho family

GTPases by type VI secretion system (T6SS)-derived effectors (Table 1.2) (Rosales-Reyes et al., 2012b). Morphology alteration is observed in murine macrophages by live *B. cenocepacia*, which involves deactivation of Rho GTPases in a T6SS-dependent manner (Flannagan et al., 2012; Rosales-Reyes et al., 2012b). The T6SS-derived effector TecA deamidates Rho GTPases and leads to cytoskeleton disruption and inflammation (Aubert et al., 2016).

While it has been shown that Bcc species readily form strong biofilms *in vitro* (Caraher et al., 2007; Scoffone et al., 2016), Schwab et al. (2014) have reported that only single bacteria or small clusters are observed either in phagocytes or mucus in bronchial sections from CF lung tissues. This may indicate that biofilm formation is not a virulence factor associated with *B. cenocepacia* infections.

To facilitate motility and invasion, Bcc is equipped with whip-like polar flagella (Table 1.2) (Tomich et al., 2002). The flagellin filament of flagella is specifically recognised by TLR5 and the major ligand that induces a pro-inflammatory response in airway epithelial cells (de C Ventura et al., 2008; Hayashi et al., 2001). Consequently, blocking the association between flagellin and TLR5 improves lung injury in CF (Blohmke et al., 2008).

Another major virulence factor, lipopolysaccharide (LPS, Table 1.2) can activate TLR4, leading to a cytokine/chemokine response through activation of MAPK and NF- $\kappa$ B (Bamford et al., 2007). Compared to other Gram-negative bacteria, Bcc LPS has unique properties and confers protection against antibiotics and antimicrobial peptides (Loutet et al., 2006; 2010; Vinion-Dubiel and Goldberg, 2003). The positively charged 4-amino-4-deoxy-arabinose is present in the lipid A and core oligosaccharide, which neutralises the anionic charge of phosphate residues in LPS (De Soyza et al., 2008). Consequently, 4-amino-4-deoxy-arabinose contributes at least partially to the resistance to cationic antibiotics and antimicrobial peptides (Hamad et al., 2012; Scott et al., 1999).

In addition, clinical Bcc isolates are commonly coated extracellular by mucoid exopolysaccharide (Table 1.2) (Herasimenka et al., 2007). Exopolysaccharide are associated with chronicity, and confer protection by inhibiting neutrophil chemotaxis and production of ROS, as well as scavenging ROS (Bylund et al., 2006). However, CF individuals exclusively infected with nonmucoid isolates (such as *B. cenocepacia* J2315) is correlated with a faster rate of lung function decline compared to patients infected with mucoid Bcc (Zlosnik et al., 2011).

Taken together, Bcc species are highly virulent and readily to adapt from natural environment to dwell inside the human host with the assistance of a wide range of virulence factors.

**Table 1.2 Key virulence factors of Bcc**

<b>Virulence factors</b>	<b>Functions</b>	<b>Reference</b>
Cable pili and the 22-kDa adhesin	Targeting cytokeratin 13 and transmigration across the airway epithelium, inflammation	Urban et al., 2005; Goldberg et al., 2011
Exopolysaccharides	Phagocytosis inhibition, chemotaxis inhibition, ROS inhibition	Bylund et al., 2006; Conway et al., 2004
Flagella	Motility, adhesion and invasion, and inflammation	Tomich et al., 2002; Urban et al., 2004
LPS	Inflammation, intrinsic antibiotic/antimicrobial peptides resistance and phagocytosis inhibition	Silipo et al., 2007; Loutet et al., 2006; Vinion-Dubiel and Goldberg, 2003; Saldías et al., 2009
Peptidoglycan-associated lipoprotein	Adhesion and inflammation	Dennehy et al., 2017
Quorum sensing	Regulation of the expression of siderophore, zinc metalloproteases, swarming motility and biofilm formation	Lewenza et al., 1999; Huber et al., 2001; Kooi et al., 200
RpoE	Phagocytosis inhibition, stress response regulator, and polymyxin B resistance	Flannagan and Valvano, 2008
RpoN	Phagocytosis inhibition, biofilm formation, motility, nitrogen metabolism and exopolysaccharide synthesis	Saldías et al., 2008; Fazli et al., 2017; Lardi et al., 2015; Liu et al., 2017
Siderophores and Ftr <sub>Bcc</sub> ABCD	Iron uptake	Mathew et al., 2014; Thomas, 2007
T6SS	Actin cytoskeleton disruption, phagocytosis inhibition, delaying NADPH oxidase complex assembly, regulation of T2SS	Flannagan et al., 2012; Rosales-Reyes et al., 2012a; Rosales-Reyes et al., 2012b
Type IV secretion system	Intracellular survival and replication	Sajjan et al., 2008
Zinc metalloproteases	Degrading extracellular matrix (type IV collagen, fibronectin), neutrophil protease inhibitor, macroglobulin, and antimicrobial peptides	Kooi et al., 2005; Kooi et al., 2006; Kooi and Sokol, 2009

## 1.2 Microbial recognition by the innate immune system and host response

The respiratory tract is the second largest mucosal surface in human body and is constantly exposed to the outer environment. It is anatomically divided into the upper airways, the lower airways (the trachea and bronchi) and alveoli (Patwa and Shah, 2015). The major role of the respiratory system is gas exchange, which occurs at the alveolar-capillary membrane. The blood-air barrier is extremely thin (average thickness of normal lungs, 1.7  $\mu\text{m}$ ) to maximise oxygen diffusion (Divertie et al., 1975). The average volume of gas inhaled into a person's lungs (respiratory minute volume) at rest is about 6 litre/min. Inevitably, small particles and microbes are ventilated into the conducting tract with air, which results in the respiratory system being a favourable entrance for pathogens. Thus, in order to provide warm, hydrated and almost sterile air to the peripheral alveoli, the conducting airways are delicately designed with multiple physical and chemical barriers. True threats must be quickly detected and removed, but unnecessary inflammatory responses should be avoided. Hence, the threshold of sensitivity is critical for normal respiratory functions; otherwise the airway is constantly hyper-responsive and chronically inflamed, causing diseases such as asthma and COPD. Hence, apart from mounting effective immune responses, the respiratory tract is tightly regulated to minimise unfavourable tissue damage and maintain homeostasis of the respiratory system.

### 1.2.1 *Cells of the innate immune system in the airway*

Individual cells types shape and orchestrate the innate immunity. From trachea to terminal alveoli, the conducting airways are predominantly lined with epithelial cells. The mucosa and, directly beneath, the epithelium compartment also contain professional immune cells, including DCs, macrophages, monocytes, neutrophils and mast cells (Yu et al., 2016). These cells serve a surveillance role and are regulated by airway epithelial cells.

#### 1.2.1.1 The multifaceted roles of airway epithelium as the first line of host defence


As the first cell type that comes into contact with environmental agents, the epithelium performs a central role in surveillance, protection and immune regulation (Whitsett and Alenghat, 2015; Weitnauer et al., 2016). The composition and morphology of epithelia vary from distal to proximal. Ciliated pseudostratified epithelia, goblet cells, undifferentiated columnar cells and basal cells constitute the epithelium in the large airways (0 to 5<sup>th</sup> division), while in small airways (6<sup>th</sup> to 23<sup>rd</sup> division) having the same

composition, ciliated epithelia and goblet cells are gradually replaced by club cells (Crystal et al., 2008). The gas exchange chamber alveoli are mainly composed of alveolar type I and type II cells. Squamous type I cells are thin and in close proximity to endothelial cells of respiratory capillaries, covering approximately 95% of the surface areas (Crapo et al., 1982; Weibel, 2015). Cuboidal type II cells are critical for producing surfactant lipids and proteins, as well as being the precursors of type I cells during repair (Barkauskas et al., 2013).

The epithelium (Figure 1.3) in the conducting airways is coated with a thin layer of mucus produced by goblet cells and submucosal glands cells (Ballard and Inglis, 2004; Ma et al., 2018). Normal mucus is a low viscous yet elastic fluid, containing water (about 98%), mucins, proteins, lipids, salts and cell debris (Fahy and Dickey, 2010). Inhaled particles are trapped by mucus, thereby being transported promptly to the pharynx by coordinated cilia beating (Kilburn, 1968). This process is a critical innate defence mechanism, namely mucociliary clearance (MCC). Defects in MCC caused by mucus hyperproduction will result in mucus plugs and entrapped microbial substances, which eventually result in pathological conditions, such as COPD, asthma and bronchiectasis (Evans et al., 2009; Ramos et al., 2014; Sears et al., 2015). In addition, the mucus layer restricts the direct contact between microbial substances and airway epithelial cells, which prevents unnecessary inflammatory responses.

Beneath the mucus layer, airway epithelium and apical junctional complexes form a physical barrier against potential pathogenic threats, controlling paracellular transport, and establishing cell polarity. This barrier is a key component of respiratory tract homeostasis as it limits the access of inflammatory ligands to subepithelial professional immune cells. Cell-cell adhesion is built up with the apical tight junctions and underlying adherens junctions (Georas and Rezaee, 2014). While the extracellular loops of tight junction proteins bind to their counterparts on the adjacent cells, the cytoplasmic domains are connected to the actin cytoskeleton by Zonula occludens-1 (Hartsock and Nelson, 2008). Tight junctions are positively or negatively regulated by different factors, such as Zonula occludens-1 exchange, TLR ligands, microbial infection (Kojima et al., 2013; Ragupathy et al., 2014; Yu et al., 2010). Adherens junctions contain transmembrane nectins and E-cadherin, the latter binding to catenins. Modulation of  $\beta$ -catenin affects the expression of E-cadherin, and thereby affects the morphology, differentiation of epithelial cells, and barrier integrity.





**Figure 1.3 Schematic of the conducting airway and alveolus.**

The conducting airway is composed of ciliated epithelial cells and secretory cells, while the gas exchange chamber alveolus is composed of type I and II alveolar cells. The airway is lined with a periciliary layer and a mucus layer which are responsible for antigen removal cooperating with cilia beating. The alveolus is coated with surfactant which reduces surface tension and has immunomodulatory effects. Under steady state, professional immune cells such as DCs, T cells, and macrophages reside within the respiratory system or in the interstitial space to perform a surveillance role. Upon activation, macrophages and DCs engulf infectious stimuli and present antigen to T cells. Adapted from Fahy and Dickey (2010), and Hussell and Bell (2014).

Airway epithelial cells, secretory cells, and professional immune cells locally produce RNS, surfactants, antimicrobial peptides, and complement products, which are multi-functional components of host defence. These antimicrobial products are described below.

The free radical nitric oxide has bacterial killing effect in the airway surface fluid at physiologic concentration (Workman et al., 2017). In addition, nitric oxide is a messenger regulating airway smooth muscle relaxation and cilia beating frequency, thereby promoting MCC (Venugopalan et al., 1998; Stout et al., 2007).

Surfactants lipids and proteins are important for normal function and physiology of the respiratory system through reducing alveolar surface tension during respiratory cycle (Whitsett et al., 2010). Moreover, surfactants recognise microbial ligands, C1q, TLRs and their co-receptors, thereby promoting opsonisation and phagocytosis, as well as anti-

inflammatory responses (Ariki et al., 2012; Watford et al., 2001). Alveolar type II cells are the major sources of surfactants; club cells have also been reported to produce surfactants (Calkovska et al., 2015).

Antimicrobial peptides are constitutively produced or induced upon infection. Defensins are small cationic molecules (3-5 kDa) and divided into 2 subgroups,  $\alpha$ -defensins and  $\beta$ -defensins (hBDs). In airway epithelial cells, hBD-1 is constitutively expressed and not inducible under inflammatory conditions, whereas hBD-2 expression is only induced by inflammatory stimuli (Kao et al., 2004; Singh et al., 1998). Apart from an antimicrobial effect, hBD-3 is a chemoattractant for macrophage and works synergistically with LL-37 (Abou Alaiwa et al., 2014; García et al., 2001). Furthermore, lysozyme degrades peptidoglycan in Gram-positive bacteria cell wall, and thus is less effective on Gram-negative bacteria (Nash et al., 2006). Lysozyme has synergistic effects with lactoferrin and secretory leucocyte protease inhibitor (SPLI) (Singh et al., 2000). In addition to iron sequestering, lactoferrin impairs biofilm formation, bacterial adhesion, and has immunomodulatory effects (Jahani et al., 2015; Singh, 2004; Siqueiros-Cendon et al., 2014). The antiprotease activity of SPLI protects the epithelium from neutrophil elastase under inflammatory conditions (Hiemstra, 2002).

Lastly, the complement system is activated in response to environmental stimuli in the airway epithelium (Kulkarni et al., 2018). Complement has diverse immunomodulatory functions, including enhancing phagocytosis, connecting the innate and adaptive immunity and regulating homeostasis (Hajishengallis et al., 2017). For example, the level of C3a in sputum isolated from CF individuals is positively correlated with the lung function (Sass et al., 2015). This may due to C3a inhibition on neutrophil recruitment and activity (Shinjyo et al., 2009).

When the amount of foreign antigens in the conducting airways exceeds a certain threshold, to accurately and rapidly recognise invading stimuli, the airway epithelium is equipped with pattern recognition receptors (PRR), which will be described in 1.2.2. Activation of PRR signalling by pathogen associated molecular patterns (PAMP) results in expression of pro-inflammatory cytokines/chemokines, which amplifies the host response such as promoting phagocytosis. Inevitably, antimicrobial mechanisms such as oxidative burst and release of lytic enzymes cause respiratory tissue damage. Therefore, in order to avoid unnecessary damage under homeostatic conditions, the epithelium uses distinct strategies to adjust sensitivity, including secretion of anti-inflammatory molecules, regulating the expression level and asymmetrical expression of PRRs on polarised mucosal barrier.

As a consequence of low or absent expression of TLR2, TLR4, and their co-receptors CD36, CD14, and MD-2, airway epithelial cells fail to initiate a strong pro-inflammatory response to TLR2 and TLR4 ligands (Becker et al., 2000; Ioannidis et al., 2013; Jia et al., 2004; Mayer et al., 2007). The cellular localisation, together with the expression level of TLRs regulate ligand accessibility and tolerance. Asymmetric expression of TLRs on the apical and basolateral surface has been observed in airway epithelium (Ioannidis et al., 2013). Thus, the restricted access to the sensors on the basolateral surface by paracellular junctions only allows incoming pathogens to activate TLRs when they breach the barrier (Humlicek et al., 2007). From bronchi to alveoli, TLR5 expression is shifted from the basolateral to the apical surface, which is correlated with the level of sterility in different parts of the respiratory tract (Shikhagaie et al., 2014). The activation threshold of airway epithelium can be adjusted by pre-exposing to certain stimuli such as respiratory syncytial virus (Monick et al., 2003) or cigarette smoke (Pace et al., 2008), which upregulate the expression and membrane localisation of TLR4, and thereby sensitise airway epithelial cells in response to LPS.

Overall, as a central regulator in host defence, the airway epithelium plays multiple roles to maintain the normal physiology of respiratory system, including barrier function and MCC, secreting antimicrobial and immunomodulatory molecules into the airway surface fluid, surveillance and quickly initiating host defence in cooperation with professional immune cells against infectious agents.

#### 1.2.1.2 The roles of macrophages in host defence against airway infections

Macrophages resident in the lungs at steady state can be divided into interstitial macrophages and alveolar macrophages (Lehnert et al., 1985). Interstitial macrophages can be found in the space between alveolar cells and capillary endothelial cells, while alveolar macrophages are located on the luminal side of alveoli (Figure 1.3). Compared to alveolar macrophages, interstitial macrophages have higher turnover rate, and higher capacity of producing cytokines in response to IFN- $\gamma$  and LPS (Cai et al., 2014). Interstitial macrophages also contribute to homeostasis by generating IL-10 and inhibiting DCs activity (Bedoret et al., 2009).

Alveolar macrophages serve as sentinels and regulators, due to their phagocytic capacity, expression of PRRs, and ability to generate cytokines and chemokines (Kopf et al., 2015). Under homeostatic conditions, alveolar macrophages mediate the tolerance threshold to harmless exogenous antigens (Thepen et al., 1989). The immunomodulatory factors generated by alveolar macrophages directly suppress T cells, include prostaglandins

(Pica et al., 1996; Ruggeri et al., 2000), nitric oxide (Bingisser et al., 1998), IL-10 (Mittal et al., 2015), and TGF- $\beta$  (Yoshimura and Muto, 2011). Alveolar macrophages appear to directly inhibit antigen presentation and migration of DCs (Holt et al., 1993; Jakubzick et al., 2006). By competitively engulfing incoming antigens, alveolar macrophages limit the antigen uptake by DCs and thereby block DCs antigen presentation to T cells (Archambaud et al., 2010).

Alveolar macrophages are in close proximity to the alveolar lumen, and hence their activity is tightly regulated by ligands on the surface of epithelial cells (Hussell and Bell, 2014). By cell-cell interactions, latent TGF- $\beta$  binds to  $\alpha_v\beta_6$ -integrin on airway epithelial cells and is subsequently transformed into bioactive TGF- $\beta$ , which suppresses the expression of cytokines in alveolar macrophages (Sheppard, 2006). However, once activated, alveolar macrophages interrupt the cell-cell interactions and thus the brake on cytokine production by TGF- $\beta$  is released (Morris et al., 2003). In addition, signal-regulatory protein- $\alpha$ , mannose receptor, and CD200 also negatively regulate alveolar macrophages function (Hussell and Bell, 2014; Snelgrove et al., 2008; Zhang et al., 2004).

Once the negative signalling is overridden by infectious agents, alveolar macrophages switch from a tolerogenic mode to inflammatory mode, exhibiting higher phagocytic ability, higher oxidative burst, and stronger expression of pro-inflammatory cytokines. Neutrophils and monocytes are recruited by the signals released from alveolar macrophages (Nomura et al., 2001). To prevent dissemination, alveolar macrophages rapidly engulf apoptotic epithelial cells and apoptotic neutrophils (Rubins, 2003; Grabiec and Hussell, 2016). This process protects the host from toxic products leaked from neutrophils. Once the inflammatory response is unleashed, the outcome depends on the interactions between virulence factors of pathogens and the corresponding host response. Hence, alveolar macrophages serve as a gatekeeper of inflammatory responses in the alveoli.

#### 1.2.1.3 The roles of neutrophils in airway infections

Neutrophils are generated from the bone marrow and circulating in the bloodstream. When infection occurs, airway epithelial cells and alveolar macrophages release inflammatory mediators (such as IL-8, IFNs, C3a, and C5a) and neutrophils respond to the call. Upon interactions between endothelial receptors and the ligands on neutrophils, neutrophils migrate to the site of infection following the concentration gradient of chemoattractants (Thammavongsa et al., 2015).

Once recruited to the battlefield, neutrophils rapidly engulf opsonised bacteria and challenge them with various toxic products in the granules, including antimicrobial

peptides, ROS, nitric oxide, and proteolytic enzymes (van Kessel et al., 2014). Upon phagocytosis, neutrophils initiate apoptosis and release 'eat me' signals (phosphatidylserine) to macrophages, to put a brake on excessive inflammatory responses (Fox et al., 2010). In addition, neutrophils immobilise and kill extracellular bacteria by releasing neutrophil extracellular traps which contains DNA and antimicrobial peptides (Spaan et al., 2013b).

However, the bactericidal activity of neutrophils is equally harmful to the host if out of control. Therefore, in the resolving phase, negative regulators (such as the sialic acid binding Ig-like lectin E, and NADPH oxidase) and macrophage phagocytosis are important to eliminate neutrophils in due course (Davidson et al., 2013; McMillan et al., 2013; Kruger et al., 2015). Prolonged neutrophil presence leads to chronic inflammation, tissue injury, and delayed healing; diseases such as COPD, CF, asthma, and pulmonary fibrosis are associated with the failure to remove apoptotic neutrophils (Hodge et al., 2003; Morimoto et al., 2012; Vandivier et al., 2002).

Overall, neutrophils play a key role in first line defence but require tight regulation to prevent the detrimental effect on the host.

### *1.2.2 TLR-mediated pathogen recognition and host defence*

#### *1.2.2.1 Overview of PRRs*

The sentinels of the lower respiratory system, airway epithelial cells and respiratory-resident phagocytes express a wide range of conserved PRRs to detect exogenous PAMPs and endogenous damage-associated molecular patterns, as well as modulate efficient immune responses. PRRs include TLRs, nucleotide-binding oligomerization domain-like receptors (NLRs), C-type lectins, and retinoid acid-inducible gene-I receptors (RLRs) (Takeuchi and Akira, 2010). Located in the cytoplasm, NLRs recognise bacterial ligands (peptidoglycan, flagellin), viral RNA, and fungi, leading to activation of MAPK and NF- $\kappa$ B signalling, inflammasome formation, and type I IFNs production (Kim et al., 2016). C-type lectins binds a wide range of carbohydrates ligands from fungi, bacteria, and viruses (Hoving et al., 2014). Activation of C-type lectins leads to activation of transcription factors, phagocytosis, and the development of T helper 1 and T helper 17 responses (Hardison and Brown, 2012). RLRs recognises intracellular viral double-stranded RNA, following by activation of IFN regulatory factor (IRF)-3 and NF- $\kappa$ B, and ultimately leads to production of IFN- $\beta$  and antiviral immune responses (Matsumiya and Stafforini, 2010). All PRRs play a key role in sensing exogenous antigens and rapidly initiating the host

response to remove the incoming threats and keeping damage under control. To maintain the homeostasis, PRRs are required to discriminate between self/non-self, commensal/pathogenic antigens, and exhibit tolerance under certain threshold. In the following section, the first discovered PRRs, TLRs, and their interactions with PAMPs will be the focus.

#### 1.2.2.2 Overview of TLRs

The discovery of TLRs in the 1990s was of significant importance to the field of innate immunity and was recognised by awarding the 2011 Nobel Prize in Physiology or Medicine to Bruce Beutler and Jules Hoffmann. TLRs are the key molecules in surveillance, signalling transduction and ultimately induction of immune responses. Their dysfunction is associated with allergic inflammation (Clarke et al., 2014), chronic inflammation, tumorigenesis (Korneev et al., 2017), and autoimmune diseases (Mohammad Hosseini et al., 2015).

So far there have been 10 TLRs (Table 1.3) identified in humans, with each one recognising unique ligands from bacteria, viruses, and fungi. TLRs are widely expressed in leukocytes and non-immune cells (including mucosal epithelial cells, fibroblasts, and endothelial cells), and their expression is regulated by microbial stimuli and cytokines (McClure and Massari, 2014; Zarembek and Godowski, 2002). TLR1, 2, 4-6 are primary transmembrane receptors that sense extracellular antigens, whereas TLR3, 7-10 are expressed on the membrane of endoplasmic reticulum (steady-state) and trafficking to endolytic compartments (active-state) to engage with internalised viral ligands. Despite recognising structurally distinct ligands, TLRs share common architecture. The ligand-binding ectodomain contains leucine-rich repeats, while the cytoplasmic domain contains Toll/IL-1 receptor (TIR) domain for signal transduction (Botos et al., 2011). Upon ligands binding, the extracellular domain dimers form 'm' shape which brings the C-terminal of leucine-rich repeats and intracellular domains in close proximity (Botos et al., 2011). TLRs and downstream signalling cascades are bridged by adaptor proteins, including myeloid differentiation primary response 88 (MyD88), TIR domain-containing adapter-inducing IFN- $\beta$  (TRIF), MyD88 adaptor-like (Mal) (Valkov et al., 2011), TRIF-related adaptor molecule (TRAM), and Sterile- $\alpha$  and HEAT-Armadillo motif-containing protein (Carlsson et al., 2016). All TLRs except TLR3 share MyD88-mediated signalling, whereas TLR3 and TLR4 recruit TRIF; all TLRs lead to activation of MAPK and NF- $\kappa$ B signalling.

**Table 1.3 Human TLRs and their exogenous ligands.**

<b>TLR</b>	<b>Localisation</b>	<b>PAMPs</b>	<b>Origin of the ligands</b>	<b>Reference</b>
TLR1	Cell surface	Triacyl lipopeptides	Gram-negative bacteria, <i>Mycoplasma</i>	Shimizu et al., 2007
TLR2	Cell surface	Lipoprotein, peptidoglycan, lipoteichoic acid, phospholipomannan	Bacteria, <i>Mycoplasma</i> , fungi	Into et al., 2004; Jouault et al., 2003; Schwandner et al., 1999; Love et al., 2010; Muller-Anstett et al., 2010
TLR3	Endolysosome	dsRNA, Poly I:C	Viruses	Alexopoulou et al., 2001
TLR4	Cell surface	LPS	Gram-negative bacteria	Chow et al., 1999
TLR5	Cell surface	Flagellin	Bacteria	Hayashi et al., 2001
TLR6	Cell surface	Diacyl lipoprotein	Gram-positive bacteria	Into et al., 2004
TLR7	Endolysosome	ssRNA, guanosine	Viruses	Diebold et al., 2004; Zhang et al., 2016
TLR8	Endolysosome	ssRNA	Viri, bacteria	Heil et al., 2004; Cervantes et al., 2013
TLR9	Endolysosome	Unmethylated CpG-DNA, chromatin-IgG complexes	Bacteria, viruses	Leadbetter et al., 2002; Rutz et al., 2004
TLR10	Endolysosome	dsRNA	Viruses	Lee et al., 2018


### 1.2.2.3 TLR-mediated signalling pathways

The adaptor protein for most TLRs, MyD88 plays a central role in TLR-mediated signalling. MyD88 deficiency increases bacterial burden and dampers host defence to *S. aureus* infection (Dudek et al., 2016; Takeuchi et al., 2000; Weiss et al., 2005). Activated TLRs recruit MyD88 by interaction between their TIR domains. As shown in Figure 1.4, MyD88 recruits and activates IL-1 receptor-associated kinase (IRAK) 4, via death domain-death domain association. IRAK4 activates IRAK1/2, leading to autophosphorylation of the IRAKs complex and subsequently the activation of TNF receptor associated factor (TRAF) 6 (Lin et al., 2010). TRAF6 forms a ubiquitin chain with E2 enzyme Ubc13-Uev1A, which promotes polyubiquitination of TRAF6 and inhibitor of nuclear factor  $\kappa$ -B kinase (IKK)  $\gamma$

(Chen, 2005). Ubiquitinated TRAF6 activates TGF- $\beta$ -activated kinase (TAK) and TAK1-binding (TAB), whereby the TAK1/TAB complex leads to the activation of the three-tier MAPK module, MAP3K, MAP2K and finally MAPK including stress-related kinase (Erk) 1/2, p38, and c-Jun N-terminal kinase (JNK) (Schröder et al., 2001; Wang et al., 2001; Dhanasekaran et al., 2007). Members of MAPK family control cell proliferation, survival, and differentiation (Li et al., 2003; D'Souza et al., 2008; Khiem et al., 2008). Activation of MAPK facilitates phosphorylation and activation of transcription factor activator protein (AP)-1 (Akira et al., 2006). Activated IKK complex contributes to phosphorylation and degradation of I $\kappa$ B $\alpha$ , depriving its sequestration of p65 NF- $\kappa$ B, and thereby p65 NF- $\kappa$ B translocates into the nucleus and regulates transcription of pro-inflammatory cytokines (Kawai and Akira, 2007). Notably, MAPK and NF- $\kappa$ B exhibit differential activation thresholds in response to TLR4 ligand (Gottschalk et al., 2016). Activation of NF- $\kappa$ B occurs at a much lower dose of lipid A (component of LPS) compared to MAPK. Although activated NF- $\kappa$ B leads to gene expression and macrophage priming, TNF production is only activated in a MAPK-dependent manner. Hence, while NF- $\kappa$ B is responsible for sensing sub-threshold antigens, MAPK tightly controls the sensitivity threshold and production of pro-inflammatory cytokines. In addition to activating MAPK and NF- $\kappa$ B signalling pathways, intracellular TLRs (TLR7, TLR8, and TLR9) regulate type I IFNs production via TRAF6. Upon sensing exogenous nucleic acids sequestered in the endolysosome, recruited MyD88 activates IRAK complex together with TRAF6, followed by activation of IRF7 (Uematsu and Akira, 2007). IRF7 subsequently translocates into the nucleus to mediate type I IFNs expression (Cervantes et al., 2011). The detailed TLR7/8/9-mediated signalling is described by Kawasaki and Kawai (2014).

TLR3 and TLR4 are associated with MyD88-independent pathway and lead to production of type I IFNs and IL-10 (Häcker et al., 2006). Activation of TLR4 via MyD88-dependent signalling results in an early-phase activation of NF- $\kappa$ B. Following TLR4/MD-2/CD14 endocytosis, TRIF and TRAM are recruited to the complex on endolysosome membrane (Fitzgerald et al., 2003; Tanimura et al., 2008). TRAF6 and receptor-interacting protein kinase-1 are recruited to TRIF, resulting in the late-phase activation of NF- $\kappa$ B (Kagan et al., 2008; Weinlich and Green, 2014). Both phases of NF- $\kappa$ B activation are required for pro-inflammatory cytokine production (Kawai and Akira, 2010). TRIF also interacts with non-canonical IKKs TRIF and two protein kinases-binding kinase-1 and IKK $\epsilon$ , through TRAF3, which leads to activation of IRF3 and IRF7, as well as production of type I IFNs and IL-10 (Uematsu and Akira, 2007; Häcker et al., 2011; Ullah et al., 2016).





**Figure 1.4 MyD88-dependent and MyD88-independent TLR signalling pathways.**

The MyD88-dependent TLR signalling pathways mainly result in production of inflammatory cytokines and chemokines, whereas MyD88-independent signalling pathways are responsible for type I IFNs production to promote anti-viral responses. Unlike other TLRs, TLR4 requires both pathways at distinct cellular localization and leads to biphasic activation of NF-κB to induce production of inflammatory cytokines (own original artwork).

#### 1.2.2.4 Function of TLRs

As described above, the main roles of TLRs are sensing infectious stimuli, recognition of self and non-self antigens, and induce the production of cytokines and chemokines. In addition, the outcome of TLR signalling includes modulation of adaptive immunity, and production of antimicrobial peptides.

The activation of TLRs on DCs leads to DC maturation, antigen presentation to antigen-specific T cells, and thereby induces activation and differentiation of T cells (Manicassamy and Pulendran, 2009; Schweighoffer et al., 2017). Hence, the activation of naïve T cells by TLR signalling bridges between innate immunity and adaptive immunity. DC maturation

features increases in major histocompatibility complex, production of costimulatory molecules and pro-inflammatory cytokines. Both MyD88 and TRIF are involved in DC maturation (Bechelli et al., 2016; Hu et al., 2015). The production of type I IFN contributes to cross-priming by DCs (Le Bon et al., 2003). Moreover, TLRs can directly regulate proliferation and cytokine production by T/B cells (Reynolds et al., 2010; Schweighoffer et al., 2017).

TLR-mediated signal transduction leads to production of cytokines (such as IL-1, IL-6, IL-10, IL-13, TNF $\alpha$ , and TGF $\beta$ ), chemokines (such as IL-8, monocyte chemoattractant protein1, macrophage inflammatory protein 2- $\alpha$ , and chemokine ligand 5), and type I IFNs (mainly IFN $\alpha$  and IFN $\beta$ ) (McClure and Massari, 2014). These secreted products play an important role in shaping the innate and adaptive immune responses, inflammation, survival, and proliferation (Li et al., 2010). Some products serve as pro-inflammatory regulators (such as IL-1, IL-8, and TNF $\alpha$ ), whereas IL-10 and TGF $\beta$  suppress the immune response. In addition to inducing antiviral enzymes, type I IFNs modulate DCs, B cells, and T cells function in a direct or indirect manner (Trinchieri, 2010). However, in response to bacterial infection, type I IFNs promote anti-inflammatory responses through suppressing IL-1 production and macrophages (Mayer-Barber et al., 2011).

Induction of antimicrobial peptides by TLR signalling was first reported in drosophila (Tauszig et al., 2000). hBD1-4 (Ryu et al., 2013), inducible nitric oxide synthase (He et al., 2006; Lewis et al., 2011), LL-37 (Rivas-Santiago et al., 2008), and SPLI (Vroling et al., 2011) are inducible in response to TLR agonists.

#### 1.2.2.5 Negative regulation of TLR signalling

Activation of TLR signalling is a key event in innate immunity. However, uncontrolled TLR-mediated pro-inflammatory responses may cause immunopathology and autoimmune disease. For example, expression of TLRs and their responsiveness are increased in macrophages and peripheral blood monocytes from individuals with rheumatoid arthritis, systemic lupus erythematosus, or type 1 and 2 diabetes (Huang and Pope, 2009; Dasu et al., 2010; Devaraj et al., 2008; O'Gorman et al., 2015). Hence, it is necessary to have negative regulation of TLR-mediated responses, at multiple stages including soluble decoy TLR, binding sites overlapping, suppression of signal transduction, and inhibition of transcription (Figure 1.5). The multi-layered negative regulation indicates two possible explanations: first, each negative regulator is indispensable but insufficient to finely tune TLR signalling; second, negative regulators function in different tissue, or at different times (Liew et al., 2005).

Competition of binding sites happens both extracellularly and intracellularly. Extracellular soluble TLR2 and TLR4 have been found in human body fluids and the release is upregulated in response to PAMPs (Iwami et al., 2000; LeBouder et al., 2003; Oever et al., 2014). The expression of soluble TLR requires metalloproteinase that causes the shedding of ectodomain of TLR2 (Langjahr et al., 2014). By competitively binding to TLR2 ligand peptidoglycan as a decoy receptor, soluble TLR2 dampers activation of NF- $\kappa$ B and production of pro-inflammatory cytokines (Iwaki et al., 2002). Furthermore, soluble TLR2 is a biomarker to discriminate inflammatory diseases caused by infectious and non-infectious agents (Oever et al., 2014). The immunomodulatory effect of soluble TLR2 is detailed by Henrick et al. (2016).


Furthermore, TLR signalling can be blocked intracellularly by molecules containing the TIR domain, such as B-cell adaptor for PI3K (BCAP), single immunoglobulin IL-1R-related molecule (SIGIRR), and ST2. By TIR-TIR interaction, BCAP binds to TLR adaptor proteins; phosphorylated BCAP recruits PI3K to cytoplasmic membrane and initiates PI3K/Akt activation, thereby bridging TLR to PI3K/Akt signalling (Ni et al., 2012). There might be 3 mechanisms involved in anti-inflammatory regulation of BCAP/PI3K. First, activated Akt phosphorylation of glycogen synthase kinase (GSK)-3 $\beta$  contributes to production of the anti-inflammatory cytokine IL-10 (Zhang et al., 2016). Furthermore, the transcription factor forkhead box protein O (Foxo) 1 is known to enhance production of pro-inflammatory cytokines in macrophages and airway epithelial cells (Fan et al., 2010a; Seiler et al., 2013). Activation of PI3K/Akt leads to phosphorylation and nucleus exclusion of Foxo1 (Brunet et al., 1999). Lastly, a protein-protein interaction study showed that the association between BCAP and MyD88 can interfere with TIR-domain signalosome and MyD88-IRAKs interaction (Guyen-Maiorov et al., 2015). Additionally, PI3K/Akt promotes the pro-inflammatory response by phosphorylating IKK (Bai et al., 2009; Lin et al., 2011); or by phosphorylating GSK-3 $\beta$ , the inhibitor of NF- $\kappa$ B (Cremer et al., 2011). The other two TLR negative regulators SIGIRR and ST2 are transmembrane receptors containing a TIR domain, and hence interfere with TLR- and IL-1R-mediated inflammatory responses, with the exception of ST2 which only disrupts MyD88-dependent signalling (Wald et al., 2003; Brint et al., 2004; Nold-Petry et al., 2015).

Inhibition of signal transduction occurs at multiple stages. IRAK-M is a catalytically inactive kinase expressed in macrophages and monocytes that interferes with TLR signalling at different steps (Kobayashi et al., 2002). Upon stimulation with TLR ligands, IRAK-M blocks dissociation of IRAK1/4 from MyD88, and consequently the formation of IRAKs and TRAF6 (Kobayashi et al., 2002). Nakayama et al. (2004) reported that induction of tolerance by pretreatment with peptidoglycan in macrophages is associated with IRAK-M.

IRAK-M decreases IRAK1 kinase activity, as well as blocking recruitment of IRAK1 to MyD88.

Ubiquitinases and deubiquitinases are associated with post-translational modification in TLR signalling. SOCS1 and SOCS3 have ubiquitin ligase activity (Hovsepian et al., 2013; Yu et al., 2015). Hence, by polyubiquitination, SOCS1 and SOCS3 lead to proteasomal degradation of their targets, including Mal (Mansell et al., 2006), TRAF3/6 (Rothlin et al., 2007; Zhou et al., 2015), and p65 NF- $\kappa$ B (Strebovsky et al., 2011). The anti-inflammatory role by SOCSs is reviewed by Inagaki-Ohara (2014). Another ubiquitin ligase Triad3A causes proteolytic degradation of TLR4 and TLR9 (Chuang and Ulevitch, 2004), as well as TLR adaptors TRIF and Mal (Bibeau-Poirier and Servant, 2008). Overall, ubiquitinases downregulate the level of signal proteins, and thus disrupt or abolish TLR signal transduction. Ubiquitination of TRAF6 is an important step to activate TRAF6 for recruiting and activating TABs/TAK1 complex. Polyubiquitinated TRAF6 can be a substrate for deubiquitinases, such as A20, cylindromatosis D (CYLD), and deubiquitinating enzyme A (DUBA, deubiquitinates TRAF3) (Kayagaki et al., 2007; Bibeau-Poirier and Servant, 2008; Sun, 2010). For example, by removing the ubiquitin chain from TRAF6 and disassociating TRAF6 and Ubc13/UbcH5c, A20 inhibits LPS-induced NF- $\kappa$ B pathway (Shembade et al., 2010).

Detailed TLR endogenous inhibitors are summarised in Liew et al., (2005) and Kondo et al., (2012a). Taken together, negative regulators of TLR signalling contribute to immune homeostasis, protecting the host from overreactive inflammation and yet ensuring sufficient host defence. Individual negative regulators have unique and non-redundant roles as deletion of any individual regulators results in hyperresponsiveness. Although there is overlapping of target proteins, regulators possibly work in a synergistic manner to induce sufficient inhibition. In addition to MyD88-dependent TLR signalling, regulators such as Triad3A and DUBA act via a MyD88-independent pathway. Notably, as TRAFs are adaptor proteins in TNFR family, inhibition of TRAF3 or TRAF6 suppresses TNFR signalling simultaneously (Kondo et al., 2012b). Hence, the pan-inhibition activity of negative regulators connects multiple pro-inflammatory pathways and have a stronger inhibitory effect. Artificial inhibitors of TLR signalling have great potential to treat chronic inflammation and autoimmune diseases; however, so far no TLR antagonists have been approved for clinical treatment (Gao et al., 2017). Furthermore, pathogens have evolved to disturb TLR signalling by producing multiple enzymes, such as kinase, protease, ubiquitinases, and deubiquitinases (McGuire and Arthur, 2015). Thus, it is favourable for pathogens to invade and colonise without initiating host immune responses.



**Figure 1.5 Negative regulators of MyD88-dependent TLR signalling pathways.**

Negative regulators (in red boxes) play nonredundant roles at distinct steps of TLR signalling. Generated from shedding of TLR ectodomains, sTLRs competitively bind to PAMPs as extracellular decoy receptors. Molecules with TIR domain (BCAP, SIGIRR, and ST2) interfere with TLR/MyD88/Mal complex through competing the binding sites. BCAP also activates PI3K which leads to production of IL-10 and inactivation of transcription factor Foxo1. IRAK-M dampens disassociation of IRAKs and MyD88. Ubiquitinases (SOCS1/3 and Triad3A) and deubiquitinases (A20 and CYLD) disrupt posttranslational modification and thus block signal transduction (own original artwork).

### 1.3 Hypothesis and aims

The introduction above has described host defences and the multiple mechanisms that pathogens have evolved to circumvent these responses.

Bacterial infections in the LRTI mainly exist in polymicrobial communities. Microbial interactions within airway microbiome may alter disease course and clinical outcomes. While Bcc colonisation negatively affects the survival of CF individuals, *S. aureus* infection has the opposite effect and may delay lung disease progression. The underlying mechanism remains largely unknown. It was hypothesised that *S. aureus* interacts with the host defence by manipulating innate signalling pathways, thereby alleviating pathogenesis caused by Bcc virulence. To test this hypothesis, human airway epithelial cells were challenged with *S. aureus* and Bcc single-species infection or co-infection. Immune responses by epithelial cells were recorded and compared.

#### Aims:

- To assess the modulation of single-species *S. aureus* and Bcc on host signalling cascade and inflammatory responses
- To study the effect of *S. aureus* / *B. cenocepacia* co-infection on host signalling and inflammatory responses
- To interpret the host target of Staphylococcal immunomodulatory factors
- To identify the nature of immunomodulatory molecules released by *S. aureus*

## **Chapter 2 Material and Methods**

## 2.1 Human Airway Epithelial Cell Lines

Human bronchial epithelial 16HBE14o (16HBE, obtained from Prof Dieter C. Gruenert, University of California San Francisco, US) was isolated from surface epithelium of 2<sup>nd</sup> division bronchi of a 1-year old male heart-lung transplant patient and transformed by simian vacuolating virus 40 large T-antigen (Cozens et al., 1994). 16HBE cells retain tight junction, cilia structure, and cystic fibrosis transmembrane conductance regulator expression (Cozens et al., 1994). Human airway epithelial Calu-3 was purchased from ATCC (HTB-55). Isolated from the pleural effusion of a 25-year old male with a pulmonary adenocarcinoma, Calu-3 cells display epithelial morphology, barrier integrity and cystic fibrosis transmembrane conductance regulator under defined conditions (Fogh et al., 1977; Kreft et al., 2015).

16HBE cells and Calu-3 cells were maintained in complete medium containing Dulbecco's Modified Eagle Medium (DMEM), supplemented with 10% (v/v) heat inactivated fetal bovine serum (FBS), 100 units/mL penicillin and 100 µg/mL streptomycin (Gibco). Cells were incubated in 75 cm<sup>2</sup> flasks (Nunc) at 37°C in a humidified atmosphere with 5% CO<sub>2</sub>.

For dissociation, cells were rinsed with 5 mL Dulbecco's phosphate buffered saline (DPBS), followed by incubating with 5 mL 0.25% trypsin-EDTA (Gibco) for 10 minutes at 37°C. After adding 5 mL complete medium to neutralise trypsin, the number of viable cells were determined using trypan blue exclusion. Briefly, 50 µL cell suspension was mixed with same volume trypan blue (0.4%), and loaded on a haemocytometer. Viable cells in the four large squares were counted and cell density (cells/mL) was calculated as: number of viable cells  $\times 2 \times 10^4 / 4$ . After centrifugation at 1000 *g* for 5 minutes, cells were resuspended in complete medium and plated out at a density of  $5 \times 10^5$  cells/well into 12-well plates or  $2.5 \times 10^5$  cells/well into 24-well plates.

For storage, cell suspensions were adjusted to  $4 \times 10^6$  cells/mL in freeze medium (50% DMEM, 40% FBS, 10% DMSO, v/v), transferred into cryovials and stored in liquid nitrogen.

## 2.2 Bacteria Strains

The bacterial strains used in this study are listed in Table 2.1. Bacterial cultures were inoculated on Luria-Bertani agar (LB, 1% tryptone, 0.5% yeast extract, 1% NaCl, 1.5% agar-agar, w/v) from -80°C stocks every week, incubated at 37°C for 1 day and kept at 4°C. Due to its slow growing nature, *Burkholderia* strains were incubated at 37°C for 2 days and maintained at room temperature. A single colony of the strain was transferred to LB broth at 37°C with shaking for about 16 hours. Overnight cultures were diluted at 1



in 20 and cultured until exponential phase as determined by the growth curve. Bacteria were centrifuged at 5000 *g* for 10 minutes and washed once with sterile PBS (0.01 M phosphate buffer, 0.0027 M KCl and 0.137 M NaCl). The bacterial cell density was adjusted based on optical density (OD) measured at a wavelength of 600 nm to meet different multiplicity of infection (MOI, the ratio of bacteria to cells) values. When tryptone soya broth (TSB) was used, bacteria were cultured in TSB (1.7% pancreatic digest of casein, 0.3% enzymatic digest of soya bean, 0.5% NaCl, 0.25% K<sub>2</sub>HPO<sub>4</sub>, 0.25% glucose) at 37°C with shaking for 16 hours.

Heat-killed bacteria were prepared by adjusting the OD<sub>600</sub> of cultures to the desired MOI and heating these on a heating block at 100°C for 1 hour. Bacterial supernatants were prepared by centrifuging overnight cultures at 5000 *g* for 10 minutes and filtered through a 0.2 µm membrane. Heat-inactivated supernatants were prepared by heating samples at 95 °C for 10 minutes. To fractionate excreted products from *S. aureus*, supernatants were submitted to 10-kDa- and 3-kDa-cutoff centrifugal concentrators (Vivaspin 20, Sartorius), centrifuged at 5000 *g* for 17 minutes and 180 minutes respectively following the manufacturer's instructions. Supernatants were separated into three fractions: >10000 molecular weight cut-off (MWCO), 3000-10000 MWCO, and <3000 MWCO. Filtrates and concentrates were adjusted to the original volume with LB, followed by filter sterilisation. Immobilised proteinase K (PK) was reconstituted in Tris-Cl (50 mM, pH 8) / CaCl<sub>2</sub> (10 mM) and used at a concentration of 100 µg/mL at 37°C for 1 hour. To prepare inactivated PK, PK was heated at 99°C for 1 hour. Immobilised PK mixture was subsequently removed by centrifugation at 250 *g* for 10 minutes. All bacterial products were aliquoted and stored at -80°C.

For long-term storage of bacterial strains, overnight cultures were transferred to a cryovial with sterile glycerol (final glycerol concentration 25%, v/v) to limit cellular damage, vortexed and stored at -80°C.

**Table 2.1 Bacterial strains used in this study.**

Strain	Characteristics	Source	Reference
<b><i>S. aureus</i></b>			
MRSA252	Hospital-acquired MRSA, fatal post-op septicemia	E.J.Feil	Holden et al., 2004
MSSA NCTC 6571	Methicillin-sensitive <i>S. aureus</i>	NCTC	
MR31	Clinical isolate	R.C. Massey	Trzciński et al., 1997

**Table 2.1 (continued)**

MR81	Clinical isolate	R.C. Massey	Trzciński et al., 1997
MRSA USA400	Community-associated <i>S. aureus</i>	R.C. Massey	Voyich et al., 2006
MRSA41		R.C. Massey	Collins et al., 2010
MSSA C316		R.C. Massey	Robinson et al., 2005
MSSA NCTC 10788		NCTC	
MSSA Newman	Isolated from a secondary osteomyelitis infection of a TB patient	R.C. Massey	Baba et al., 2008
MSSA RN4220	Derivative of NCTC8325-4, with <i>agrA</i> mutation	R.C. Massey	Nair et al., 2011
MSSA SH1000	Laboratory strain, derivative of NCTC8325 by complementation of <i>rsbU</i>	R.C. Massey	O'Neill, 2010
MSSA160		R.C. Massey	Feil et al., 2003
MSSA209		R.C. Massey	Collins et al., 2010
US119		R.C. Massey	
<b>Bcc</b>			
<i>B. multivorans</i> DSMZ 13243	Isolated from a CF patient	DSMZ	
<i>B. cenocepacia</i> J2315	Isolated from a CF patient, epidemic ET12 lineage	J. R. W. Govan	Holden et al., 2009
<i>B. anthina</i> J2552	Botanical isolate	J. R. W. Govan	Palfreyman et al., 1997
<b>Reference strains</b>			
<i>E. coli</i> HB101	Non-invasive K-12 strain		Boyer and Roulland-Dussoix, 1969
<i>Salmonella</i> Typhimurium NTB6		Radboud collection	Boleij et al., 2011b

## 2.3 Biofilm Assay

Biofilm formation was assessed as described before (Coffey and Anderson, 2014). Briefly, overnight cultures were diluted 1:100 with LB, and 100  $\mu$ L aliquots were added to a 96-well microtiter plate. After 24 hours incubation at 37°C on a 3-dimensional rotator, planktonic bacteria were removed, and the biofilms were gently rinsed with PBS twice. Biofilms were stained with 125  $\mu$ L of 0.1% (w/v) crystal violet for 10 minutes. Excess crystal violet was removed by washing the plate three times in trays filled with lukewarm tapwater, after which the biofilm was dried in a 60°C oven. Crystal violet was dissolved with 150  $\mu$ L of 20% (v/v) acetic acid and 125  $\mu$ L of each sample was transferred to a new plate. The absorption was read at a wavelength of 550 nm.


## 2.4 Measurement of Epithelial Barrier Integrity

### 2.4.1 Measurement of Transepithelial Electric Resistance (TEER)

Calu-3 cells were plated into Transwell permeable supports (12 mm diameter, Corning) at  $5 \times 10^5$  cells/mL, 0.3 mL per insert. The basolateral chamber was supplied with 1 mL complete medium. Medium in the apical chamber was removed 48 hours after seeding to induce the formation of polarised monolayer with tight junction.

TEER was monitored by EVOM Volt/Ohm Meter and STX2 electrode (World Precision Instrument). STX2 electrode was sterilised in 100% ethanol and washed once with DMEM. As shown in the schematic (Figure 2.1), serum-free DMEM (300  $\mu$ L) was added to the apical chamber. After 30 minutes incubation, the long electrode was placed into the basolateral chamber and the short electrode in the apical chamber without damaging the monolayer. After subtracting the readout of inserts with medium (background) from raw data, they were normalised for area, giving TEER values in  $\text{Ohm.cm}^2$  as in the following equation:  $(\text{total resistance} - \text{background resistance}) \times 1.12\text{cm}^2$ . Higher value of resistance indicates higher confluency of cells on the inserts as less ions can transport through the membrane. Calu-3 cells achieved a polarised state within 7 days when the TEER reached  $>400 \text{ ohm.cm}^2$ .

Polarised Calu-3 cells were infected apically with MRSA252 and *B. multivorans* 13243 (0.5 mL) for 24 hours. TEER was recorded at time 0, 3 and 24 hours post infection.



**Figure 2.1 Schematic of measuring TEER for airway epithelial cell monolayer.**

STX2 electrode is used to measure the TEER. The total readout includes the electrical resistance of the cell monolayer, the membrane and culture medium.

#### 2.4.2 Flux of Fluorescein Isothiocyanate (FITC) Labelled-Dextran

Calu-3 cells were infected with MRSA252 and *B. multivorans* 13243 as described above. At 24 hours postinfection, the inserts were washed twice with Hanks' Balanced Salt Solution (HBSS) gently to remove phenol red. Two-hundred-microlitre FITC-dextran (molecular weight 3000-5000, 500 µg/mL in HBSS) were applied to the apical side and 600 µL HBSS to the basolateral side. After 30 minutes incubation at 37 °C, medium from the basolateral chamber was collected and fluorescence intensity was determined using a FLUOstar OPTIMA plate reader (BMG Labtech) at 485 nm excitation wavelength and 530 nm emission wavelength. The readout was normalised by subtracting the background (HBSS), and was divided by maximum (the insert without monolayer) to obtain the value of percent epithelial permeability.

## 2.5 Invasion Assay

To study the role of PI3K/Akt signalling, epithelial cells were pretreated with LY294002 or wortmannin for 30 or 60 minutes until the end of invasion period. 16HBE cells were infected with MRSA252, MSSA NCTC 6571, *B. multivorans* 13243, *B. cenocepacia* J2315, *B. anthina* J2552, *E. coli* HB101 or *S. Typhimurium* NTB6 at MOI 50 for 2 hours. The number of bacteria in the inocula were determined by viable count. 16HBE cells were rinsed three times with sterile DPBS to remove unattached pathogens and lysed with 100 µL saponin (1% w/v in trypsin-EDTA). The lysates were plated on LB agar plates to count the number of bacteria that were attached to the epithelial cell membrane. Alternatively, following 2 hours infection and DPBS rinse, medium was supplemented with antibiotics (Table 2.2) to eliminate extracellular bacteria for the time indicated in Table 2.2. To ensure more than 99.9% bacterial killing effect, the last DPBS wash was plated on LB agar plates as well. Subsequently, epithelial cells were lysed with 100 µL saponin and internalised

bacteria were determined by viable count. Adherence is shown as a percentage of the inoculum, and internalisation as a percentage of adherence.

**Table 2.2 Antibiotics treatment for bacterial strains.**

Strains	Antibiotics	Time
<i>S. aureus</i> isolates	Lysostaphin (20 µg/mL) + vancomycin (25 µg/mL)	1 h
<i>B. multivorans</i> 13243	Ceftazidime (1 mg/mL) + amikacin (1 mg/mL)	2 h
<i>B. anthina</i> J2552		
<i>B. cenocepacia</i> J2315	Ciproflaxin (1 mg/mL) + amikacin (1 mg/mL)	2 h
<i>E. coli</i> HB101	Gentamicin (200 µg/mL)	1 h
<i>S. Typhimurium</i> NTB6	Gentamicin (200 µg/mL) + ampicillin (50 µg/mL)	1 h

## 2.6 Western Blot Analysis

After stimulation, 16HBE or Calu-3 cells were rinsed with cold DPBS twice and lysed with 100 µL RIPA buffer (1% Triton X-100, 1% sodium deoxycholate, 25 mM Tris-Cl pH 8.0, 150 mM NaCl), supplemented with PhosSTOP (Roche) and protease inhibitor cocktail (2 mM AEBSF, 0.3 µM aprotinin, 116 µM bestatin, 14 µM E-64, 1 µM leupeptin 1 mM EDTA; Sigma) just before use. Lysates were collected, centrifuged at 13 000 *g* for 10 minutes at 0°C and the protein concentrations were quantified using Pierce™ BCA Protein Assay Kit (Thermo Scientific) according to the manufacturer's instructions. Briefly, 10 µL of sample was diluted 1 in 10 with Milli-Q water, and 25 µL of diluted sample was pipetted into a 96-well microtiter plate in triplicate and mixed with 200 µL Working Reagent (0.08% cupric sulfate w/v). Standards ranged from 25 to 2000 µg/mL. The plate was covered and incubated at 37°C for 30 minutes. The absorbance was read at a wavelength of 550 nm.

Protein samples with equivalent total protein concentrations were mixed with 5 × loading buffer (250 mM Tris-Cl pH 6.8, 500 mM dithiothreitol, 10% SDS, 0.25% bromophenol blue, 50% glycerol, 2% β-mercaptoethanol added before use, v/v). After 95°C heat-treated for 5 minutes, aliquots were resolved on a 10% SDS-PAGE gel at 90 volts for 30 minutes and then increased to 110 volts for about 1 hour (Mini-Protein Tetra Cell System, Bio-Rad). Proteins were transferred to a nitrocellulose membrane at 150 mA for 1.5 hours (Mini Trans-Blot Cell, Bio-Rad). Running buffer is composed of 190 mM glycine, 25 mM Tris base, 0.1% w/v SDS, and transfer buffer is composed of all these chemicals supplemented with 20% v/v methanol.

To minimise nonspecific binding of the primary antibodies, the membrane was blocked with 5% (w/v) skimmed-milk in TBS (50 mM Tris-Cl, 150 mM NaCl, pH 7.5) for 1 hour at room temperature. Membranes were probed with primary antibodies anti phospho-Akt, phospho-GSK-3 $\beta$ , phospho-NF- $\kappa$ B-p65, phospho-p38, phospho-Erk, phospho-I $\kappa$ B $\alpha$ , or I $\kappa$ B $\alpha$  (Cell Signaling Technology), respectively at 4°C with gentle rotation overnight. For anti phospho-I $\kappa$ B $\alpha$  and I $\kappa$ B $\alpha$ , same amount of proteins was loaded twice on separate gels and probed separately. After washing with 3 $\times$ TBST (TBS-0.1% v/v Tween-20) and TBS (each wash 5 minutes), the membranes were then probed with the secondary antibodies, IRDye 800 donkey anti-mouse IgG or 680 donkey anti-rabbit IgG (1:10000 in 5% BSA-TBST) for 1 hour. The blots were washed with 3  $\times$  TBST and TBS. Quantitative analysis of the signals from each antibody was performed using the Li-Cor Odyssey Clx imaging system. The membranes were subsequently probed with primary antibodies anti Akt, p38, or Erk, respectively (Cell Signaling Technology), and secondary antibodies as previous. Signal intensity was normalised as a percentage of total Akt, p38 or I $\kappa$ B $\alpha$  on the same membrane.

## **2.7 Detection of Cytokine Production**

At the end of stimulation, conditioned medium of 16HBE cells was centrifuged at 5000 *g* for 10 minutes and supernatants were analysed for IL-8 using a commercially available ELISA kit (eBioscience) according to the manufacturer's instructions. Briefly, Maxisorp ELISA plates (Nunc) were coated with 100  $\mu$ L capture antibody at 4°C overnight. The wells were washed three times with Wash Buffer (0.05% v/v Tween-20, PBS) and blocked with 200  $\mu$ L diluent for 1 hour at room temperature. For each assay, human IL-8 standard was reconstituted using 1 mL distilled water and diluted with ELISA Diluent to 250 pg/mL. The standard curve included 8 points by performing serial 2-fold dilutions of the top standard (range, 2-250 pg/mL). One-hundred microliter standards, samples, and ELISA Diluent (blank) were added to the plate in duplicate. After 2 hours incubation, the wells were washed as previous. The wells were incubated with 100  $\mu$ L detection antibody for 1 hour, followed by 3 washes. After 30 minutes incubation with 100  $\mu$ L Avidin-HRP, the wells were rinsed 5 times. One-hundred microliter TMB Solution was added to the wells until 15 minutes later the reaction was stopped by adding 50  $\mu$ L Stop Solution (2N H<sub>2</sub>SO<sub>4</sub>). The optical density was read at 450 nm and 550 nm, using a FLUOstar OPTIMA (BMG Labtech) microplate reader. The values of 550 nm were subtracted from those of 450 nm; the values of the blank wells were subtracted from those of standards and samples. Standard curve was generated by graphing concentration of IL-8 versus optical density and used to determine the concentration of IL-8 in the samples.

## 2.8 Cell Viability Assay

To confirm that the modulation of IL-8 production by heat-killed *S. aureus* or the supernatants was not due to cell damage, epithelial metabolic activity was assessed using the MTT assay (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide). 16HBE cells were plated in 96-well plates (5000 cells/well) and incubated for 24 hours. At 24 hours post-stimulation with 10% (v/v) *S. aureus* supernatants or heat-killed *S. aureus*, cells were cultured with 12 mM MTT-DMEM for 2 hours. Formazan formed from MTT in viable cells was dissolved in DMSO for 10 minutes before the optical density was read at 550 nm wavelength.

To determine cell damage caused by live bacteria, 16HBE cells were stimulated with *S. aureus* strains (MOI 0.5) and *B. cenocepacia* J2315 (MOI 5). 16HBE were seeded at  $2 \times 10^4$  cells/well in 96-well plates one day before experiments (0.1 mL DPBS into the peripheral wells of the 96-well plates). At 2 hours postinfection, cells were rinsed with DPBS twice and then incubated for 24 hours in 0.1 mL phenol red-free DMEM to minimise background absorbance. Lactate dehydrogenase (LDH) release was measured with an LDH Cytotoxicity Assay Kit according to manufacturer's instructions (Pierce). Briefly, 45 minutes before collecting supernatants, 10  $\mu$ L sterile water was added to one set of triplicate wells as spontaneous LDH activity control; 10  $\mu$ L Lysis Buffer was added to another set of triplicate wells as maximum LDH activity control. After incubated at 37°C for 45 minutes, 50  $\mu$ L supernatants were mixed with 50  $\mu$ L reaction mixture in a new 96-well plate and incubated at room temperature for 30 minutes in the dark. The reaction was stopped by adding 50  $\mu$ L Stop Solution. The absorbance was read at 450 nm and 600 nm, using a FLUOstar OPTIMA (BMG Labtech) microplate reader. The 600 nm absorbance value was subtracted from the 450 nm absorbance value. Background control value (DMEM with no cell) was subtracted from average of each sets of readouts and calculated %Cytotoxicity using the following equation.

$$\% \text{Cytotoxicity} = 100 \times (\text{samples value} - \text{spontaneous LDH activity control}) / \text{maximum LDH activity control}$$

## 2.9 Statistical Analysis

Data were expressed as the mean  $\pm$  standard error of the mean (SEM)  $n \geq 3$  independent biological repeats. Results were compared using one-way or two-way ANOVA, followed by Tukey or Dunnet multiple comparisons test, with  $p < 0.05$  considered significant. The Pearson correlation score with two-tailed test of significance was used for rank correlations.

## 2.10 Reagents and Consumables

**Table 2.3 Reagents and consumables for mammalian cell and bacteria culture.**

Product	Supplier	Catalog number
<b><i>Reagents and consumables for cell culture</i></b>		
12-well clear sterile plate	Thermo, UK	150628
24-well clear sterile plate	Corning, UK	3527
96-well black plate	Greiner Bio-one, UK	655087
96-well clear sterile plate	Corning, UK	CLS3595
DMEM	Gibco, UK	41966052
DPBS	Gibco, UK	14190250
FBS	Gibco, UK	10082147
HBSS	Gibco, UK	14025092
Penicillin-Streptomycin	Gibco, UK	15140122
T75 cell culture flasks	Nunc, UK	156499
Transwell cell culture inserts	Sigma, UK	CLS3460
Trypsin-EDTA (0.25%)	Gibco, UK	25200072
Tweezer electrode	World Precision Instrument	STX2
<b><i>Reagents and consumables for bacterial cultures</i></b>		
Agar-agar	Fisher, UK	10548030
Immobilised proteinase K	Sigma, UK	82452
LB broth, Miller	Fisher, UK	BP1426
PBS tablets	Sigma, UK	P4417
Tryptone soya broth	Oxoid, UK	CM0129
Vivaspin, 10000 MWCO	Sartorius, UK	VS2001
Vivaspin, 3000 MWCO	Sartorius, UK	VS2091



**Table 2.4 Drugs for *in vitro* treatment.**

Product	Supplier	Catalog Number
Flagellin, FLA-ST	Invivogen, UK	tlrl-stfla
hTLR5-Fc	Invivogen, UK	fc-htlr5
Lipopolysaccharides	Sigma, UK	L2880
LPS-RS	Invivogen, UK	tlrl-rslps
LY294002	Sigma, UK	L9908
Pam3CSK4	Invivogen, UK	tlrl-pms
Recombinant human IL-1 $\beta$	Peprotech, UK	200-01B
Wortmannin	Sigma, UK	W1628

**Table 2.5 Reagents for western blot.**

Product	Supplier	Catalog Number
2-mercaptoethanol	Sigma, UK	M3148
30% acrylamide/bis	Bio-rad, UK	1610154
Ammonium persulfate	Fisher, UK	10744171
Glycine	Fisher, UK	10080160
Prestained protein standards	Bio-rad, UK	1610375
Protease inhibitor cocktail	Sigma, UK	P2714
Roche PhosSTOP tablets	Sigma, UK	4906845001
SDS	Sigma, UK	L3771
TEMED	Fisher, UK	10549960
Tris base	Sigma, UK	252859

**Table 2.6 List of primary antibodies.**

Antigen target	Product Number <sup>a</sup>	Dilution	Diluent	Host
Akt	2920	1:2000	5% skimmed-milk-TBST	Mouse
Erk1/2	4695	1:1000	5% BSA-TBST	Rabbit
IκBα	4814	1:1000	5% skimmed-milk-TBST	Mouse
p38	8690	1:1000	5% BSA-TBST	Rabbit
Phospho-Akt (S473)	4060	1:2000	5% BSA-TBST	Rabbit
Phospho-Akt (T308)	13038	1:1000	5% skimmed-milk-TBST	Rabbit
Phospho-Erk1/2	9106	1:2000	5% skimmed-milk-TBST	Mouse
Phospho-GSK-3β	5558	1:1000	5% BSA-TBST	Rabbit
Phospho-IκBα	9246	1:1000	5% skimmed-milk-TBST	Mouse
Phospho-p38 (T180/Y182)	9216	1:2000	5% skimmed-milk-TBST	Mouse
Phospho-p65	3033	1:1000	5% BSA-TBST	Rabbit

<sup>a</sup> All the products were purchased from Cell Signaling Technology, MA, US

**Table 2.7 List of secondary antibodies.**

Product	Product Number	Supplier
IRDye 800CW donkey anti-mouse	92632212	Li-Cor, UK
IRDye 680RD donkey anti-rabbit	92668073	Li-Cor, UK

**Table 2.8 Assay kits.**

Product	Product Number	Supplier
IL-8 human ELISA kit	eBioscience, US	88-8086-88
Pierce BCA protein assay kit	Thermo, UK	23225
Pierce LDH cytotoxicity assay kit	Fisher, UK	13454269

# **Chapter 3 Investigating Host Response to Bacterial Infections in Human Airway Epithelial Cells**

### 3.1 Introduction

The lung is the major organ of the respiratory system in humans. It is exposed to inhaled microorganisms and particles continuously. To resist infectious attack, the lung has developed sophisticated defence mechanisms. The physical barrier which comprises of airway epithelial cells and the mucus layer plays a major role in limiting entry and assisting clearance of pathogens. The coordinated beating of the cilia on the apical surface of epithelial cells enables removal of particles that are trapped by mucus (Bustamante-Marin and Ostrowski, 2017). Paracellular junctional complexes play an essential role in controlling the transport of water, ions and immune cells, and separating airborne pathogens from the interior of the body to maintain homeostasis in the lung (Brune et al., 2015). However, the epithelial barrier is compromised when the traversal of pathogens occurs (Kim et al., 2005; Malik et al., 2015). To study the barrier function, cells can be cultured on a porous membrane, which is the most common method to represent the morphology and tight junctions of airway epithelial cells *in vitro* (Grainger et al., 2006). In this it is important that the cells used do polarise and form tight junctions as cells would normally do *in vivo*; not many pulmonary cell lines do, but the Calu-3 cell line is one of the few exceptions (Shen et al., 1994). When a confluent monolayer is achieved, the apical side is separated from the basolateral membrane and therefore displays a certain TEER value, which is a measure of the integrity of airway epithelial. This integrity may be compromised by bacteria, but so far little work has been done on effect of MRSA and Bcc, two major lung infection species.

Airway epithelial cells also serve as a sentinel to detect intruding pathogens through a wide range of PRRs (Takeuchi and Akira, 2010). Activation of PRRs lead to a sequence of events, including activation of downstream signalling pathways, initiation of the expression of proinflammatory cytokines, and recruitment and activation of immune cells to the site of infection. Eventually pathogens are resolved by phagocytosis, or antimicrobial substances released by professional phagocytes if the microbial clearance is successful (Nagl et al., 2002). In this process, IL-8 plays an important role at the early stage of infection by chemoattracting granulocytes such as neutrophils. Hence, the level of IL-8 is a biomarker of pro-inflammatory responses and inflammation.

The PI3K/Akt pathway is one of the downstream signalling cascades activated by TLRs, regulating a wide range of cellular functions, such as survival, proliferation and the inflammatory response (Hawkins and Stephens, 2015; Rosadini and Kagan, 2017). For example, Akt is required for the proinflammatory response to *B. cenocepacia* in human peripheral blood mononuclear cells (PBMC) using a PI3K inhibitor (Cremer et al., 2011).

In contrast, inhibition of PI3K/Akt signalling abolished anti-inflammatory cytokine IL-10 production but did not affect proinflammatory cytokine TNF $\alpha$  production by PBMCs in response to *S. aureus* isolates (Peres et al., 2015). Therefore, the discrepancy of PI3K/Akt signalling might indicate that it has pleiotropic roles in regulating inflammatory response, depending on the stimulus and other signals generated.

*S. aureus* and Bcc are the leading cause of lower respiratory tract infections. Despite being a commensal microbe in the respiratory tract, *S. aureus* is able to colonise the host in a pathogenic manner, which leads to pneumonia with an intense and exaggerated host response. The major challenge of *S. aureus* infection is rapidly evolved antibiotic resistance (Foster, 2017). After acquisition of the resistance gene *SCCmec*, a subgroup of *S. aureus* becomes resistant to most  $\beta$ -lactam antibiotics, namely MRSA. The incidence of individuals with CF harbouring MRSA increased from 5% to 26% from 2000 to 2010, and this has plateaued ever since (Cystic Fibrosis Foundation, 2017). On the other hand, the prevalence of Bcc is about 3% among persons with CF. The acquisition of *Burkholderia* infections are most likely from the natural environment such as soil, and some epidemic strains are well adapted to infectious disease and have greater ability for interpatient transmission (Govan et al., 1993; LiPuma et al., 2002). While some patients have no deterioration, about 20% of infected patients develop cepacia syndrome either soon after the first positive isolate or many years after (Blackburn et al., 2004). Cepacia syndrome is a fatal illness characterised as rapid deterioration of pulmonary function, bacteremia and sepsis. Pre-operative Bcc colonisation is also a significant marker for post lung transplantation mortality (De Soyza et al., 2010). The difficulty of treating Bcc is recognised with their intrinsic and acquired antibiotic resistance. Multi-antibiotic resistance is observed in *Burkholderia* species due to the expression of  $\beta$ -lactamase, efflux pump and outer membrane permeability barrier (Fraser-Pitt et al., 2016; Rhodes and Schweizer, 2016).

In chronic lung infections, microbial pathogens tend to form so-called biofilms to promote long-term survival under the stress coming from antibiotics treatment and the immune clearance (Archer et al., 2011). These biofilms are sessile communities of bacteria embedded within a self-produced matrix on biotic or abiotic surfaces. Biofilm extracellular polymeric substances comprise polysaccharides, proteins, glycoproteins, glycolipids, and extracellular DNA. Bacteria in biofilm displays altered phenotypes with respect to gene expression, protein production and growth rate (Donlan and Costerton, 2002). By adapting to the sessile type of life, microorganisms in biofilm benefit from several advantages over their planktonic counterparts. The majority of cells in biofilm communities are dormant due to the lack of oxygen and nutrients. Hence, the low metabolic rate and

quiescent state downregulate the susceptibility to antimicrobial agents that inhibit synthesis in the process of cell division (Høiby et al., 2010). In addition, the diffusion barrier in biofilms plays a critical role to reduce the penetration of antimicrobial factors. Indeed, antimicrobial agents can be inactivated or absorbed by the matrix layer before reaching the inner layer of bacteria. For example, a study proved that the polysaccharide Poly- $\beta$ -1,6-N-acetyl-D-glucosamine in the biofilm matrix restricts fluid transport and sequesters the biocide cetylpyridinium chloride within biofilms of *Actinobacillus pleuropneumoniae* and *S. epidermidis* (Ganeshnarayan et al., 2009). In the final stage of the life cycle of a biofilm, bacteria can detach from the community and disseminate to uninfected sites. Biofilm dispersal is regulated by environmental factors, such as shear stress, and biological signalling. It is a vital step between the persistence and dissemination of bacterial pathogens (Guilhen et al., 2017). Overall, biofilm formation represents a key virulence factor of microorganisms.


This chapter was aimed at investigating the host response to bacterial pathogens in airway epithelial cell lines. First, the disruptive effect on *in vitro* epithelial monolayer model by *S. aureus* and *B. multivorans* were characterised. One of the major virulence factors, biofilm formation, was compared between *S. aureus* and Bcc. Finally, the role of PI3K in the production of the neutrophil chemoattractant IL-8 in response to *S. aureus* and Bcc was assessed by the use of PI3K inhibitors.

### **3.2 Airway Barrier Function Is Compromised by Bacterial Pathogens**

Bacterial invasion compromises the barrier integrity of the tight junctions and epithelial cell death. To assess the integrity of infected Calu-3 monolayers, TEER and FITC-dextran flux were monitored.

TEER levels did not change in the first 3 h infection either with MRSA252 or *B. multivorans* 13243, but a pronounced drop was achieved at 24 h. A difference was observed between MRSA252 and *B. multivorans*. At 24 h, a CFU-dependent decrease was monitored in MRSA252-treated groups, whereas *B. multivorans*-treated monolayers showed the same level of reduction at all dose levels examined (Figure 3.1A and B).

To further assess airway epithelial barrier compromise, the diffusion of FITC-dextran (4 kDa) from the apical to basolateral chamber was measured at 24 h postinfection. FITC-dextran detected from the basolateral chamber showed a dose-dependent increase induced by both strains, and *B. multivorans*-infected monolayers showed greater flux than MRSA252.




**Figure 3.1 Compromised Calu-3 barrier by MRSA252 or *B. multivorans* 13243 invasion.**

Polarised Calu-3 monolayers were exposed to increasing CFU of MRSA252 and *B. multivorans* 13243 (0.5 mL apically). **(A and B)** TEER levels were recorded at 3 and 24 h post infection. N=3 independent experiments, performed in duplicate. Error bars indicate mean  $\pm$  SEM. Data were analysed using two-way ANOVA with Dunnett's multiple comparisons test between untreated and infected monolayers. \*,  $p < 0.05$ . **(C and D)** Diffusion of FITC-dextran through compromised monolayers at 24 h. Data are presented as percent of maximum diffusion (inserts with no cells) and analysed by one-way ANOVA with Dunnett's multiple comparisons test. Compared to untreated monolayers. \*,  $p < 0.05$ .

### 3.3 Evaluation of Biofilm Formation by Bcc and *S. aureus* sp.

To assess biofilm formation of Bcc and *S. aureus* isolates, crystal violet staining was used (Figure 3.2). Cystic fibrosis-isolated strains *B. multivorans* 13243 (absorbance readings of  $1.35 \pm 0.26$ ) and *B. cenocepacia* J2315 ( $0.60 \pm 0.09$ ) formed abundant biofilms compared to the environmental isolate *B. anthina* J2552 ( $0.10 \pm 0.03$ ). *S. aureus* strains formed same levels of biofilm ( $0.92 \pm 0.42$ ) except MSSA SH1000 ( $2.80 \pm 0.23$ ), which formed 2 times more biofilm than the average biofilm formation. Notably, SH1000 and RN4220 share the same ancestral strain NCTC 8325, but that RN4220 only produced half the amount of biofilm as compared to SH1000. Strain RN4220 differs from SH1000 with defective *agrA* and *rsbU* genes, which are important virulence regulators (Traber and Novick, 2006; Bæk et al., 2013). Compared to RN4220, SH1000 has stronger haemolytic and proteolytic activity, and produces golden pigmentation (Bæk et al., 2013). Overall, heterogeneity of biofilm formation ability was seen between Bcc and *S. aureus* isolates.




**Figure 3.2 Biofilm formation by Bcc and *S. aureus* strains.**

Overnight bacterial cultures were diluted, plated into 96-well polystyrene plates, and incubated at 37°C on a 3-dimensional rotator for 24 h. Biofilm structure was quantified by staining with crystal violet and absorbance read at 550 nm. Data are presented as mean  $\pm$  SEM from three independent experiments.



### 3.4 Modification of IL-8 Production by *B. cenocepacia* and *S. aureus* sp. in 16HBE Cells

To study the effect of bacterial pathogens on IL-8 production, 16HBE cells were infected with *B. cenocepacia* J2315 (MOI 5) and *S. aureus* strains (MOI 0.5) for 24 h. IL-8 concentration in the culture medium was determined using ELISA. Under basal conditions 16HBE cells produced low level of IL-8, indicating it has endogenous IL-8 expression (Figure 3.3A). CF-isolated pathogen *B. cenocepacia* J2315 induced a greater level of IL-8 (increased by 102%,  $p < 0.05$ ) compared to the baseline. Interestingly, *S. aureus* isolates possessed different abilities of inducing IL-8 in 16HBE cells. Strains such as Newman, NCTC 6571 and MRSA252 induced much less IL-8, whereas some strains stimulated abundant levels of IL-8 (SH1000, C316, RN4220, and USA400,  $p < 0.05$ ). However, bacterial pathogens cause damage to epithelial cells, which could potentially result in reduced IL-8 accumulation. To eliminate this possibility, the LDH assay was used to assess cytotoxicity of *B. cenocepacia* J2315 and *S. aureus* isolates (Figure 3.3B). Under basal conditions, a minimal amount of LDH was released ( $1.9 \pm 1.1$  %). Even though the inoculum CFU of *B. cenocepacia* J2315 was 10 times higher than *S. aureus*, it caused the lowest level of cell damage ( $11 \pm 1.6$  %). This might due to the slow growing nature of J2315 (Kaza et al., 2010), which results in minimal amount of toxins secretion. Among the cytotoxic *S. aureus* isolates, Newman and NCTC 10788 were isolated from a human lesion, MRSA41 is hospital-acquired MRSA, while C316 and USA400 are community-associated *S. aureus*. Most *S. aureus* strains that significantly reduced IL-8 production did not show major cytotoxicity, indicating that the inhibition of IL-8 production was not due to cell death. As biofilm formation is a key virulence factor, we compared IL-8 production in response to *B. cenocepacia* J2315 or *S. aureus* isolates versus biofilm development *in vitro* (Figure 3.3C). However, no correlation was observed between these two variables ( $r^2 = 0.06689$ ,  $p = 0.3520$ ). Taken together, *S. aureus* species show heterogeneity in induction of IL-8 in 16HBE cells, and it is not correlated with biofilm formation.




**Figure 3.3 IL-8 production and cytotoxicity caused by *B. cenocepacia* J2315 and *S. aureus* sp. in 16HBE cells.**

**(A)** IL-8 secretion in response to *B. cenocepacia* J2315 (MOI 5) and *S. aureus* sp. (MOI 0.5) for 24 h. Data are presented as mean  $\pm$  SEM from three independent experiments in duplicate. Statistical analysis was performed using two-way ANOVA with Dunnett's multiple comparisons test. **(B)** Cytotoxicity of *B. cenocepacia* J2315 (MOI 5) and *S. aureus* sp. (MOI 0.5) by measuring LDH release. Data are plotted as mean  $\pm$  SEM from three independent experiments performed in triplicate. Statistical analysis was performed using one-way ANOVA with Dunnett's multiple comparisons test. \*,  $p < 0.05$  compared to the control. **(C)** Scatter plot of IL-8 production in response to *B. cenocepacia* J2315 or *S. aureus* isolates versus biofilm formation. Correlation between biofilm formation and IL-8 was calculated using Pearson correlation score.

### 3.5 PI3K is required for the proinflammatory response to *B. cenocepacia* J2315

To define the role of PI3K/Akt on host response to *S. aureus* or Bcc infections, 16HBE cells were pretreated with LY294002 or wortmannin for 1 hour and infected with *S. aureus* or Bcc at an MOI of 50. IL-8 concentration in the cultured medium was quantified using ELISA. A heterogeneity was seen in Bcc-induced IL-8 production in the absence of PI3K inhibitors (Figure 3.4). Despite *B. multivorans* 13243 and *B. cenocepacia* J2315 isolated from CF individuals, J2315-infected 16HBE cells produced threefold more IL-8 compared to 13243-infected. As expected, the chemokine response to botanical isolate *B. anthina* J2552 was the lowest, which was consistent with biofilm formation. A slightly higher concentration of IL-8 was seen in response to MRSA252 in comparison with MSSA NCTC 6571. Little or no reduction was seen with the presence of LY294002 or wortmannin, with the exception of J2315. In consistence with Cremer et al (2011), pretreatment with LY294002 or wortmannin caused significant reduction in the J2315-induced IL-8 response (decreased by 34% and 26%, respectively). Thus, these results indicate that the proinflammatory response to J2315 requires PI3K activity in 16HBE cells.

Due to its strong ability of inducing pro-inflammatory chemokine, *B. cenocepacia* J2315 was chosen to investigate pro-inflammatory responses and modulation on host signalling for future experiments.



**Figure 3.4 IL-8 response to *S. aureus* and Bcc with the presence or absence of PI3K inhibitors in 16HBE cells.**

Cells were treated with LY294002 (30  $\mu$ M) or wortmannin (100 nM) for 1 hour before infected with *S. aureus* or Bcc at an MOI of 50. At 2 h postinfection, 16HBE cells were rinsed with DPBS twice and incubated in fresh DMEM for 18 h. IL-8 production was quantified by ELISA. Data are presented as mean  $\pm$  SEM from three independent experiments performed in duplicate. \*,  $p < 0.05$  compared to J2315-infected cells without any PI3K inhibitors; #,  $p < 0.05$  compared to the control; using two-way ANOVA followed by Dunnett's multiple comparisons test.

### 3.6 Discussion

In this chapter, we investigated the host response to Bcc and *S. aureus* isolates, respectively, including the disruption of the airway barrier, induction of pro-inflammatory chemokine IL-8, and cell damage.

Primary airway epithelial cells and cell lines are widely used in host-pathogen interaction studies. Culturing Calu-3 cells on a permeable membrane has proven to represent a great similarity to primary human bronchial epithelial cells phenotype, with cilia-like structures, secretory granules and a high TEER (Grainger et al., 2006). Human bronchial epithelial cells are considered to be the most physiological model, but they have a highly differentiated phenotype due to donor variation, experiment and passage. Calu-3 cells have a more reproducible TEER and phenotype, although data have shown some distinctive differences, such as ZO-1 (a tight junction marker) expression and  $\beta$ -tubulin IV (a ciliated cell marker, expression localised to the cell boundaries) localisation (Stewart et al., 2012). The modulation of MRSA252 and *B. multivorans* 13243 on TEER value and FITC-dextran diffusion of Calu-3 monolayers proves the disruptive effect on the integrity of airway epithelial monolayers. Studies working on Calu-3 cells show that *S. aureus* isolates and *B. multivorans* 13243 caused large falls in TEER (Kiedrowski et al., 2016; Duff et al., 2006). Furthermore, *S. aureus*-secreted products are potent at breaching tight junctions of epithelial barriers (Malik et al., 2015). Duff et al. (2006) demonstrate that although all the tested Bcc isolates reduced TEER within 4 h, translocation speed of Bcc from the apical to the basolateral chamber differs. In order to study the translocation in our model, we cultured Calu-3 cells on 3  $\mu$ m pore size polyester membrane to allow pathogens to pass through. Despite exhibiting a high TEER value ( $\sim 600 \text{ Ohm.cm}^2$ ), Calu-3 cells formed 'leaky' monolayers on 3  $\mu$ m membranes compared with 0.4  $\mu$ m filter cultures. Therefore, filter pore size can affect cell morphology and cell densities. In a preliminary experiment, translocation was tested with MRSA252, MSSA NCTC 6571, *B. multivorans* 13243, *E. coli* HB101 and *S. Typhimurium* NTB6 at an MOI of 50 for 6 h apically. No translocation was observed except the invasive reference strain *S. Typhimurium* NTB6 (Table 3.1).

**Table 3.1 Translocation in airway epithelial monolayer**

Unit: CFU	Time	MRSA252	MSSA NCTC 6571	<i>B. multivorans</i> 13243	<i>E. coli</i> HB101	<i>S. Typhimurium</i> NTB6
Inoculum	0 h	$5.8 \times 10^6$	$4.8 \times 10^6$	$5.3 \times 10^6$	$5.4 \times 10^6$	$5.6 \times 10^6$
Basolateral chamber	2 h	0	0	0	0	$1.5 \times 10^4$
	4 h	0	0	0	0	$3.9 \times 10^6$
	6 h	0	0	0	0	$3.8 \times 10^7$
Apical chamber	6 h	$3.3 \times 10^7$	$2.9 \times 10^7$	$3.6 \times 10^8$	$6.0 \times 10^7$	N/A

Biofilm formation is a major challenge of bacterial eradication, particularly in chronic lung infections. We showed differential biofilm development with a selection of Bcc and *S. aureus* strains. As expected, CF-associated isolates developed more robust biofilms compared to the environmental isolate J2552. *B. cenocepacia* J2315 belongs to the epidemic lineage ET12, which makes up the majority of *B. cenocepacia* isolates from CF patients and accounts for a dramatically shorter survival expectancy (Jones et al., 2004). Isolates of *B. multivorans* and *B. cenocepacia* have been shown to form stronger biofilm and are more invasive of 16HBE cells than *B. cepacia*, *B. stabilis* and *B. vietnamiensis* (Caraher et al., 2007). A study (Cunha et al., 2004) described the isolation of 15 Bcc strains from 21 CF patients and the testing the abilities of biofilm formation *in vitro*. Although the results suggested that the virulence of lung infections caused by Bcc is not correlated with biofilm formation, three patients who died during 7-year surveillance period were hosting biofilm-forming strains. Our results show that *S. aureus* strains have different abilities of biofilm development, and it is uncoupled with their abilities of IL-8 induction, which suggests the proinflammatory response of airway epithelial cells caused by *S. aureus* depends on virulence determinants other than biofilm formation.

The recognition of microorganisms is the first step of host defence. The Gram-negative pathogen *B. cenocepacia* J2315 has prototypical agonists for TLR2, 4 and 5, which are lipopeptide, LPS and flagellin, respectively (West et al., 2009); whereas Gram-positive *S. aureus* species are mainly detected by TLR2. TLR2/1 and TLR2/6 heterodimers recognise a wide range of *S. aureus* cell wall components, such as teichoic acid, peptidoglycan and lipoteichoic acid (Fournier and Philpott, 2005). The stimulation of Toll-like receptors by conserved microbial surface ligands initiates sequential events, including the activation of the NF- $\kappa$ B pathway, proinflammatory cytokine and chemokine secretion, and activation and recruitment of professional immune cells. We showed that *B. cenocepacia* J2315 elicited a robust IL-8 production, whereas heterogeneity was seen in *S. aureus* isolates-

induced IL-8 production. This is consistent with the finding of differential abilities of nasal *S. aureus* isolates to induce proinflammatory cytokine TNF $\alpha$  production by PBMCs (Peres et al., 2015). Mayer et al. (2007) demonstrate that human bronchial epithelial cell line BEAS-2B cells had dose-dependent IL-8 production in response to Gram-negative pathogens whereas they were hypo-responsive to Gram-positive bacteria such as *S. aureus*. TLR2 and its co-receptor CD36 co-transfection restored IL-8 responses to TLR2 agonists. Therefore, the expression of TLR2 and CD36 are limiting factors of the pro-inflammatory response to *S. aureus*. However, in our system, a few MSSA strains (SH1000 and RN4220) are strong inducers of IL-8 production, which suggested they might express different surface ligands.

Inhibition of PI3K activity with pharmacological inhibitors did not change IL-8 production in *S. aureus*- or Bcc-stimulated 16HBE cells, with the exception of *B. cenocepacia* J2315, suggesting only J2315 is capable of inducing a robust proinflammatory response of 16HBE cells that is PI3K-dependent. One possible explanation is the scale of IL-8 response induced by J2315 is much higher than the other strains, and therefore the suppressing effect of PI3K inhibitors is inclined to be larger. The concentrations of LY294002 (30  $\mu$ M) and wortmannin (100 nM) used here are typical concentrations for inhibiting PI3K activity (Davies et al., 2000). The PI3K regulation of the proinflammatory response to *B. cenocepacia* in macrophages has been reported previously (Cremer et al., 2011). *B. cenocepacia* stimulates NF- $\kappa$ B activity by activating PI3K/Akt signalling, followed by inactivating the downstream NF- $\kappa$ B repressor GSK3 $\beta$ . However, PI3K/Akt plays different roles in the immune response depending on the cell type and stimulus. In neutrophils, PI3K promotes migration, phagocytosis, respiratory burst and survival (Martin et al., 2015; Luo and Mondal, 2015), whereas in monocytes (Luyendyk et al., 2008), macrophages (Androulidaki et al., 2009) and dendritic cells (Fukao et al., 2002), PI3K activity negatively regulates the pro-inflammatory response to maintain homeostasis. Indeed, defining the role of PI3K in inflammation as a positive or negative regulator is oversimplified, as different PI3K and Akt isoforms finely tune the inflammatory responses in distinct ways as complex as the inflammatory response itself. For example, while Akt1 promotes M2 macrophage polarisation, Akt2 contributes to M1 phenotype polarisation (Androulidaki et al., 2009; Arranz et al., 2012; Vergadi et al., 2014). Akt1<sup>-/-</sup> macrophages are hypo-responsive to LPS, exhibit higher levels of iNOS and enhanced bacterial clearance (Arranz et al., 2012), whereas Akt2<sup>-/-</sup> macrophages are hyper-responsive to LPS, suppress TLR4 signalling through the anti-inflammatory microRNA miR-146a (Vergadi et al., 2014). Furthermore, the p110 $\delta$  isoform of PI3K promotes neutrophil flux and inflammation in macrophages (Konrad et al., 2008). PI3K p110 $\delta$  plays an opposing

role in bone marrow-derived dendritic cells by inducing internalisation of TLR4 and degradation of the adaptor molecule TIRAP, which switches from proinflammatory phase to anti-inflammatory phase, and thereby, enhances anti-inflammatory cytokines (IL-10 and IFN- $\beta$ ) production (Aksoy et al., 2012). Furthermore, PI3K p110 $\delta$  is found to play a dominant role in IL-10 production in response to *S. aureus* through the PI3K/Akt/mTOR pathway (Peres et al., 2015). In this project, we intended to investigate the anti-inflammatory response to *S. aureus* and Bcc by measuring IL-10 production. However, 16HBE cells do not express IL-10 at mRNA or protein level (Massengale et al., 1999), and we could not detect any IL-10 production using ELISA (data not shown).

### **3.7 Conclusions**

The results presented in this chapter demonstrate the host response of airway epithelial cells stimulated by *S. aureus* and *Burkholderia cepacia* complex isolates. Barrier function of polarised airway epithelial cells was disrupted by MRSA252 and *B. multivorans* 13243. *S. aureus* isolates had differential abilities of inducing IL-8 production, which did not correlate with biofilm development. *B. cenocepacia* J2315 induced a strong IL-8 production, in part through PI3K signalling. Therefore, the role of PI3K/Akt in infection process needs further investigation.

# **Chapter 4 Investigating the Role of PI3K/Akt, MAPK and NF- $\kappa$ B Signalling in Bacterial Infection of Human Airway Epithelial Cells**



## 4.1 Introduction

Bacterial adherence to the airway lumen is an essential step of infection, followed by internalisation depending on the lifestyle of the pathogen (Silva, 2012). Multiple bacterial surface structures are involved in adhesion and invasion, such as cell wall-anchored proteins (Foster et al., 2014), the cable pili (Essex-Lopresti et al., 2005), and flagella (Bucior et al., 2012). These adhesins interact with host receptors and activate host kinase pathways in order to trigger actin rearrangement, which enables invasion and uptake of bacteria by endocytosis (Colonne et al., 2016). By doing so, intracellular bacteria are temporally protected from antimicrobial substances (such as nitric oxide, defensins, and lysozymes) in the extracellular milieu, as well as evade recognition and phagocytosis by immune cells (Eisele and Anderson, 2011). Pathogens exploit the intracellular niche as a reservoir for proliferation, and subsequent dissemination which also requires the manipulation of actin rearrangement (Colonne et al., 2016). Hence, subverting the interaction between pathogen and cytoskeleton may provide a novel antimicrobial treatment.

The microbial detectors (PRRs) are widely expressed in phagocytes and non-professional phagocytes such as epithelial cells and fibroblasts (Takeuchi and Akira, 2010). As a family of well characterised PRRs, TLRs are involved in host response to bacteria, viruses, and parasites. Activation of TLR leads to sequence of events: activation of downstream signalling cascades (including the MAPK and NF- $\kappa$ B pathways) (Takeuchi and Akira, 2010), nuclear translocation of transcription factors (such as NF- $\kappa$ B, AP-1, IRFs) (Troutman et al., 2014), and production of cytokines/chemokines and antimicrobial substances (Medzhitov, 2001; Selsted and Ouellette, 2005). In addition to the innate immune response, TLRs induce antigen uptake and presentation. Thus, TLR provides double protection to the host by (1) removing invading pathogens by directly phagocytosis and (2) triggering the adaptive immune response if the innate immune response is not sufficient to eradicate pathogens. However, bacterial pathogens have evolved several tactics to circumvent immune clearance by manipulating TLR signalling.

PI3K/Akt finely tunes TLR signalling both positively and negatively. TLR2-mediated NF- $\kappa$ B activation requires PI3K activity, as blocking PI3K using pharmacological inhibitors hinders phosphorylation of p65 NF- $\kappa$ B (Strassheim et al., 2004). While PI3K/Akt is involved in the pro-inflammatory response, this signalling cascade prevents excessive inflammation by promoting the anti-inflammatory response (Fukao and Koyasu, 2003). For example, the transcription factor FoxO1 enhances TLR4-mediated signalling, which involves activation of Akt (Fan et al., 2010b). However, activated Akt negatively regulates

FoxO1, thereby putting the brake on TLR4 signalling. Although the role of PI3K in inflammation remains controversial, it is undeniable that owing to its role in cellular proliferation and survival, PI3K is beneficial to the host during an immune response (Weichhart and Säemann, 2008).

In addition, PI3K activity is critical for cytoskeleton rearrangement and bacterial invasion (Kierbel et al., 2005; Oviedo-Boyso et al., 2011). Therefore, inhibiting PI3K activity may be a potential target to reinforce host barrier. However, although many studies have been done with myeloid cells, the role of PI3K/Akt in airway epithelial cells in response to bacterial pathogens remains unclear.


The aims of this chapter were to investigate modulation of PI3K, MAPK and NF- $\kappa$ B signalling by *S. aureus* and Bcc in bronchial epithelial cells, and particularly the role of PI3K in bacterial invasion.

## **4.2 PI3K activity regulates *S. aureus* invasion of airway epithelial cells**

To assess *S. aureus* and Bcc adhesion, human airway epithelial 16HBE cells were infected by MRSA252, MSSA NCTC 6571, or *B. multivorans* 13243 at an MOI of 50 for 2 hours. *E. coli* HB101 and *S. Typhimurium* NTB6 were used as the negative and positive control, respectively. Adhered bacteria were counted by determining the viable count of the epithelial cell lysates and presented as a percentage of the inoculum (Figure 4.1A). The result showed that cellular adherence by MRSA252 was 1-fold higher than MSSA NCTC 6571, whereas only minimal amount of *B. multivorans* 13243 ( $4.6\% \pm 1.4\%$ ) attached to 16HBE cells. To assess *S. aureus* and Bcc internalisation, 16HBE cells were exposed to MRSA252, MSSA NCTC 6571, or *B. multivorans* 13243 (MOI 50) for 2 hours, followed by 1- or 2-hour antimicrobial treatment to eliminate extracellular bacteria. The number of internalised bacteria was determined using viable count and shown as a percentage of adherence (Figure 4.1B). Although less amount of MSSA NCTC 6571 and *B. multivorans* 13243 adhered on epithelial cells, their percent of internalisation was higher than MRSA252.


To investigate the role of PI3K/Akt in bacterial adherence and invasion, two structurally distinct pharmacological inhibitors of PI3K, LY294002 (Vlahos et al., 1994) and wortmannin (Arcaro and Wymann, 1993) were used. In Figure 4.2A and B, 16HBE cells were treated with 10  $\mu$ M LY294002 30 minutes prior to bacterial infections. LY294002 upregulated marginally the adherence of MRSA252 and *S. Typhimurium* NTB6, whereas *B. multivorans* 13243 attachment was reduced by 60%. Only minimal numbers of MRSA252 and MSSA NCTC 6571 entered into epithelial cells, and this was decreased

with LY294002. The invasion assay was also performed with a higher concentration of LY294002 (30  $\mu$ M) and wortmannin (100 nM) (Figure 4.2C and D). MRSA252 and MSSA NCTC 6571 maintained the same level of adherence, whereas LY294002 caused a modest reduction, and wortmannin caused a modest increase of adherence of *B. multivorans* 13243. LY294002 reduced the invasion of MRSA252, MSSA NCTC 6571 and *B. multivorans* 13243 by more than 50%. In contrast, wortmannin only caused a decrease for *B. multivorans* 13243. Further invasion assays were performed with Bcc isolates (Figure 4.2E and F). In comparison with the CF-related pathogens *B. multivorans* 13243 and *B. cenocepacia* J2315, the environmental isolate *B. anthina* J2552 did not adhere to epithelial cells. Although the pretreatment of PI3K inhibitors was extended to 1 hour, no effect was observed on adherence. Internalisation of *B. anthina* J2552 was blocked by LY294002. Altogether, in the process of invasion, PI3K is not involved in adherence but is involved in internalisation, particularly for *S. aureus*.



**Figure 4.1 Pilot study of adherence to and invasion of airway epithelial 16HBE cells.**

**(A)** Adherence of MRSA252, MSSA NCTC 6571, or *B. multivorans* 13243 to 16HBE cells at 2 hours post infection (MOI 50). *E. coli* HB101 and *S. Typhimurium* NTB6 were used as the negative control and the positive control, respectively. **(B)** Invasion of 16HBE cells by MRSA252, MSSA NCTC 6571, or *B. multivorans* 13243 was quantified after 1 or 2 hours antibiotic treatment. Data are presented as mean  $\pm$  SEM from three independent experiments performed in duplicate. \*,  $p < 0.05$  compared to *E. coli* HB101, using two-way ANOVA followed by Dunnett's multiple comparisons test.



**Figure 4.2 Inhibition of bacterial adherence and internalisation by pharmacological inhibitors of PI3K.**


**(A and B)** After 30 minutes treatment with LY294002 (10  $\mu$ M), 16HBE cells were infected with MRSA252, MSSA NCTC 6571, *B. multivorans* 13243, *E. coli* HB101 or *S. Typhimurium* NTB6 (50 MOI) for 2 hours with the presence of LY294002. 16HBE cells were washed 3 times with DPBS to remove nonadherent bacteria and subsequently lysed in 1% saponin (100  $\mu$ L) to quantify adhered bacteria using viable count. The percentage of adhered bacteria was normalised against the inoculum. Alternatively, extracellular adherent bacteria were killed with antibiotics for another hour (2 hours for Bcc strains) to measure the amount of internalised bacteria. The percentage of internalised bacteria was calculated against the number of adhered bacteria. Adherence **(A)** and invasion **(B)** are shown as mean  $\pm$  SEM from four independent experiments, each performed in duplicate. Data were analysed using two-way ANOVA following Bonferroni's multiple comparisons test. \*,  $p < 0.05$ .

**(C and D)** After 30 minutes treatment with LY294002 (30  $\mu$ M) or wortmannin (100 nM), 16HBE cells were challenged with MRSA252, MSSA NCTC 6571, *B. multivorans* 13243, *E. coli* HB101 or *S. Typhimurium* NTB6 (50 MOI) with the presence of inhibitors. Adherence **(C)** and invasion **(D)** were assessed as previous and are shown as mean  $\pm$  SEM from three independent experiments in duplicate. Data were analysed using repeated measures two-way ANOVA following Dunnett's multiple comparisons test.

**(E and F)** After 1 hour treatment with LY294002 (30  $\mu$ M) or wortmannin (100 nM), 16HBE cells were infected with *B. multivorans* 13243, *B. cenocepacia* J2315, *B. anthina* J2552, *E. coli* HB101 or *S. Typhimurium* NTB6 (50 MOI) with the presence of inhibitors. Adherence **(E)** and invasion **(F)** were assessed as previous and are shown as mean  $\pm$  SEM from three independent experiments, each performed in duplicate. Data are analysed using repeated measures two-way ANOVA following Dunnett's multiple comparisons test.


### 4.3 Modulation of PI3K/Akt signalling by *S. aureus* isolates

Akt is the major downstream target of PI3K and activated at Ser473 and Thr308 through a dual phosphorylation mechanism. In order to study the role of PI3K, airway epithelial 16HBE cells were infected with planktonic (Figure 4.3) or biofilm (Figure 4.4) MRSA252 or MSSA NCTC 6571 cultures, and phospho-Akt (Ser473) and phospho-Akt (Thr308) were analysed by immunoblotting. Although planktonic *S. aureus* cultures were used at a high dose ( $10^8$  CFU/mL), they failed to activate Akt from 5 to 30 minutes. With their biofilm counterparts, phospho-Akt (Ser473) peaked at 15 minutes but no significant difference was observed (Figure 4.4A and C).



**Figure 4.3 Planktonic MRSA252 or MSSA NCTC 6571 do not affect PI3K activity.**

Human airway epithelial 16HBE cells were starved overnight. LY294002 (10  $\mu$ M) overnight was used as negative control and TNF $\alpha$  (100 ng/mL) for 15 minutes was used as positive control. Cells were infected with MRSA252 or MSSA NCTC 6571 culture ( $10^8$  CFU/mL) for the time indicated. Cell lysates were analysed by immunoblotting to quantify phospho-Akt (Ser473 and Thr 308). The fluorescence signals were normalised against total Akt and are presented as mean  $\pm$  SEM from three independent experiments.




**Figure 4.4 Biofilm MRSA252 or MSSA NCTC 6571 do not affect PI3K activity.**

Human airway epithelial 16HBE cells were starved overnight. LY294002 (10  $\mu$ M) overnight was used as the negative control and TNF $\alpha$  (100 ng/mL) for 15 minutes was used as the positive control. Cells were infected with MRSA252 or MSSA NCTC 6571 culture (10<sup>8</sup> CFU/mL) for the time indicated. Cell lysates were analysed by immunoblotting to quantify phospho-Akt (Ser473 and Thr308). The signals were normalised against total Akt. Data are presented as mean  $\pm$  SEM from three independent experiments. \*,  $p < 0.05$  compared with the control using repeated measures one-way ANOVA followed by Dunnett's multiple comparisons test.

#### **4.4 *B. cenocepacia* J2315 and *S. aureus* supernatants modulating PI3K/Akt, MAPK and NF- $\kappa$ B signalling in human airway epithelial cells**

To investigate the effect of *B. cenocepacia* on MAPK signalling, 16HBE cells were challenged with the robust IL-8 inducer J2315 (Figure 3.3), at various MOIs (Figure 4.). With an MOI of 5, one-fold upregulation of phosphorylation of p38 was observed at 30 and 60 minutes post infection. In contrast, phosphorylation of Erk was not changed by any treatment. With 50 MOI, the level of phosphorylation of p38 and Erk did not change. J2315 failed to activate NF- $\kappa$ B signalling, as no increase in phosphorylation of I $\kappa$ B $\alpha$  was observed. Furthermore, higher MOI (100) of J2315 did not activate phosphorylation of I $\kappa$ B $\alpha$  or p65.


Since it has been reported that *S. aureus* supernatants suppress TLR2 agonist Pam<sub>3</sub>CSK4-activated NF- $\kappa$ B pathway in human bronchial epithelial cells, to replicate this in our system, 16HBE cells were stimulated with different concentrations of MSSA NCTC 6571 supernatant (Chekabab et al., 2015). When the cells were treated with lower dilution of MSSA NCTC 6571 supernatant (10% v/v, Figure 4.A and B), only a minimal increase in phosphorylation of p38 was observed at 60 minutes post stimulation, whereas phosphorylation of Erk was significantly inhibited at 15 minutes. In contrast, 25% (v/v) MSSA NCTC 6571 supernatant downregulated phosphorylation of p38 at 5 and 30 minutes, and recovered back to baseline at 240 minutes (Figure 4.C). The bacterial culture medium trypticase soy broth (TSB, 10% v/v, 2 hours stimulation) has previously been shown to significantly downregulate Erk signalling in airway epithelial S9 cells but not 16HBE cells (Below et al., 2009). In Figure 4.C and D, 4 hours treatment with 25% (v/v) LB did not change phosphorylation of p38 but increased phosphorylation of Erk by 50%. This suggests the upregulation of phospho-Erk may result from LB but not *S. aureus* supernatant.



**Figure 4.5 *B. cenocepacia* J2315 has no effect on MAPK or NF-κB signalling.**

Airway epithelial 16HBE cells were stimulated with *B. cenocepacia* J2315 at various MOIs (**A and B**, MOI=5; **C-F**, MOI=50; **G and H**, MOI=100) for the indicated time. (**A and B**) TNFα (20 ng/mL) and IL1β (20 ng/mL) for 20 minutes were used as the positive control. (**E-H**) TNFα (20 ng/mL) for 5 minutes was the positive control. Phosphorylation of p38, Erk, IκBα, and p65 were detected by western blotting and the fluorescence signals were normalised against p38. Data are shown as mean ± SEM from three independent experiments and analysed using repeated measures one-way ANOVA following Dunnett's multiple comparisons test.






**Figure 4.6 Analysis of Erk and p38 by MSSA NCTC 6571 supernatant.**

16HBE cells were incubated with 10% (v/v) (**A and B**) or 25% (v/v) (**C and D**) MSSA NCTC 6571 conditioned medium for the indicated time. In **C and D**, cells were treated with 25% v/v LB medium for 4 hours as a control for LB. Western blots on cell lysates were done to measure phospho-p38, phospho-Erk. Membranes were reprobbed with loading control total p38 and Erk antibodies. Data are shown as mean  $\pm$  SEM from three independent experiments. \*,  $p < 0.05$  compared with the control using repeated measures one-way ANOVA following Dunnett's multiple comparisons test.


To optimise the dose of stimuli, 16HBE cells were treated with increasing dose of *B. cenocepacia* J2315 or MSSA NCTC 6571 supernatant for 1 hour (Figure 4.7). A dose-dependent increasing trend was observed with J2315. In contrast, *S. aureus* supernatant failed to regulate Erk signalling.



**Figure 4.7 MAPK response to *B. cenocepacia* J2315 live bacteria or MSSA NCTC 6571 supernatant.**


Phosphorylation of p38 and Erk of 16HBE cells after stimulated with *B. cenocepacia* J2315 (**A and B**) or MSSA NCTC 6571 supernatant (v/v, **C and D**) for 1 hour. Phospho-p38 and phospho-Erk of cell lysates were measured by western blot. Membranes were reprobbed with loading control total p38 and Erk antibodies. Data are shown as mean  $\pm$  SEM from three independent experiments and analysed using repeated measures one-way ANOVA following Dunnett's multiple comparisons test. \*,  $p < 0.05$ .

As surface structures (such as lipopeptides, flagellin and LPS) of *B. cenocepacia* play important roles in host-pathogen interactions, the potentially damaging effect of centrifugation in the bacteria preparation process should be considered (Peterson et al., 2011). In Figure 4., 16HBE cells were infected with unwashed J2315 at an MOI of 5 for up to 180 minutes and phosphorylation of Akt was observed with maxima at 120 minutes. The Akt downstream target GSK-3 $\beta$  was phosphorylated marginally compared to the control. For the MAPK family, phosphorylation of p38 showed a 1-fold increase at 120 minutes, whereas phosphorylation of Erk peaked at 5 minutes, followed by a decrease and then a secondary rise at 120 minutes. For NF- $\kappa$ B signalling, J2315 upregulated phosphorylation of p65, particularly after 120 minutes. Except for NF- $\kappa$ B, TNF $\alpha$  failed to activate PI3K/Akt and MAPK signalling as a positive control. Due to differential signalling response to unwashed J2315 culture comparing with washed culture (Figure 4.), a further experiment was performed to directly compare the difference. As shown in Figure 4., washed and unwashed J2315 cultures were used to stimulate 16HBE cells for 5 and 120 minutes. Although no significant difference was shown in phosphorylation of Akt and GSK-3 $\beta$ , a higher level of phosphorylation of Akt was observed with unwashed culture at 120 minutes. Both washed and unwashed cultures had 2-fold higher phosphorylation of p38 at 120 minutes compared with the control (Figure 4.C). In contrast, with washed culture, phosphorylation of Erk showed a 1-fold increase comparing 120 to 5 minutes, whereas with unwashed culture a 1-fold increase in phosphorylation of Erk exhibited both at 5 and 120 minutes, which is consistent with Figure 4.D. Similar to the previous results, washed and unwashed culture induced robust phosphorylation of p65 at 120 but not 5 minutes.



**Figure 4.8 Host signalling modulation by unwashed *B. cenocepacia* J2315 culture.**

16HBE cells were infected by *B. cenocepacia* J2315 unwashed culture at an MOI of 5 for the indicated time. TNF $\alpha$  (20 ng/mL) for 5 minutes was used as the positive control. Phosphorylation of Akt (**A**), GSK-3 $\beta$  (**B**), p38 (**C**), Erk (**D**), p65 (**E**) of cell lysates were measured by western blot and normalised against total p38. (**F**) Western blot analysis for levels of phosphorylated Akt, phosphorylated GSK-3 $\beta$ , phosphorylated p38, phosphorylated Erk, phosphorylated p65, and p38 in cell lysate following *B. cenocepacia* J2315 stimulation. Protein bands representative of three independent experiments. Data are shown as mean  $\pm$  SEM. \*,  $p < 0.05$  compared with the control using repeated measures one-way ANOVA following Dunnett's multiple comparisons test.



**Figure 4.9 Comparison of washed and unwashed *B. cenocepacia* J2315 culture on stimulating 16HBE signalling.**

Overnight culture of *B. cenocepacia* J2315 were cultured into exponential phase. The culture was divided into two portions. Washed culture was centrifuged to collect bacteria and washed by DPBS once. Washed and unwashed culture were adjusted to the same optical density. 16HBE cells were challenged by *B. cenocepacia* J2315 at an MOI of 5 for 5 or 120 minutes. TNFα (20 ng/mL) for 5 minutes was used as the positive control. Phosphorylation of Akt (**A**), GSK-3β (**B**), p38 (**C**), Erk (**D**), p65 (**E**) of cell lysates were measured by western blot and normalised against total p38. (**F**) Western blot analysis for levels of phosphorylated Akt, phosphorylated GSK-3β, phosphorylated p38, phosphorylated Erk, phosphorylated p65, and p38 in cell lysate following *B. cenocepacia* J2315 stimulation. Protein bands representative of three independent experiments. Data are shown as mean ± SEM. \*,  $p < 0.05$  compared with the control using repeated measures one-way ANOVA following Tukey's multiple comparisons test.

## 4.5 Discussion

In this chapter, we investigated cellular invasion and signalling cascades mediated by *S. aureus* and Bcc isolates in human bronchial epithelial 16HBE cells.

*S. aureus* is identified as an extracellular pathogen associated with skin and soft tissue infections in the traditional classification, whereas Bcc are dual intracellular/extracellular style pathogens (Wiersinga et al., 2006). However, there is growing evidence of an intracellular phase of *S. aureus* (Garzoni and Kelley, 2009). Here we assessed invasion by *S. aureus* and *B. multivorans*, as well as two control strains, in 16HBE cells. At 2 hours post infection, adherence of MRSA252, MSSA NCTC 6571 and *B. multivorans* 13243 showed 45%, 25% and 10% of *S. Typhimurium* NTB6, respectively, whereas internalisation was 3%, 15% and 20%. This suggests their infective lifestyles in the host are consistent with the previous study (Silva, 2012). Phosphoinositide in the cell is essential for membrane dynamics, actin polymerisation and vesicle trafficking. Therefore, by disrupting phosphoinositide homeostasis, pathogens are able to enhance their uptake by host cells (Ham et al., 2011). Gessain et al. (2015) gave further insight into the critical role of PI3K. Despite no intrinsic PI3K activity in placental epithelial cells, a surface protein of *Listeria monocytogenes* InlB activates PI3K/Akt, which enables bacterial invasion (Gessain et al., 2015),

To elucidate the role of PI3K in airway epithelial cell membrane permissiveness, we pretreated 16HBE cells with PI3K specific inhibitors LY294002 or wortmannin, followed by an invasion assay. LY294002 inhibits PI3K activity by competing the ATP binding site of catalytic subunit p110 reversibly, whereas wortmannin irreversibly causes covalent modification of p110 kinase and leads to a conformation rearrangement (Walker et al., 2000). LY294002 (30 minutes treatment) failed to affect adherence of tested isolates, with the exception of *B. multivorans* 13243. However, the downregulation effect was not consistent with a longer pretreatment period (1 h). This part requires more replicates to validate the results. Of note, although only minimal amounts of *S. aureus* entered into epithelial cells, a consistent decrease was seen with LY294002, suggesting internalisation of *S. aureus* requires PI3K activity. This result is consistent with a study working with polarised MDCK cells and *P. aeruginosa* strain K (Kierbel et al., 2005). In addition, although pharmacological inhibition of Akt subverts *S. aureus* entry into endothelial cells as well, inhibition of Akt phosphorylation on Thr308 residue does not affect entry, suggesting *S. aureus* internalisation may be associated with phosphorylation of Akt on Ser473 (Oviedo-Boyso et al., 2011). Unexpectedly, although wortmannin is a more potent (IC<sub>50</sub>=5 nM) PI3K inhibitor compared with LY294002 (IC<sub>50</sub>=1.4 µM), in our model

wortmannin failed to cause changes similar to LY294002. We found similar results in Figure 3.4, in which wortmannin did not reduce bacteria-induced IL-8, except for *B. cenocepacia* J2315. This might be due to the instability of wortmannin in the presence of nucleophiles (Yuan et al., 2007).

PI3K/Akt, MAPK and NF- $\kappa$ B are known to play critical roles in response to bacterial pathogens (Krachler et al., 2011). However, pathogens have evolved to exploit these pathways in order to promote survival and proliferation inside the host. First, we investigated activation of PI3K by planktonic or biofilm culture of *S. aureus* isolates. 16HBE cells endogenously express phosphorylated-Akt at a relative high level, which might leave a very small window of unphosphorylated-Akt for *S. aureus*. This might explain why no visible trend was seen in cells stimulated with *S. aureus*. Then we assessed the ability of *B. cenocepacia* J2315 to activate PI3K. Activation of Akt leads to phosphorylation and inactivation of GSK-3 $\beta$  (Ser9). Phosphorylation of Akt (Ser473) peaked at 2 hours, but did not upregulate phosphorylation of GSK-3 $\beta$ . As a key regulator of numerous cellular functions, positive or negative dysregulation of PI3K/Akt in epithelial cells by bacteria has been reported in many studies. For example, invasion of *P. aeruginosa* to either the apical or basolateral surface of polarised airway epithelial Calu-3 cells via pili or flagellin requires PI3K/Akt signalling, and ultimately leads to entry into host cells (Bucior et al., 2012). Apart from invasion, PI3K is a bacterial target to manipulate programmed cell death, depending on the cell types and stage of infection (Knodler et al., 2005). At early stages of infection, the *Salmonella* effector SopB is proved to be anti-apoptotic by continual activation of Akt in HeLa cells, whereas in macrophages, the effector SipB induces cell death in macrophages, thereby escaping phagocytosis (Hernandez et al., 2003).

We further performed a pilot study of the time and dose response to *B. cenocepacia* J2315 and *S. aureus* cell-free supernatant on host signalling. As IL-8 expression is associated with NF- $\kappa$ B (Elliott et al., 2001) and MAPK signalling pathways (Bhattacharyya et al., 2011), these pathways might be involved in *B. cenocepacia* and *S. aureus* infections. The *B. cenocepacia* ET-12 lineage strain K56-2 has been reported to significantly upregulate phosphorylation of Akt, Erk and NF- $\kappa$ B at an MOI of 5 in 16HBE cells within 30 minutes of stimulation (Gillette et al., 2013). In parallel, *B. cenocepacia* BC7 (isolated from a CF patient who died of cepacia syndrome and is able to induce a robust IL-8 response), triggers phosphorylation of p38 and JNK in immortalised CF bronchial epithelial IB3 cells, whereas an environmental *B. cepacia* strain fails to show any activation of these pathways (Sajjan et al., 2008). This indicates that activation of MAPK signalling is related to more intense lung inflammation caused by Bcc isolates.

In contrast, in our system *B. cenocepacia* J2315 failed to activate these pathways even at a high MOI. During the process of preparing the strain, in order to remove metabolites and secreted products, bacterial cultures were centrifuged twice (5000 g for 10 minutes), which might potentially cause damage to bacterial surface structures. Therefore, we performed same experiments using J2315 culture without centrifugation steps. Phosphorylation of Akt, p38, Erk and p65 increased at 120 minutes post-infection. Of note, J2315 without the centrifugation step caused a transient Erk phosphorylation at 5 minutes post-infection and peaked again at 120 minutes, which was reproduced in Figure 4.9. Centrifugation is likely to damage surface structures such as LPS and flagellin. D'Elia et al. observed a similar bi-phasic activation pattern of Erk and p38 in response to *B. pseudomallei* *in vitro* and *in vivo* (D'Elia et al., 2017). Thirty-minutes post-infection, murine alveolar macrophages exhibit transient phosphorylation of p38 and Erk1, and at 2 h post-infection exhibited sustained activation. D'Elia et al. (2017) proposed that the transient activation at the early stage is associated with host defence and during the latter stage of infection host cells might initiate cell death as indicated by continuous Erk activation. *B. pseudomallei* might modulate the dynamics of Erk activation to delay apoptosis, thereby contributing to intracellular survival. Another member of MAPK family, JNK, has shown to be involved in J2315-induced TNF $\alpha$  and IL-1 $\beta$  production in macrophages. In our experiments we were unable to detect phosphorylation of JNK. In addition, we investigated the effect of *S. aureus* supernatant on host signalling cascades, as it has been reported that *S. aureus* supernatant is able to induce different cellular response in human tracheal gland cells compared to live bacteria, particularly a huge upregulation of IL-6 and IL-8 production (Moreilhon et al., 2005). However, in our system, MSSA NCTC 6571 supernatant failed to show consistent modulation on MAPK signalling. Although same conditions (10% v/v, 60 minutes stimulation) were used in Figure 4.6A and B and Figure 4.7C and D, the downregulation on phosphorylation of Erk was not reproduced. This might be caused by repeated propagation of cell lines, or phenotype variance of bacterial culture.

## 4.6 Conclusion

In this chapter we assessed *S. aureus* and Bcc isolates to adhere to and enter into airway epithelial cells. Of note, *S. aureus* internalisation but not adhesion is PI3K-dependent. Hence, PI3K is a critical target for barrier permissiveness by *S. aureus*. In addition, a preliminary study on the ability of *S. aureus*, Bcc live bacteria and *S. aureus* supernatant to modulate host signalling was carried out. PI3K/Akt, MAPK and NF- $\kappa$ B signalling was unchanged when 16HBE cells were infected with bacterial cultures prepared with



centrifugation, whereas undamaged *B. cenocepacia* J2315 culture induced a bi-phasic activation of Erk during the early and the latter stage of infection. This may indicate that *B. cenocepacia* suppresses Erk activity to delay host apoptosis in order to promote survival and proliferation.

# **Chapter 5 Investigating Host Response to Polymicrobial Infections in The Airway**

## 5.1 Introduction

Over the past decades, abundant research has focused on single-species microorganisms and single virulence factors. However, the complex nature of infectious diseases such as airway infections is often associated with multiple species including bacteria, fungi and viruses. Yet we have much less understanding regarding polymicrobial communities within the infection sites, the net impact on progression and manifestations of infectious diseases and host responses to polymicrobial infections. Hence, elucidating the mechanisms of host-pathogen interactions in the context of polymicrobial communities is required to protect against such infection diseases.

Synergism and antagonism are terms that can be applied to describe the cooperative / competitive polymicrobial interactions in disease progression. According to Murray et al. (2014), synergism is 'an interaction of two or more microbes in an infection site that results in enhanced disease compared to infections containing the individual microbe acting alone'. In contrast, the definition of antagonism is 'the presence of two or more species protects the host from disease that occurs when only one species is present' (Nguyen and Oglesby-Sherrouse, 2016). Synergistic interactions between *S. aureus* and *P. aeruginosa* have been discovered in many studies, including increased antimicrobial tolerance and enhanced virulence (Dalton et al., 2011; DeLeon et al., 2014; Fugère et al., 2014; Hoffman et al., 2006). For example, clinical isolates of *P. aeruginosa* and *B. cenocepacia* co-culture exhibit stronger biofilm formation and elicit increased pro-inflammatory responses in a murine chronic lung infection model (Bragonzi et al., 2012). Conversely, *S. aureus* has demonstrated an antagonistic effect in polymicrobial infections. In a 5-year predictive multivariate survivorship model of cystic fibrosis patients, Liou et al. (2001) discussed some known risk factors such as age, gender, and frequency of acute pulmonary exacerbations, and revealed that harbouring *S. aureus* promotes survivorship, whereas *B. cepacia* has a negative effect on survivorship. Indeed, Bcc species are notoriously associated with 'cepacia syndrome', which is characterised with rapid lung function decline and mortality, particularly *B. cenocepacia* (Boussaud et al., 2008). However, it is peculiar that *S. aureus* has protective effects and the reason remains unclear. It may be due to low-level pro-inflammatory or even anti-inflammatory property of *S. aureus*, partially antagonising excessive inflammation and tissue damage resulting from *P. aeruginosa* and Bcc infections (Chavakis et al., 2007). Consistent with this finding, a study by Ahlgren et al. (2015) compared clinical outcomes (for example, lung function and exacerbation rate) of a cohort of 84 CF adults grouped into *S. aureus* infection only, *P. aeruginosa* infection only, and neither *S. aureus* nor *P. aeruginosa*. Patients with *S.*

*aureus* infections had a better clinical outcome and the presence of *S. aureus* appeared a marker of less severe disease in patients compared to those who are infected with *P. aeruginosa* (Ahlgren et al., 2015).

*S. aureus* is known to possess a plethora of immunomodulation strategies. For example, cell wall component SpA is able to activate TNFR1, elicit IL-8 expression and an excessive inflammatory response and tissue damage (Gomez et al., 2004). *S. aureus* also manipulates host responses by blocking neutrophil recruitment (Spaan et al., 2013b), phagocytosis and complement activation (Jongerius et al., 2010b; Jusko et al., 2014), as well as disrupting the adaptive immune response (Wang et al., 2012). Hence, understanding the molecular mechanism how *S. aureus* manipulates the immune system to evade microbial killing may be useful for developing novel treatments against *S. aureus* infection, particularly MRSA-related infections.

The surveillance of innate immune system TLRs recognise numerous microbial ligands and trigger protective immune responses. TLRs occur either as a homodimer such as TLR4 and TLR5, or form heterodimer such as TLR1/2 and TLR2/6. All the TLRs comprise an extracellular leucine-rich repeat domain and a cytoplasmic TIR domain (Lim and Staudt, 2013). TLR and IL-1 receptor belong to the IL-1R/TLR superfamily as they contain homologous TIR domain which is responsible for signalling transduction, and thus signalling pathways are highly conserved between TLRs and IL-1R (Cohen, 2014). IL-1R and all TLRs except TLR3 recruit MyD88 as an adaptor protein, and ultimately lead to activation of NF- $\kappa$ B. Upon activation of TLRs, MyD88 is recruited to the receptors via its TIR domain, and interacts with IRAK1/2/4 via its death domain. Phosphorylation of IRAKs results in the activation of TRAF6, which is an E3 ubiquitin ligase. TRAF6, together with UEV1A and UBC13, synthesises ubiquitin chain and subsequently activates the kinase complex containing TAB1/TAB2/TAK1, which contributes to the activation of MAPK family and nuclear translocation of transcription factors (such as NF- $\kappa$ B and AP-1) and ultimately mounts a pro-inflammatory response (Chen, 2005). Modulation of any signal molecules would affect the intensity and duration of the response and cause harmful outcomes. Therefore, it is vital to fine tune TLR-mediated innate immune response to ensure a protective yet not excessive inflammatory response.


The aims of this chapter were to determine the immunomodulation effect of *S. aureus* on co-infection of *B. cenocepacia*, to reveal the potential molecular mechanisms, as well as to characterise the nature of *S. aureus*-secreted immunomodulation molecules.

## 5.2 Suppression of *B. cenocepacia*-induced IL-8 production by *S. aureus* supernatants

Following co-stimulation with *B. cenocepacia* J2315 and *S. aureus* supernatants or heat-killed bacteria, the neutrophil chemoattractant IL-8 released by airway epithelial 16HBE cells was measured using ELISA. As shown in Chapter 3, *B. cenocepacia* J2315 is able to induce a robust IL-8 production. To optimise MOI (the ratio of bacteria to cells), 16 HBE cells were infected with an MOI ranging from 0.04 to 25 (Figure 5.1A) and all groups showed significant upregulation of IL-8, with a peak at an MOI of 5. Therefore, we chose an MOI of 5, for 24 hours, for the subsequent experiments.


Interestingly, although supernatants (LB medium) of MRSA252 or MSSA NCTC 6571 did not change IL-8 production compared to control, they significantly downregulated IL-8 induced by J2315 (23% and 31% less than J2315-treated group, respectively). To determine whether this effect is medium-specific, the same experiment was performed using supernatant from bacteria cultured in TSB (Figure 5.1C). In contrast, supernatants from *S. aureus* cultured in TSB maintained the level of IL-8 production with the presence or absence of J2315, which indicates that the anti-inflammatory effect is medium-specific. Similarly, J2315-induced IL-8 production did not change with supernatant of MRSA252 biofilm (cultured in LB) but decreased marginally with MSSA NCTC 6571 biofilm supernatant ( $p < 0.05$ , Figure 5.1D).

Surprisingly, heat treatment (95°C, 10 minutes) did not reverse the anti-inflammatory property of *S. aureus* supernatants (Figure 5.1E), whereas heat-killed (100°C, 1 hour) *S. aureus* was unable to modulate IL-8 production (Figure 5.1F). To validate the results, the same experiments were done with a series of *S. aureus* isolates, either supernatants (Figure 5.2A) or heat-killed bacteria (Figure 5.2B). Supernatants from all the tested isolates downregulated the level of IL-8 to varying levels ( $p < 0.05$ ), whereas heat-killed bacteria had no effect on J2315-induced IL-8 production. These data indicate that anti-inflammatory substances exhibit in secreted products but not *S. aureus* cell surface structures. To confirm the suppressing effect is not due to cell damage, an MTT assay was used to assess cell viability (Figure 5.2C and D). Of note, supernatants of SH1000 and RN4220, which caused a doubling in IL-8 production (Figure 5.2A), significantly reduced cell viability. This indicates the correlation between pro-inflammatory secreted products and their virulence. In contrast, heat-killed *S. aureus* did not affect cell viability.



**Figure 5.1 The anti-inflammatory effect of *S. aureus* on *B. cenocepacia* J2315-induced IL-8.**

(A) Dose response of 16HBE cells to *B. cenocepacia* J2315 live bacteria for 24 hours. 16HBE cells were treated with *B. cenocepacia* J2315 at an MOI of 5, in the presence or absence of (B) *S. aureus* supernatants 10% (v/v in LB), (C) 10% (v/v in TSB), (D) 10% biofilm supernatants (v/v in LB), (E) heat-inactivated supernatants 10% (v/v in LB), (F) heat-killed *S. aureus* (5 MOI) for 24 hours. The control group was treated with 10% (v/v) LB in B, D and E; 10% (v/v) TSB in C. IL-8 release was measured by ELISA. Data represent mean  $\pm$  SEM from 3 independent experiments performed in duplicate. Significance was compared to (A) the control or (B-F) J2315-treated group using 2-way ANOVA followed by Dunnett's multiple comparisons test.




**Figure 5.2 IL-8 response to co-stimulation with *B. cenocepacia* J2315 and different *S. aureus* supernatants or heat-killed *S. aureus*.**

16HBE cells were stimulated with *B. cenocepacia* J2315 (5 MOI), in the presence or absence of (A) *S. aureus* supernatants (10% v/v) for 24 hours, or (B) heat-killed *S. aureus* (5 MOI) for 24 hours. The control group was treated with 10% (v/v) LB in A. IL-8 release was measured by ELISA. Data represent mean  $\pm$  SEM from 3 independent experiments performed in duplicate. Significance compared to J2315-treated group using 2-way ANOVA followed by Dunnett's multiple comparisons test. (C, D) MTT assay was used to assess cellular cytotoxicity caused by *S. aureus* supernatants and heat-killed *S. aureus*. Results are expressed as percent of the control, representing mean  $\pm$  SEM (n=3). Significance compared to the control using 1-way ANOVA followed by Dunnett's multiple comparisons test.


In order to study the modulation on host signalling by this co-stimulation model, 16HBE cells were treated with *B. cenocepacia* J2315 (50 MOI) and 25% (v/v) MSSA NCTC 6571 supernatant for 5 minutes (Figure 5.3A and B). J2315 caused a marginal downregulation on phosphorylation of p38 and Erk, whilst *S. aureus* supernatant caused a rapid phosphorylation, particularly with Erk. Co-stimulation led to activation of p38 and Erk, which likely results from the effect of *S. aureus*. Following 1 hour pretreatment of *S. aureus* supernatant (25% v/v), 16HBE cells were infected with J2315 (50 MOI) for 5 minutes (Figure 5.3C, D and E). J2315 and *S. aureus* induced a marginal increase in phosphorylation of p38, whilst dramatically increasing phosphorylation of Erk and I $\kappa$ B $\alpha$ , which indicates *S. aureus* supernatant at a higher concentration is able to induce persistent phosphorylation of Erk and I $\kappa$ B $\alpha$ . However, when co-stimulation extended to 2 hours some trends became visible (Figure 5.4), although not achieving statistical significance. *B. cenocepacia* J2315 caused activation of Akt, p38, Erk and p65, whereas *S. aureus* did not change any tested signalling. It is worth noting that unlike MRSA252 supernatant, MSSA NCTC 6571 supernatant abolished phosphorylation of p38, Erk and p65 caused by *B. cenocepacia* J2315. Therefore, these findings suggest that *S. aureus* supernatants may suppress the pro-inflammatory effect of *B. cenocepacia* by blocking p38, Erk and NF- $\kappa$ B.





**Figure 5.3** MAPK response to co-stimulation with *B. cenocepacia* J2315 and MSSA NCTC 6571 supernatant.

(A-B) 16HBE cells were treated with LB (25% v/v) as control, J2315 (50 MOI), *S. aureus* supernatant (25% v/v) for 5 minutes. (C-E) After 1 hour pretreatment with *S. aureus* supernatant (25% v/v), 16HBE cells were infected with J2315 (50 MOI). The control was treated with 25% (v/v) LB for 1 hour. Phosphorylation of p38, Erk and IκBα of cell lysates were measured using western blot and normalised against total p38, Erk and IκBα, respectively. Data are shown as mean ± SEM from three independent experiments. \*,  $p < 0.05$  compared with the control using one-way repeated measures ANOVA following Dunnett's multiple comparisons test.




**Figure 5.4** PI3K, MAPK and NF- $\kappa$ B response to co-stimulation with *B. cenocepacia* J2315 and MRSA252 or MSSA NCTC 6571 supernatant.

16HBE cells were treated with LB (10% v/v) as the control, J2315 (5 MOI), with the presence or absence of *S. aureus* supernatant (10% v/v) for 2 hours. Phosphorylation of Akt (**A**), GSK-3 $\beta$  (**B**), p38 (**C**), Erk (**D**) and p65 (**E**) of cell lysates were measured using western blot and normalised against total p38. (**F**) Western blot analysis for levels of phosphorylated Akt, phosphorylated GSK-3 $\beta$ , phosphorylated p38, phosphorylated Erk, phosphorylated p65, and p38 in cell lysate following *S. aureus* supernatants and *B. cenocepacia* J2315 co-stimulation. Protein bands representative of three independent experiments. Data are shown as mean  $\pm$  SEM. \*,  $p < 0.05$  compared with the control using one-way repeated measures ANOVA following Dunnett's multiple comparisons test.

### 5.3 Characterisation of anti-inflammatory factors in *S. aureus* supernatants

It has been confirmed that the anti-inflammatory factors in *S. aureus* supernatant are heat-stable and not associated with cell wall structures (Figure 5.1E-F). To determine the nature of the effectors, *S. aureus* supernatants were digested with the broad-spectrum serine protease, PK or heat-inactivated PK (Figure 5.5A). Co-stimulation with *B. cenocepacia* J2315 and PK-treated supernatants did not restore IL-8 production to the level as J2315-treated group; on the contrary, IL-8 production dropped to the same level of the control. This indicates that the effectors are not proteinaceous. To assess the approximate molecular weight, *S. aureus* supernatants were separated into different molecular cut-off fractions (Figure 5.5B). Although there was a weak positive control response in this series of experiments, supernatants <3 kDa still abolished IL-8 production back to the same level as the control, which indicates the anti-inflammatory molecules are small. In order to elucidate whether TLR is exclusively involved in *S. aureus*-induced anti-inflammatory response, 16HBE cells were stimulated with up to 10 ng/mL IL-1 $\beta$ . However, IL-1 $\beta$  only induced half the amount of IL-8 induced by *B. cenocepacia* J2315. In addition, IL-1 $\beta$  / *S. aureus* supernatants co-stimulation caused a marginal increase rather than a decrease (Figure 5.5C). Therefore, a further experiment was done by co-stimulating with IL-1 $\beta$  and TNF $\alpha$ , which resulted in a similar level of IL-8 compared to J2315-treated group (Figure 5.5D). However, no effect of *S. aureus* supernatants was observed on IL-1 $\beta$  / TNF $\alpha$ -induced IL-8 production, indicating IL-1R and TNFR are not associated with this process and the suppressive effect does not appear to be a non-selective downregulation of IL-8 production.

Taken together, the anti-inflammatory molecules in *S. aureus* supernatants are characterised as small, heat-resistant, non-proteinaceous, non-cell wall components.



**Figure 5.5 Characterisation of anti-inflammatory effectors in *S. aureus* supernatants.**

(A) *S. aureus* supernatants were treated with immobilised proteinase K for 1 hour at 37°C, or treated with heat-inactivated proteinase K (99°C for 1 hour) as the positive control. 16HBE cells were co-stimulated with *B. cenocepacia* J2315 (5 MOI) and proteinase K-treated *S. aureus* supernatants (10% v/v) for 24 hours.


(B) *S. aureus* supernatants were filtered through selective membranes. 16HBE cells were treated with *B. cenocepacia* J2315 (5 MOI) and *S. aureus* supernatant fractions containing metabolites of >10 kDa, >3 kDa and <3 kDa for 24 hours. IL-8 release was measured by ELISA. Data represent mean  $\pm$  SEM from 3 independent experiments performed in duplicate. Significance compared to J2315-treated group using 2-way ANOVA followed by Dunnett's multiple comparisons test.

(C) After 1 hour pretreatment of *S. aureus* supernatants (10% v/v), 16HBE cells were stimulated with increasing dose of IL-1 $\beta$  for 24 hours. Data represent mean  $\pm$  SEM from 3 independent experiments performed in duplicate. Significance was compared to the control using 2-way ANOVA followed by Dunnett's multiple comparisons test.

(D) 16HBE cells were co-stimulated with IL-1 $\beta$  (1 ng/mL) / TNF $\alpha$  (10 ng/mL) and *S. aureus* supernatants (10% v/v) for 24 hours. *B. cenocepacia* J2315 (5 MOI) was the positive control. Data represent mean  $\pm$  SEM from 3 independent experiments performed in duplicate. \*,  $p < 0.05$  compared to IL-1 $\beta$  / TNF $\alpha$ -treated group or J2315-treated group using 2-way ANOVA followed by Dunnett's multiple comparisons test.

#### 5.4 16HBE cells are hypo-responsive to Pam<sub>3</sub>CSK4, LPS but not flagellin

As a Gram-negative bacterium with flagella, *B. cenocepacia* J2315 potentially binds to and activates TLR2, TLR4 and TLR5 (de C Ventura et al., 2008; West et al., 2008). To determine which TLR is involved in *B. cenocepacia* / *S. aureus* polymicrobial infection setting, 16HBE cells were co-stimulated with TLR2, TLR4 and TLR5 agonists, Pam<sub>3</sub>CSK4, LPS and flagellin, respectively, and with the presence or absence of MSSA NCTC 6571 supernatant (Figure 5.6). None of the agonists induced IL-8 production as well as J2315, and only flagellin triggered a dose-dependent, significantly upregulated IL-8 response, indicating the pro-inflammatory property of J2315 is associated with multiple virulence factors. Moreover, with all agonists a marginal drop of the IL-8 response was seen with the presence of *S. aureus* supernatant.




**Figure 5.6 Pro-inflammatory response to TLR2, 4, or 5 agonists and *S. aureus* supernatant.**

16HBE cells were co-stimulated with increasing dose of Pam<sub>3</sub>CSK4, LPS, flagellin and MSSA NCTC 6571 supernatant (10% v/v) for 24 hours. *B. cenocepacia* J2315 (5 MOI) was the positive control. IL-8 release was measured by ELISA. Data represent mean  $\pm$  SEM from 3 independent experiments performed in duplicate. \*,  $p < 0.05$  compared to control using 2-way ANOVA followed by Dunnett's multiple comparisons test.


To investigate host signalling responses to Pam<sub>3</sub>CSK4 and flagellin, the time course of the response was assessed. No effect was seen with Pam<sub>3</sub>CSK4 (10 µg/mL) up to 60 minutes (Figure 5.7), whereas flagellin (1 µg/mL) caused a transient phosphorylation of Akt, GSK-3β, p38, Erk and p65 between 15 and 30 minutes (Figure 5.8), which is consistent with the IL-8 response. This indicates that flagellin might play a major role in the *B. cenocepacia* induced pro-inflammatory response.

Following pretreatment with the TLR4 antagonist LPS-RS, stimulation with J2315 showed significant downregulation in IL-8 production (14% decrease compared to the J2315-treated group), to the same level as *S. aureus* supernatants (Figure 5.9A). In contrast, co-stimulation with TLR5 antagonist hTLR5fc and J2315 did not have any effect compared to the J2315-treated group (Figure 5.9B).



**Figure 5.7 Time course of Pam<sub>3</sub>CSK4 on PI3K, MAPK and NF-κB activation.**


16HBE cells were treated with Pam<sub>3</sub>CSK4 (10 µg/mL) for 5-60 minutes, TNFα (20 ng/mL, 5 minutes) as the positive control. Phosphorylation of Akt (**A**), GSK-3β (**B**), p38 (**C**), Erk (**D**) and p65 (**E**) of cell lysates were measured using western blot and normalised against total p38. (**F**) Western blot analysis for levels of phosphorylated Akt, phosphorylated GSK-3β, phosphorylated p38, phosphorylated Erk, phosphorylated p65, and p38 in cell lysate following Pam<sub>3</sub>CSK4 stimulation. Protein bands representative of three independent experiments. Data are shown as mean ± SEM. \*,  $p < 0.05$  compared with control using one-way repeated measures ANOVA following Dunnett's multiple comparisons test.



**Figure 5.8 Time course of flagellin on PI3K, MAPK and NF-κB activation.**

16HBE cells were treated with flagellin (1 µg/mL) for 5-180 minutes, TNFα (20 ng/mL, 5 minutes) as the positive control. Phosphorylation of Akt (**A**), GSK-3β (**B**), p38 (**C**), Erk (**D**) and p65 (**E**) of cell lysates were measured using western blot and normalised against total p38. (**F**) Western blot analysis for levels of phosphorylated Akt, phosphorylated GSK-3β, phosphorylated p38, phosphorylated Erk, phosphorylated p65, and p38 in cell lysate following flagellin stimulation. Data are shown as mean ± SEM from three independent experiments. \*,  $p < 0.05$  compared with control using one-way repeated measures ANOVA following Dunnett's multiple comparisons test.





**Figure 5.9 TLR4 antagonist LPS-RS inhibits *B. cenocepacia* J2315-induced IL-8 production but not TLR5 antagonist hTLR5 fc.**

**(A)** Following pretreatment with TLR4 antagonist LPS-RS (2 µg/mL) for 1 hour, 16HBE cells were stimulated with LPS (10 µg/mL) or *B. cenocepacia* J2315 (5 MOI) for 24 hours. Cells were also co-stimulated with *S. aureus* supernatants (10% v/v, 1 hour pretreatment) and *B. cenocepacia* J2315 (5 MOI) for 24 hours as positive controls. **(B)** After 24 hours pretreatment with TLR5 antagonist hTLR5-fc (100 ng/mL), 16HBE cells were stimulated with TLR5 agonist flagellin (1 µg/mL) or *B. cenocepacia* J2315 (5 MOI) for 24 hours. Cells were co-stimulated with *S. aureus* supernatants (10% v/v, 1 hour pretreatment) and *B. cenocepacia* J2315 (5 MOI) for 24 hours as positive controls. IL-8 release was measured by ELISA. Data represent mean ± SEM from 3 independent experiments performed in duplicate. \*,  $p < 0.05$  compared to *B. cenocepacia*-J2315 treated group using 2-way ANOVA followed by Dunnett's multiple comparisons test.

## 5.5 Discussion

Several lines of evidence indicate that *S. aureus* induces an anti-inflammatory response to facilitate persistence inside the host. For example, the secreted protein SSL3 blocks TLR2 activity and thereby inhibits IL-8 production *in vitro* (Bardoel et al., 2012), as well as antagonising TLR2 function as a virulence factor during *in vivo* infection of mice (Koymans et al., 2017). Additionally, *S. aureus*-induced IL-10 production enhances persistence by hindering local pro-inflammatory responses in an acute murine skin infection model (Leech et al., 2017). Because of the large reservoir of immunomodulatory effectors (Thammavongsa et al., 2015), multi-drug resistance, and biofilm formation, *S. aureus* is a major threat to human health in infectious diseases.

Extensive studies have focused on the interactions between *S. aureus* and *P. aeruginosa* as they are two prevalent pathogens involved in CF lung infections. However, *S. aureus*/Bcc interactions and their impact on host response have not been assessed before. Herein, we demonstrate that in the context of *S. aureus* / *B. cenocepacia* co-infection, supernatants from *S. aureus* isolates dampen *B. cenocepacia* live bacteria-induced IL-8 production in airway epithelial cells, indicating that *S. aureus* releases anti-inflammatory compounds into the broth. Interestingly, all the supernatants from *S. aureus* isolates block J2315-induced IL-8 production to different levels ( $p < 0.05$ ). Hyperinflammatory isolates such as SH1000, RN4220 and USA400, where the live bacteria induced strong IL-8 production (Figure 3.3A), exhibit inhibitory effect on IL-8 production as well. This indicates the components may be widely expressed in *S. aureus*.

In agreement with our findings, Chekabab et al. (2015) demonstrated that *S. aureus* supernatants suppress *P. aeruginosa* supernatant-induced cytokine (IL-8 and CXCL-2) production through inhibiting TLR1/2 and TLR4, but not TLR5. However, the inhibition effect is medium-specific, as it presents with LB-cultured supernatant but not TSB-cultured supernatant, which is in parallel with our results. LB and TSB are rich media, commonly used to culture aerobic microorganisms such as *S. aureus*. Compared to TSB, LB has simple ingredients and is not buffered. Unfortunately, we are unable to confirm which medium reflects the physiological condition of the airway milieu. Several studies have applied synthetic cystic fibrosis sputum medium (Kumar and Cardona, 2016) or artificial sputum medium (Fung et al., 2010) to resemble the sputum from CF patients and observed changes in gene expression responsible for virulence. Since differences in compositions of culture media may change metabolism and subsequently affect modulation of metabolites on host response, *S. aureus* should be cultured in a medium that reflects the nutrition of the infected airway. Nevertheless, *S. aureus* grows poorly in

synthetic CF sputum medium and thus is not possible to compare the differences of *S. aureus* supernatant cultured in synthetic CF sputum medium and LB (Bernier et al., 2016).

*S. aureus* biofilm supernatants are able to impair macrophage phagocytosis (Hanke and Kielian, 2012) and *S. aureus* biofilms programme macrophages to the alternative activated M2 phenotype (Thurlow et al., 2011). It was unexpected that in our results *S. aureus* biofilm supernatants had very little effect on J2315-induced IL-8, except a marginal downregulation by MSSA NCTC 6571 biofilm supernatant ( $p < 0.05$ ). Supernatants from planktonic and biofilm *S. aureus* contain different features (such as proteases in biofilm supernatant) which result in different signalling activation and cytokine production. Compared to the planktonic counterparts, biofilm supernatants cause apoptosis, downregulate phosphorylation of p38 and JNK, upregulate phosphorylation of Erk, and induce higher level of cytokines at 4 hours but lower level at 24 hours post-stimulation in human keratinocytes (Secor et al., 2011). Differential host responses directed towards planktonic and biofilm supernatants indicate that *S. aureus* modifies its expression of virulence factors at different stages of infection. Brady et al. (2018) developed epicutaneous (planktonic) and subcutaneous mesh (biofilm) *S. aureus* infection models in mice and compared cytokine production at 7 days and 21 days. Although the day 7 biofilm model caused higher levels of CXCL1, CXCL2, IL-1 $\alpha$  and IL17A production compared to the planktonic counterpart, the host response was ineffective and insufficient for microbial clearance. On day 21, *S. aureus* biofilm shifted to a dormant state, which results in *S. aureus* being less detectable to the immune system as shown by a decrease in cytokines released (Brady et al., 2018).

In further experiments, the effects of heat-treatment, heat-killed *S. aureus*, proteinase K-treatment, and different molecular weight cut-off fractions were examined in order to determine the nature of the anti-inflammatory compounds in *S. aureus* supernatants. The compounds were characterised as small (<3 kDa), heat-resistant, non-proteinaceous, and non-cell wall components, in agreement with previous work (Chekabab et al., 2015). When co-stimulated with J2315, IL-8 production elicited by proteinase K-treated *S. aureus* supernatants showed a 50% decrease compared to deactivated proteinase K-treated *S. aureus* supernatants. This indicates that the anti-inflammatory components is less likely to be peptides or proteins. The decrease may result from pro-inflammatory components in *S. aureus* supernatants spontaneously degraded by proteinase K. Apart from releasing immunosuppression modulators, *S. aureus* produces multiple virulence factors that have pro-inflammatory effects. For example, MRSA-derived small, heat- and protease-resistant molecules induce nitric oxide in human sinonasal epithelial cells (Carey et al., 2015). The level of nitric oxide and inflammation status are positively correlated as nitric oxide plays

a critical role in microbial killing, wound healing as well as facilitating chemokine production (Trifilieff et al., 2000). In contrast with our observations, *S. aureus* supernatants are reported to induce a stronger magnitude of cytokine, chemokine and prostaglandin E<sub>2</sub> expression than live bacteria (Moreilhon et al., 2005). These findings suggest that *S. aureus* as a prevalent pathogen modulates the immune response to benefit colonisation and infections.

Inhibition of IL-8 production by *S. aureus* supernatants has been previously reported in TNF $\alpha$ -treated HUVEC cells (Tajima et al., 2006). The inhibitory effect of supernatants is also abolished by proteinase K or trypsin treatment but, in contrast with our findings, is sustained after heat-treatment. Thus, the anti-inflammatory component is defined as heat-stable protein. Of note, *S. aureus* supernatants are treated with proteinase K and trypsin, respectively; however, they did not describe whether the final supernatants contain proteinase K. Serine proteases such as trypsin activate proteinase-activated receptor-2 which subsequently activates NF- $\kappa$ B (Buddenkotte et al., 2005). Although there is no evidence showing that proteinase K causes activation of NF- $\kappa$ B, it may be that stimulating HUVEC cells with residual proteinase K (10  $\mu$ g/mL according to their method) initiated an inflammatory response that masked the *S. aureus* inhibitory activity.

Immunosuppression is also observed with commensal *S. salivarius*, which dwells in the oral and nasopharyngeal cavity. *S. salivarius* secretes small heat-resistant proteins which hinder p65 nuclear translocation and subsequently IL-8 production, both under baseline condition and on co-stimulation with different stimuli (TNF $\alpha$ , IL-1 $\beta$ , *P. aeruginosa*, flagellin, LL-37) (Cosseau et al., 2008; Kaci et al., 2014; 2011). The abilities of commensal bacteria to cause a low pro-inflammatory response, induce an anti-inflammatory response and regulate gene expression involved in airway epithelial tight junctions contribute to immune tolerance and homeostasis in the mucosal barrier. Our results showed that the inhibitory effect of *S. aureus* secreted products was TLR-specific, as *S. aureus* supernatants did not affect IL-1 $\beta$ /TNF $\alpha$ -induced IL-8 production. In contrast, *S. aureus* supernatants suppress TNF $\alpha$ -induced IL-8 production in the HUVEC model. However, the supernatants are obtained using a different method (incubated for 1 hour in mammalian cell culture medium) which may elicit different metabolites (Tajima et al., 2006). Overall, *S. aureus* may induce anti-inflammatory responses to prevent phagocytosis and hinder the adaptive immune response. Thus, it is critical to elucidate the immune evasion mechanisms used by *S. aureus* to design novel treatments.

Following co-stimulation with *B. cenocepacia* J2315 and *S. aureus* supernatant, we assessed activation of PI3K, MAPK and NF- $\kappa$ B and found that MSSA supernatant

abolished phosphorylation of p38, Erk and p65 caused by J2315. MRSA252 supernatant did not affect either baseline or J2315-induced phosphorylation, which correlates with its less downregulation effect on IL-8 production compared to MSSA NCTC 6571. Our results are in partial agreement with the results by Chekabab et al., (2015) as they showed *S. aureus* supernatants block NF- $\kappa$ B activation but not p38. However, Chekabab et al. (2015) stimulated airway epithelial Beas-2B cells with synthetic TLR2/1 agonist Pam<sub>3</sub>Csk4, whereas *B. cenocepacia* contains more complicated ligands. Additionally, the downstream signalling cascades engaged in the anti-inflammatory response may be cell type-specific. Nasal *S. aureus* isolates utilise PI3K and Erk signalling cascades to induce IL-10 production whereas they induce TNF $\alpha$  in a p38-dependent pathway in human PBMC (Peres et al., 2015). This suggests that through expressing unique ligands in the process of infection, *S. aureus* manipulates different signalling cascades to initiate immune tolerance or phagocytosis.

Chekabab et al. (2015) suggest that the inhibitory effect of *S. aureus* supernatant is through TLR signalling. In order to determine which TLR signalling was modulated by *S. aureus*-secreted products, first we stimulated 16HBE cells with TLR2/1, TLR4 and TLR5 agonists, respectively. Pam<sub>3</sub>CSK4, LPS and flagellin caused 22%, 62% and 151% increases in IL-8 production at the highest dose (10  $\mu$ g/mL) compared to the control. The result was unexpected, as in Beas-2B cells Pam<sub>3</sub>CSK4 was found to be the most efficacious IL-8 inducer, LPS showed a dose-dependent manner of inducing IL-8 and flagellin did not affect IL-8 production (Chekabab et al., 2015). The hypo-responsiveness to TLR2/1 and TLR4 agonists may result from low expression and location of the receptors and the accessory molecules, as in polarised primary human airway epithelial cells TLR2 is mainly expressed on the basolateral surface and TLR4 is weakly expressed on the apical surface (Ioannidis et al., 2013). For example, airway epithelial Beas-2B cells display lower sensitivity to *S. aureus* and TLR2 agonists compared to Gram-negative bacteria in terms of IL-8 production, which is associated with low expression of TLR2 and missing expression of the co-receptor CD36; whereas monocytes exhibit a robust TNF $\alpha$  response to both Gram positive and negative bacteria (Mayer et al., 2007). This indicates that mucosal epithelia may use this mechanism to prevent an excessive inflammatory response and maintain homeostasis. In contrast, *B. cenocepacia* J2315 can induce TLR2 and TLR4, but not TLR5 expression in Beas-2B cells, whereas only TLR5 is required for a pro-inflammatory response (de C Ventura et al., 2008).

Further we assessed the effect of Pam<sub>3</sub>CSK4 and flagellin on signalling cascades. It is not surprising that Pam<sub>3</sub>CSK4 did not affect PI3K, MAPK and NF- $\kappa$ B. In agreement with our result, Pam<sub>3</sub>CSK4 is able to trigger marginal phosphorylation of p38 but not NF- $\kappa$ B

activity and IL-8 production in human airway epithelial cells (Winder et al., 2009). Flagellin has been reported to activate phosphorylation of p38 and p65 in airway epithelial Calu-3 cells (Illek et al., 2008), phosphorylation of Akt in human colonic epithelial cells (Sang et al., 2006), and phosphorylation of p38, Erk, JNK and I $\kappa$ B $\alpha$  in mouse bone marrow-derived dendritic cells (Feuillet et al., 2006). In our system, airway epithelial cells were most sensitive to flagellin, as flagellin activated several signalling pathways and induced a strong IL-8 response. In the following experiment, we pretreated the cells with TLR4 and TLR5 antagonists, respectively, followed by stimulation with TLR agonists or J2315. Compared to J2315-treated cells, only the TLR4 antagonist caused a decrease in IL-8 production. However, the magnitude of reduction was similar to *S. aureus* supernatants. This suggests that *S. aureus*-secreted products may block TLR4 and subsequently suppress the *B. cenocepacia*-induced pro-inflammatory response. Recently, a family of small (~20-25 amino acids) secreted peptides, namely PSMs are able to bind to TLR4 without activation in THP-1 cells (Chu et al., 2018). PSM  $\alpha$ 1- $\alpha$ 3 suppress pro-inflammatory cytokines production induced by high mobility group box-1 protein (belongs to the danger-associated molecular pattern family) by subverting TLR4 signalling. However, although heat stability of these factors is unclear, according to our results the anti-inflammatory factors may not be peptide/protein. Thus, further characterisation needs to identify the Staphylococcal factors.

## 5.6 Conclusions

In this chapter, we demonstrate that *S. aureus* isolates produce immunomodulatory factors that subvert the *B. cenocepacia*-induced inflammatory response in airway epithelial cells. The factors are characterised as small, heat-stable, non-proteinaceous, non-cell wall components. Additionally, we demonstrate *S. aureus* subverts the *B. cenocepacia*-induced inflammatory response by attenuation of TLR4-mediated p38, Erk and NF- $\kappa$ B signalling cascades. Further experiments are needed to identify the active factors.

## **Chapter 6 General discussion**

## 6.1 Host responses to monomicrobial infections

The innate immunity protects the lower respiratory tract and controls its homeostasis. Intensity and duration of the immune response, along with pathogenesis of the pathogens, determine the clinical outcome (Strobel et al., 2016). The host-pathogen interactions in the scenario of polymicrobial infections have been acknowledged with increasing frequency, yet still require intensive investigation (Bragonzi et al., 2012; Dai et al., 2018; Flight et al., 2015; Sibley et al., 2008). This thesis began by assessing pathogenesis of individual bacterial pathogens that belong to *S. aureus* and Bcc species, and the effect on host signalling cascades and pro-inflammatory responses. Upon recognition of incoming microorganisms, certain signalling pathways are activated to initiate the expression of pro-inflammatory cytokines and chemokines as key events to conduct and amplify the immune response (Weitnauer et al., 2016; Whitsett and Alenghat, 2015).

To enhance the potential of pathogenesis, multiple virulence factors are required at different stages of infection (Foster et al., 2014; Kim et al., 2010). Thus, microbial invasion was assessed by viable count using airway epithelial cells grown on plastic. Airway epithelial cells are vulnerable to *S. aureus* invasion, and might be killed by virulent isolates (Strobel et al., 2016). We observed that epithelial cells accumulated higher amounts of *S. aureus* on the cell membrane compared to clinical Bcc isolates, whereas the latter was taken up more readily. To assess invasion on intact polarised epithelial monolayers, airway epithelial cells were cultured on transmembrane inserts, infected apically with live bacteria, and the integrity was monitored using TEER. In agreement with a previous study, our isolates of interest, MRSA252 and *B. multivorans* 13243 both breached polarised epithelial monolayers in 24 h. By disturbing the actin cytoskeleton, *S. aureus* or *B. multivorans* invade the epithelium barrier (Schwab et al., 2002; Soong et al., 2011). *B. multivorans* also leads to extensive cell death (Schwab et al., 2002), which might explain why *B. multivorans* caused higher level of FITC-dextran flux compared to *S. aureus*. This indicates that *B. multivorans* resulted in a higher barrier permeability than *S. aureus*.

As PI3K regulates cytoskeleton rearrangement (Gessain et al., 2015), we assessed invasion by treating airway epithelial cells with pharmacological PI3K inhibitors prior to live bacteria exposure. In line with a previous study (Kierbel et al., 2005), PI3K is required for *S. aureus* internalisation. In addition, because of the asymmetric localisation of TLRs on the apical and basolateral surfaces (Ioannidis et al., 2013), an invasion assay using polarised epithelial monolayers would closer reflect the physiological condition. This is further supported by the fact that *P. aeruginosa* binds to distinct receptors on the apical and basolateral surfaces, with different binding affinity (Bucior et al., 2010). Hence, a



future experiment is guaranteed to perform the invasion assay using epithelial cells grown on permeable membranes, and comparing invasion by infecting the monolayers from the apical and the basolateral surface.

As bacteria in biofilms are more resistant to antimicrobial substances and host clearance, biofilm formation is a key virulence factor for persistence and immune evasion. The strength of biofilm development by *S. aureus* is positively correlated with disease severity (Domenico et al., 2018). Both *S. aureus* and Bcc are known to form biofilm in close proximity to the apical mucosal surface (Krismer and Peschel, 2011; Schwab et al., 2002). Our results showed that *S. aureus* and Bcc developed differential amounts of biofilms. Some clinical isolates such as *B. multivorans* 13243, *B. cenocepacia* J2315, and MSSA NCTC 10788 formed more robust biofilms compared to non-invasive strains.

Heterogenicity was also observed in IL-8 production induced by *S. aureus* isolates. IL-8 is a biomarker for monitoring infections (Kraft et al., 2015). As the main receptor for recognising *S. aureus*, TLR2 activation depends on ligand abundance on the bacterial surface. Hence, differential activation of TLR2-mediated signalling by *S. aureus* clinical isolates may result from differences in bacterial metabolic activity, capsule expression and cell wall synthesis (expression of lipoproteins and SpA) (Hilmi et al., 2014; Peres et al., 2015). Bcc has prototypical ligands for TLR2, TLR4, and TLR5 (West et al., 2009). CF-related isolates *B. cenocepacia* J2315 and *B. multivorans* 13243 have been reported to induce high amounts of IL-8 in 16HBE cells, with a higher level induced by J2315 (Kaza et al., 2010). However, in our system, *B. cenocepacia* J2315 but not *B. multivorans* 13243 caused a significant increase in IL-8 production; thus, *B. cenocepacia* J2315 was chosen for further experiments.

The quantification of the IL-8 protein level was 24 h post-bacterial stimulation. However, the duration should be carefully decided, as protein levels are highly dynamic and regulated by mRNA level, translation, and degradation. Protein levels at steady state are dominated by mRNA levels, whereas under stress conditions, in addition to transcription changes, a delay occurs between changes in transcription and translation level, especially immune response proteins (Jovanovic et al., 2015; Liu et al., 2016). Hence, a time course of IL-8 protein level would be useful to decide the duration of bacterial stimulation. By doing so, the gap between different treatments might be larger.

IL-8 production is initiated by TLR-mediated downstream signalling cascades (Lim and Staudt, 2013). To elucidate which pathways are modulated by *S. aureus* and Bcc, phosphorylation of key signalling proteins following bacterial stimulation was investigated using western blot. In agreement with Gillette et al. (2013), *B. cenocepacia* J2315 led to

phosphorylation of p38, Erk and p65 NF- $\kappa$ B, whereas phosphorylation maintained the same when exposed to Staphylococcal strains and their supernatants. The minimal expression of TLR2 co-receptor CD36 restricts recognition of Gram-positive bacteria, which leads to airway epithelial cells hypo-responsive to Gram-positive bacteria but not Gram-negative bacteria (Mayer et al., 2007). On the other hand, p38 and Erk contribute to IL-8 mRNA stability in lung epithelial cells, supporting the observation that J2315 induces higher level of IL-8 compared to *S. aureus* isolates (Bhattacharyya et al., 2011).

## **6.2 Host response to polymicrobial infections and the comparison with monomicrobial infections**

*B. cenocepacia* J2315 is strong IL-8 inducer on its own, and in a chronic *P. aeruginosa*/*B. cenocepacia* co-infection model, significant higher amounts of cytokines and chemokines were induced by co-infection compared to single infections (Bragonzi et al., 2012). The net impact of polymicrobial infections leads to different host responses, and the consequences may be detrimental, insignificant or beneficial to the host (McArdle et al., 2018; Ramsey and Whiteley, 2009; Sibley et al., 2008). To elucidate the host response to *S. aureus*/*B. cenocepacia* co-infection, 2 isolates of *S. aureus* that downregulated IL-8 production without causing significant cell damage were chosen and supernatants from overnight cultures were prepared.

The data presented here is the first time to show the anti-inflammatory effect of *S. aureus* products when co-stimulated with live *B. cenocepacia* bacteria. Similar results were observed when Staphylococcal supernatants co-stimulated bronchial epithelial cells with supernatant of a close relative of *Burkholderia*, *P. aeruginosa* (Chekabab et al., 2015). Chekabab et al. (2015) demonstrated that the synthetic ligand for TLR1/2 induced IL-8 to the same level of as *P. aeruginosa* supernatant, and the response was abolished by *S. aureus* supernatant, whereas Beas-2B cells were hypo responsive to LPS or flagella. Further experiments confirmed that *S. aureus* supernatant blocked TLR2-mediated NF- $\kappa$ B signalling but not MAPK (Chekabab et al., 2015). To demonstrate which signalling pathway is involved in our system, we co-treated airway epithelial cells with *S. aureus* supernatant and *B. cenocepacia* J2315, followed by the assessment of phosphorylation using western blotting. Although not achieving statistical significance, *S. aureus* supernatant abolished phosphorylation of p38, Erk and p65 NF- $\kappa$ B. This is the first time showing the inhibitory effect of *S. aureus* on MAPK signalling.

The findings in this thesis demonstrate the antagonistic interaction between *S. aureus* and *B. cenocepacia* on inducing the pro-inflammatory response. By inhibiting MAPK and NF-

$\kappa$ B signalling cascades activated by *B. cenocepacia*, *S. aureus*-secreted products reduce IL-8 production in airway epithelial cells. This might shed light on a novel Staphylococcal immune evasion mechanism through inducing anti-inflammatory responses. Further, identification of the host target might provide a novel treatment for chronic inflammation resulted from excessive immune responses.

### **6.3 Interpretation of the host receptor for *S. aureus* anti-inflammatory factor(s)**

TLRs are the major sensors for *B. cenocepacia*, and TLR2 is the most important PRR to detect *S. aureus* (Askarian et al., 2018). Chekabab et al. (2015) reported that *P. aeruginosa* activated TLR2 is blocked by *S. aureus* which subsequently subverts TLR2-mediated NF- $\kappa$ B signalling. To distinguish whether the effect was TLR-specific, upon stimulation with IL-1 $\beta$  and TNF $\alpha$ , airway epithelial cells were treated with *S. aureus* supernatants. In contrast, while IL-1 $\beta$  / TNF $\alpha$  induced the same level of IL-8 as *B. cenocepacia* J2315, *S. aureus* supernatants did not abolish IL-8 production. Hence, the anti-inflammatory effect of *S. aureus* is TLR-specific.

To illustrate which TLR is targeted by *S. aureus*, airway epithelial cells were exposed to *S. aureus* supernatants, together with TLR2/1, TLR4 and TLR5 agonists, respectively. However, although flagellin led to a significant increase in IL-8 production, none of the other TLR agonists elicited IL-8 to the same level as *B. cenocepacia* J2315 even at a high concentration. In addition, only flagellin significantly upregulated PI3K/Akt, MAPK and NF- $\kappa$ B signalling cascades in 16HBE cells. This is consistent with a previous study that only TLR5 but not TLR2 and TLR4 plays a central role in *B. cenocepacia*-induced pro-inflammatory responses (de C Ventura et al., 2008).

The hypo-sensitivity to TLR ligands have three possible explanations. The commercially purchased TLR ligands were isolated from other microorganisms. The acylation of LPS lipid A contributes to abnormal pro-inflammatory responses to whole-cell lysates and highly purified LPS from *B. cenocepacia* (De Soyza et al., 2004; Zughaier et al., 1999). Hence, stimulation with isolated TLR ligands from *B. cenocepacia* J2315 would closely reflect the host response to *B. cenocepacia*. Secondly, the hypo-responsiveness might result from restricted expression of TLRs. The epithelial cell line 16HBE exhibits a low but inducible level of TLR2 expression and a relatively higher level of TLR4 (Bailey et al., 2009; Pace et al., 2008), which explains hypo-responsiveness to Gram-positive bacteria compared to Gram-negative bacteria (Mayer et al., 2007). In addition, epithelial cells such as BEAS-2B are deficient in TLR2 co-receptor CD36, and TLR2/CD36 transfection results

in enhanced IL-8 production (Mayer et al., 2007). TLR5 is also present in 16HBE cells (Muir et al., 2004; Parker and Prince, 2013). Lastly, the differential responses to *B. cenocepacia* and TLR ligands might result from the additional receptors that are targeted by *B. cenocepacia* (Ganesan and Sajjan, 2011). For example, the O antigen of LPS from *B. cenocepacia* activates the inflammasome in a TLR4- and caspase-1-dependent manner (Kotrange et al., 2011). In addition, *B. cenocepacia* binds to TNFR1 and subsequently elicits robust IL-8 expression through an unknown ligand in CF airway epithelial cells (Sajjan et al., 2008). Hence, in addition to TLR activation, other host receptors might be involved in the disproportionate pro-inflammatory response to *B. cenocepacia*.

Further experiments were performed with TLR antagonists. TLR4 but not TLR5 inhibitor reduced IL-8 production elicited by *B. cenocepacia* to the same level of *S. aureus* supernatants. Multiple Staphylococcal products are known to antagonise TLRs, such as TLR2 by SSL3 (Bardoel et al., 2012). The anti-inflammatory molecules present in Staphylococcal supernatant might be PSM $\alpha$ , which consists of 20 to 30 amino acids with about 2 to 3 kDa peptides (Berube et al., 2014). PSM $\alpha$ 1- $\alpha$ 3 of *S. aureus* have shown to disrupt the interaction between TLR4 and high motility group box-1 protein, and subsequently inhibit NF- $\kappa$ B signalling in HEK-Blue hTLR4 cells (Chu et al., 2018). However, PSM $\alpha$  is able to activate TLR2 signalling (Armbruster et al., 2016) and HEK-Blue hTLR4 cells (InvivoGen, USA) are inactive to TLR2 ligands. What would be the net impact of PSM $\alpha$  if cells with the presence of TLR2 and TLR4 were used?

In our system, the downregulation on IL-8 production by blocking TLR4 was minimal but significant. Alternatively, following confirmation of TLR isoforms expression in airway epithelial cells, we could compare chemokine responses to *B. cenocepacia* using wild type cells and TLR isoform knockdown cells. By doing so, it would confer an unbiased actual response showing the net impact of Staphylococcal products. In addition, using live bacteria would reflect the actual ligands shed from intact bacteria compared to using TLR agonists.

As TLR4 is the only TLRs that requires both MyD88 and TRIF, this leads to a hypothesis that the anti-inflammatory effect of *S. aureus* results from blocking MyD88- and TRIF-dependent signalling. So far, YopJ from *Yersinia pseudotuberculosis* is the only bacterial effector that blocks type I IFN and cytokine responses through suppressing MyD88 and TRIF in DC and macrophages (Rosadini et al., 2015; Rosadini and Kagan, 2015). Hence, by knocking out MyD88 and TRIF, respectively, it could be possible to evaluate the role of MyD88 and TRIF when airway epithelial cells are co-stimulated with *B. cenocepacia* and *S. aureus* supernatants.

## 6.4 Characterisation of *S. aureus* immunomodulatory factor(s)

The nature of Staphylococcal products is defined as small (less than 3kDa), heat-stable, non-proteinaceous, and not cell wall-related. So far, the immunomodulatory molecules identified in *S. aureus* are mostly peptides or proteins (Bardoel et al., 2012; Chu et al., 2018; Tajima et al., 2006; Tajima et al., 2008; Chekabab et al., 2015), in contrast to our results. In addition to the possibility that the molecules are non-proteinaceous, they might be proteinase K-resistant peptides/proteins as protease-resistant proteins have been reported before, mainly in prion and *Mycoplasma* (Abd-Elhadi et al., 2015; Butler et al., 1991; Salta et al., 2014).

Alternatively, with the assistance of in silico prediction, liquid chromatography-mass spectrometry/mass spectrometry (LC-MS/MS) could be applied to identify the Staphylococcal secretomes that interact with human TLR4 (Boleij et al., 2011a; Fels et al., 2017). While computational technique provides rapid discovery of the potential candidates based on empirical data, in some cases the predicted secretomes might be poorly expressed. With major improvements in accuracy, efficiency and sensitivity, LC-MS/MS enables identification of small organic compounds such as proteins and sugars in the field of host-pathogen interactions (Portillo et al., 2011; Han et al., 2013; Schweppe et al., 2015). However, this technique is vastly restricted by the ratio of macromolecules between host and pathogen (for example, human proteomes have about 20457 proteins while the prokaryotic counterparts have about 4423 proteins), which results in a decrease in the number of identified pathogen proteins when analysing mixed host and pathogen lysates (Fels et al., 2017). Identification of Staphylococcal immunomodulatory molecules would illustrate a novel immune evasion mechanism contributing to *S. aureus* pathogenesis, and implicate novel anti-Staphylococcal therapeutic interventions.

## 6.5 Limitations

This study has several limitations. Firstly, one difficulty throughout the project was inadequate phosphorylation to bacterial stimuli. Phosphatase inhibitors were included in the lysis buffer to inhibit phosphatase activity during cell lysis. However, the inadequate response could not be ascribed to the endogenous immune tolerance since the positive control TNF $\alpha$  failed to cause a significant increase in phosphorylation in several blots. Other pharmacological reagents such as EGF and phorbol 12-myristate-13-acetate could be applied as positive controls for Akt and MAPK activation (Rice et al., 2002).

Alternatively, techniques to assess cumulative responses such as ELISA, intracellular flow cytometry or mass spectrometry, might provide a clearer readout.

Secondly, this project was limited to human airway cell lines. Although pulmonary epithelial cell lines and culture models are commonly used, there are inconsistencies between different phenotypes of cell lines (Kaza et al., 2010), and between cell lines and primary cells (Strobel et al., 2016) in the host response. Since polymicrobial respiratory infections are highly prevalent in CF individuals, CF-isolated primary tissues and immortalised cell lines would be critical for the understanding of biochemical, physiological and genetic mechanisms of host defence underlying CF defect (Gruenert et al., 2004).

Thirdly, this study requires confirmation using *in vivo* models. Using animal models would provide a panorama of the host response to *S. aureus*/Bcc co-infection. Moreover, there is limited research using rodent models to study polymicrobial infections. In this project, due to cell damage caused by *S. aureus*, pro-inflammatory responses to *S. aureus* and Bcc live bacteria were not evaluated. Alternatively, murine models could be applied to assess host responses to either acute or chronic polymicrobial infections, by intratracheal injection or agar bead-based method, respectively (Kukavica-Ibrulj et al., 2014). These models would be useful to study microbial clearance, inflammatory responses, mortality and lung histopathology.

Lastly, the interaction between *S. aureus* secreted products and TLR4 complex requires binding assays to be validated. Since antagonising TLR4, reduced *B. cenocepacia* J2315-induced pro-inflammatory response, it is plausible that Staphylococcal products competitively bind to TLR4 complex. However, this requires separating *S. aureus* supernatant using chromatography and binding assays to confirm TLR4 is the host target (Nicholas et al., 2017; Fuchs et al., 2018).

## 6.6 Final conclusions

This thesis demonstrates the disruptive effect of two microorganism species, *S. aureus* and Bcc on human airway epithelial monolayers, as well as their modulation on host signalling cascades. *S. aureus* and Bcc are also able to adhere to and enter into airway epithelial cells. *B. cenocepacia* J2315 exposure leads to phosphorylation of MAPK and NF- $\kappa$ B cascades, which results in substantial IL-8 production. However, this robust pro-inflammatory chemokine response is subverted by secreted molecules from *S. aureus*, partially through blocking TLR4 pathway. Preliminary characterisation describes the anti-inflammatory molecules as small, heat-stable, non-proteinaceous, and not cell wall-related factors. The findings might provide valuable insight into a novel Staphylococcal

immune evasion mechanism, as well as a resolution of tissue injury resulting from excessive or chronic inflammation.

## Reference

- Abd-Elhadi, S., Honig, A., Simhi-Haham, D., Schechter, M., Linetsky, E., Ben-Hur, T., Sharon, R., 2015. Total and Proteinase K-resistant  $\alpha$ -synuclein levels in erythrocytes, determined by their ability to bind phospholipids, associate with Parkinson's disease. *Sci Rep*, 5, 11120.
- Abou Alaiwa, M.H., Reznikov, L.R., Gansemer, N.D., Sheets, K.A., Horswill, A.R., Stoltz, D.A., Zabner, J., Welsh, M.J., 2014. pH modulates the activity and synergism of the airway surface liquid antimicrobials  $\beta$ -defensin-3 and LL-37. *Proc Natl Acad Sci*, 111, 18703–18708.
- Ahlgren, H.G., Benedetti, A., Landry, J.S., Bernier, J., Matouk, E., Radzioch, D., Lands, L.C., Rousseau, S., Nguyen, D., 2015. Clinical outcomes associated with *Staphylococcus aureus* and *Pseudomonas aeruginosa* airway infections in adult cystic fibrosis patients. *BMC Pulm Med*, 15, 918.
- Ahn, D.S., Parker, D., Planet, P.J., Nieto, P.A., Bueno, S.M., Prince, A., 2014. Secretion of IL-16 through TNFR1 and calpain-caspase signaling contributes to MRSA pneumonia. *Mucosal Immunol*, 7, 1366–1374.
- Akira, S., Uematsu, S., Takeuchi, O., 2006. Pathogen Recognition and Innate Immunity. *Cell*, 124, 783–801.
- Aksoy, E., Taboubi, S., Torres, D., Delbauve, S., Hachani, A., Whitehead, M.A., Pearce, W.P., Berenjano-Martin, I., Nock, G., Filloux, A., Beyaert, R., Flamand, V., Vanhaesebroeck, B., 2012. The p110 $\delta$  isoform of the kinase PI(3)K controls the subcellular compartmentalization of TLR4 signaling and protects from endotoxic shock. *Nat Immunol*, 13, 1045–1054.
- Albrecht, V.S., Limbago, B.M., Moran, G.J., Krishnadasan, A., Gorwitz, R.J., McDougal, L.K., Talan, D.A., EMERGENCY ID NET Study Group, 2015. *Staphylococcus aureus* colonization and strain type at various body sites among patients with a closed abscess and uninfected controls at U.S. Emergency Departments. *J. Clin. Microbiol*, 53, 3478–3484.
- Alexopoulou, L., Holt, A.C., Medzhitov, R., Flavell, R.A., 2001. Recognition of double-stranded RNA and activation of NF- $\kappa$ B by Toll-like receptor 3. *Nature*, 413, 732–738.
- Alonzo, F., III, Torres, V.J., 2014. The bicomponent pore-forming leucocidins of *Staphylococcus aureus*. *Microbiol Mol Biol Rev*, 78, 199–230.
- Alonzo, F.3., Kozhaya, L., Rawlings, S.A., Reyes-Robles, T., DuMont, A.L., Myszka, D.G., Landau, N.R., Unutmaz, D., Torres, V.J., 2013. CCR5 is a receptor for *Staphylococcus aureus* leukotoxin ED. *Nature*, 493, 51–55.
- Androulidaki, A., Iliopoulos, D., Arranz, A., Doxaki, C., Schworer, S., Zacharioudaki, V., Margioris, A.N., Tsiachlis, P.N., Tsatsanis, C., 2009. The kinase Akt1 controls macrophage response to lipopolysaccharide by regulating microRNAs. *Immunity*, 31, 220–231.
- Arcaro, A., Wymann, M.P., 1993. Wortmannin is a potent phosphatidylinositol 3-kinase inhibitor: the role of phosphatidylinositol 3,4,5-trisphosphate in neutrophil responses. *Biochem J*, 296 (Pt 2), 297–301.
- Archambaud, C., Salcedo, S.P., Lelouard, H., Devilard, E., de Bovis, B., Van Rooijen, N., Gorvel, J.-P., Malissen, B., 2010. Contrasting roles of macrophages and dendritic cells in controlling initial pulmonary *Brucella* infection. *Eur J Immunol*, 40, 3458–3471.
- Archer, N.K., Mazaitis, M.J., Costerton, J.W., Leid, J.G., Powers, M.E., Shirtliff, M.E., 2011. *Staphylococcus aureus* biofilms. *Virulence*, 2, 445–459.



- Ariki, S., Nishitani, C., Kuroki, Y., 2012. Diverse functions of pulmonary collectins in host defense of the lung. *J Biomed Biotechnol*, 2012, 532071–7.
- Armbruster, N.S., Richardson, J.R., Schreiner, J., Klenk, J., Gunter, M., Kretschmer, D., Poschel, S., Schenke-Layland, K., Kalbacher, H., Clark, K., Autenrieth, S.E., 2016. PSM peptides of *Staphylococcus aureus* activate the p38-CREB pathway in dendritic cells, thereby modulating cytokine production and T cell priming. *J Immunol*, 196, 1284–1292.
- Arranz, A., Doxaki, C., Vergadi, E., Martinez de la Torre, Y., Vaporidi, K., Lagoudaki, E.D., Ieronymaki, E., Androulidaki, A., Venihaki, M., Margioris, A.N., Stathopoulos, E.N., Tsiachlis, P.N., Tsatsanis, C., 2012. Akt1 and Akt2 protein kinases differentially contribute to macrophage polarization. *Proc Natl Acad Sci*, 109, 9517–9522.
- Arya, R., Ravikumar, R., Santhosh, R.S., Princy, S.A., 2015. SarA based novel therapeutic candidate against *Staphylococcus aureus* associated with vascular graft infections. *Front Microbio*, 6, 416.
- Askarian, F., Wagner, T., Johannessen, M., Nizet, V., 2018. *Staphylococcus aureus* modulation of innate immune responses through Toll-like (TLR), (NOD)-like (NLR) and C-type lectin (CLR) receptors. *FEMS Microbiol Rev*, 42, 656–671.
- Aubert, D.F., Xu, H., Yang, J., Shi, X., Gao, W., Li, L., Bisaro, F., Chen, S., Valvano, M.A., Shao, F., 2016. A *Burkholderia* Type VI effector deamidates Rho GTPases to activate the pyrin inflammasome and trigger inflammation. *Cell Host Microbe*, 19, 664–674.
- Baba, T., Bae, T., Schneewind, O., Takeuchi, F., Hiramatsu, K., 2008. Genome sequence of *Staphylococcus aureus* strain Newman and comparative analysis of staphylococcal genomes: polymorphism and evolution of two major pathogenicity islands. *J Bacterio*, 190, 300–310.
- Bai, D., Ueno, L., Vogt, P.K., 2009. Akt-mediated regulation of NF- $\kappa$ B and the essentialness of NF- $\kappa$ B for the oncogenicity of PI3K and Akt. *Int J Oncol*, 125, 2863–2870.
- Bailey, K.L., Wyatt, T.A., Romberger, D.J., Sisson, J.H., 2009. Alcohol functionally upregulates Toll-like receptor 2 in airway epithelial cells. *Alcohol Clin Exp Res*, 33, 499–504.
- Bæk, K.T., Frees, D., Renzoni, A., Barras, C., Rodriguez, N., Manzano, C., Kelley, W.L., 2013. Genetic variation in the *Staphylococcus aureus* 8325 strain lineage revealed by whole-genome sequencing. *PLoS ONE*, 8, e77122.
- Baldan, R., Cigana, C., Testa, F., Bianconi, I., De Simone, M., Pellin, D., Di Serio, C., Bragonzi, A., Cirillo, D.M., 2014. Adaptation of *Pseudomonas aeruginosa* in cystic fibrosis airways influences virulence of *Staphylococcus aureus* *in vitro* and murine models of co-infection. *PLoS ONE*, 9, e89614.
- Ballard, S.T., Inglis, S.K., 2004. Liquid secretion properties of airway submucosal glands. *J Physiol-LONDON*, 556, 1–10.
- Bamford, S., Ryley, H., Jackson, S.K., 2007. Highly purified lipopolysaccharides from *Burkholderia cepacia* complex clinical isolates induce inflammatory cytokine responses via TLR4-mediated MAPK signalling pathways and activation of NF- $\kappa$ B. *Cell Microbiol*, 9, 532–543.
- Bardoel, B.W., Vos, R., Bouman, T., Aerts, P.C., Bestebroer, J., Huizinga, E.G., Brondijk, T.H.C., van Strijp, J.A.G., de Haas, C.J.C., 2012. Evasion of Toll-like receptor 2 activation by staphylococcal superantigen-like protein 3. *J Mol Med*, 90, 1109–1120.
- Barkauskas, C.E., Counce, M.J., Rackley, C.R., Bowie, E.J., Keene, D.R., Stripp, B.R., Randell, S.H., Noble, P.W., Hogan, B.L.M., 2013. Type 2 alveolar cells are stem cells in adult lung. *J Clin Invest*, 123, 3025–3036.

- Bechelli, J., Smalley, C., Zhao, X., Judy, B., Valdes, P., Walker, D.H., Fang, R., 2016. MyD88 mediates instructive signaling in dendritic cells and protective inflammatory response during rickettsial infection. *Infect Immun*, 84, 883–893.
- Becker, M.N., Diamond, G., Verghese, M.W., Randell, S.H., 2000. CD14-dependent lipopolysaccharide-induced  $\beta$ -defensin-2 expression in human tracheobronchial epithelium. *J Biol Chem*, 275, 29731–29736.
- Bedoret, D., Wallemacq, H., Marichal, T., Desmet, C., Quesada Calvo, F., Henry, E., Closset, R., Dewals, B., Thielen, C., Gustin, P., de Leval, L., Van Rooijen, N., Le Moine, A., Vanderplasschen, A., Cataldo, D., Drion, P.-V., Moser, M., Lekeux, P., Bureau, F., 2009. Lung interstitial macrophages alter dendritic cell functions to prevent airway allergy in mice. *J Clin Invest*, 119, 3723–3738.
- Below, S., Konkel, A., Zeeck, C., Muller, C., Kohler, C., Engelmann, S., Hildebrandt, J.P., 2009. Virulence factors of *Staphylococcus aureus* induce Erk-MAP kinase activation and c-Fos expression in S9 and 16HBE14o- human airway epithelial cells. *Am J Physiol Lung Cell Mol Physiol*, 296, L470–L479.
- Bernier, S.P., Workentine, M.L., Li, X., Magarvey, N.A., O'Toole, G.A., Surette, M.G., 2016. Cyanide toxicity to *Burkholderia cenocepacia* is modulated by polymicrobial communities and environmental factors. *Front Microbio*, 7, 725.
- Berube, B.J., Sampedro, G.R., Otto, M., Bubeck Wardenburg, J., 2014. The PSM $\alpha$  locus regulates production of *Staphylococcus aureus*  $\alpha$ -toxin during infection. *Infect Immun*, 82, 3350–3358.
- Bestebroer, J., Poppelier, M.J.J.G., Ulfman, L.H., Lenting, P.J., Denis, C.V., van Kessel, K.P.M., van Strijp, J.A.G., de Haas, C.J.C., 2007. Staphylococcal superantigen-like 5 binds PSGL-1 and inhibits P-selectin-mediated neutrophil rolling. *Blood*, 109, 2936–2943.
- Bhattacharyya, S., Gutti, U., Mercado, J., Moore, C., Pollard, H.B., Biswas, R., 2011. MAPK signaling pathways regulate IL-8 mRNA stability and IL-8 protein expression in cystic fibrosis lung epithelial cell lines. *Am J Physiol Lung Cell Mol Physiol*, 300, L81–7.
- Bibeau-Poirier, A., Servant, M.J., 2008. Roles of ubiquitination in pattern-recognition receptors and type I interferon receptor signaling. *Cytokine*, 43, 359–367.
- Bingisser, R.M., Tilbrook, P.A., Holt, P.G., Kees, U.R., 1998. Macrophage-derived nitric oxide regulates T cell activation via reversible disruption of the Jak3/STAT5 signaling pathway. *J Immunol*, 160, 5729–5734.
- Bjarnsholt, T., Jensen, P.Ø., Fiandaca, M.J., Pedersen, J., Hansen, C.R., Andersen, C.B., Pressler, T., Givskov, M., Høiby, N., 2009. *Pseudomonas aeruginosa* biofilms in the respiratory tract of cystic fibrosis patients. *Pediatr Pulmonol*, 44, 547–558.
- Blackburn, L., Brownlee, K., Conway, S., Denton, M., 2004. “Cepacia syndrome” with *Burkholderia multivorans*, 9 years after initial colonization. *J Cyst Fibros*, 3, 133–134.
- Blohmke, C.J., Victor, R.E., Hirschfeld, A.F., Elias, I.M., Hancock, D.G., Lane, C.R., Davidson, A.G.F., Wilcox, P.G., Smith, K.D., Overhage, J., Hancock, R.E.W., Turvey, S.E., 2008. Innate immunity mediated by TLR5 as a novel anti-inflammatory target for cystic fibrosis lung disease. *J Immunol*, 180, 7764–7773.
- Boleij, A., Laarakkers, C.M., Gloerich, J., Swinkels, D.W., Tjalsma, H., 2011a. Surface-affinity profiling to identify host-pathogen interactions. *Infect Immun*, 79, 4777–4783.
- Boleij, A., Muytjens, C.M.J., Bukhari, S.I., Cayet, N., Glaser, P., Hermans, P.W.M., Swinkels, D.W., Bolhuis, A., Tjalsma, H., 2011b. Novel clues on the specific association of *Streptococcus gallolyticus* subsp. *gallolyticus* with colorectal cancer. *J Infect Dis*, 203, 1101–1109.

- Botos, I., Segal, D.M., Davies, D.R., 2011. The structural biology of Toll-like receptors. *Structure*, 19, 447–459.
- Boussaud, V., Guillemain, R., Grenet, D., Coley, N., Souillamas, R., Bonnette, P., Stern, M., 2008. Clinical outcome following lung transplantation in patients with cystic fibrosis colonised with *Burkholderia cepacia* complex: results from two French centres. *Thorax*, 63, 732–737.
- Bowler, L.L., Zhanel, G.G., Ball, T.B., Saward, L.L., 2012. Mature *Pseudomonas aeruginosa* biofilms prevail compared to young biofilms in the presence of ceftazidime. *Antimicrob Agents Chemother*, 56, 4976–4979.
- Boyer, H.W., Roulland-Dussoix, D., 1969. A complementation analysis of the restriction and modification of DNA in *Escherichia coli*. *J Mol Biol*, 41, 459–472.
- Brady, R.A., Mocca, C.P., Plaut, R.D., Takeda, K., Burns, D.L., 2018. Comparison of the immune response during acute and chronic *Staphylococcus aureus* infection. *PLoS ONE*, 13, e0195342.
- Bragonzi, A., Farulla, I., Paroni, M., Twomey, K.B., Pirone, L., Lorè, N.I., Bianconi, I., Dalmastri, C., Ryan, R.P., Bevivino, A., 2012. Modelling co-infection of the cystic fibrosis lung by *Pseudomonas aeruginosa* and *Burkholderia cenocepacia* reveals influences on biofilm formation and host response. *PLoS ONE*, 7, e52330.
- Brauner, A., Fridman, O., Gefen, O., Balaban, N.Q., 2016. Distinguishing between resistance, tolerance and persistence to antibiotic treatment. *Nat Rev Microbiol*, 14, 320–330.
- Brint, E.K., Xu, D., Liu, H., Dunne, A., McKenzie, A.N.J., O'Neill, L.A.J., Liew, F.Y., 2004. ST2 is an inhibitor of interleukin 1 receptor and Toll-like receptor 4 signaling and maintains endotoxin tolerance. *Nat Immunol*, 5, 373–379.
- Brune, K., Frank, J., Schwingshackl, A., Finigan, J., Sidhaye, V.K., 2015. Pulmonary epithelial barrier function: some new players and mechanisms. *Am J Physiol Lung Cell Mol Physiol*, 308, L731–L745.
- Brunet, A., Bonni, A., Zigmond, M.J., Lin, M.Z., Juo, P., Hu, L.S., Anderson, M.J., Arden, K.C., Blenis, J., Greenberg, M.E., 1999. Akt promotes cell survival by phosphorylating and inhibiting a Forkhead transcription factor. *Cell*, 96, 857–868.
- Bucior, I., Mostov, K., Engel, J.N., 2010. *Pseudomonas aeruginosa*-mediated damage requires distinct receptors at the apical and basolateral surfaces of the polarized epithelium. *Infect Immun*, 78, 939–953.
- Bucior, I., Pielage, J.F., Engel, J.N., 2012. *Pseudomonas aeruginosa* pili and flagella mediate distinct binding and signaling events at the apical and basolateral surface of airway epithelium. *PLoS Pathog*, 8, e1002616–18.
- Buddenkotte, J., Stroh, C., Engels, I.H., Moormann, C., Shpacovitch, V.M., Seeliger, S., Vergnolle, N., Vestweber, D., Luger, T.A., Schulze-Osthoff, K., Steinhoff, M., 2005. Agonists of proteinase-activated receptor-2 stimulate upregulation of intercellular cell adhesion molecule-1 in primary human keratinocytes via activation of NF- $\kappa$ B. *J Investig Dermatol*, 124, 38–45.
- Bustamante-Marin, X.M., Ostrowski, L.E., 2017. Cilia and Mucociliary Clearance. *Cold Spring Harb Perspect Bio*, 9, a028241.
- Butler, G.H., Kotani, H., Kong, L., Frick, M., Evancho, S., Stanbridge, E.J., McGarrity, G.J., 1991. Identification and characterization of proteinase K-resistant proteins in members of the class Mollicutes. *Infect Immun*, 59, 1037–1042.

- Bylund, J., Burgess, L.-A., Cescutti, P., Ernst, R.K., Speert, D.P., 2006. Exopolysaccharides from *Burkholderia cenocepacia* inhibit neutrophil chemotaxis and scavenge reactive oxygen species. *J Biol Chem*, 281, 2526–2532.
- Cachat, J., Deffert, C., Hugues, S., Krause, K.-H., 2015. Phagocyte NADPH oxidase and specific immunity. *Clin Sci (Lond)*, 128, 635–648.
- Cai, Y., Sugimoto, C., Arainga, M., Alvarez, X., Didier, E.S., Kuroda, M.J., 2014. *In vivo* characterization of alveolar and interstitial lung macrophages in rhesus macaques: implications for understanding lung disease in humans. *J Immunol*, 192, 2821–2829.
- Calkovska, A., Uhliarova, B., Joskova, M., Franova, S., Kolomaznik, M., Calkovsky, V., Smolarova, S., 2015. Pulmonary surfactant in the airway physiology: A direct relaxing effect on the smooth muscle. *Respir Physiol Neurobiol*, 209, 95–105.
- Caraher, E., Duff, C., Mullen, T., Mc Keon, S., Murphy, P., Callaghan, M., McClean, S., 2007. Invasion and biofilm formation of *Burkholderia dolosa* is comparable with *Burkholderia cenocepacia* and *Burkholderia multivorans*. *J Cyst Fibros*, 6, 49–56.
- Carey, R.M., Workman, A.D., Chen, B., Adappa, N.D., Palmer, J.N., Kennedy, D.W., Lee, R.J., Cohen, N.A., 2015. *Staphylococcus aureus* triggers nitric oxide production in human upper airway epithelium. *Int Forum Allergy Rhinol*, 5, 808–813.
- Carlsson, E., Ding, J.L., Byrne, B., 2016. SARM modulates MyD88-mediated TLR activation through BB-loop dependent TIR-TIR interactions. *Biochim Biophys Acta*, 1863, 244–253.
- Cervantes, J.L., Dunham-Ems, S.M., La Vake, C.J., Petzke, M.M., Sahay, B., Sellati, T.J., Radolf, J.D., Salazar, J.C., 2011. Phagosomal signaling by *Borrelia burgdorferi* in human monocytes involves Toll-like receptor (TLR) 2 and TLR8 cooperativity and TLR8-mediated induction of IFN- $\beta$ . *Proc Natl Acad Sci*, 108, 3683–3688.
- Cervantes, J.L., La Vake, C.J., Weinerman, B., Luu, S., O'Connell, C., Verardi, P.H., Salazar, J.C., 2013. Human TLR8 is activated upon recognition of *Borrelia burgdorferi* RNA in the phagosome of human monocytes. *J Leukoc Biol*, 94, 1231–1241.
- Chan, P.F., Foster, S.J., Ingham, E., Clements, M.O., 1998. The *Staphylococcus aureus* alternative sigma factor  $\sigma_B$  controls the environmental stress response but not starvation survival or pathogenicity in a mouse abscess model. *J Bacterio*, 180, 6082–6089.
- Chavakis, T., Preissner, K.T., Herrmann, M., 2007. The anti-inflammatory activities of *Staphylococcus aureus*. *Trends Immunol*, 28, 408–418.
- Chekabab, S.M., Silverman, R.J., Lafayette, S.L., Luo, Y., Rousseau, S., Nguyen, D., 2015. *Staphylococcus aureus* inhibits IL-8 responses induced by *Pseudomonas aeruginosa* in airway epithelial cells. *PLoS ONE*, 10, e0137753–19.
- Chen, Z.J., 2005. Ubiquitin signalling in the NF- $\kappa$ B pathway. *Nat Cell Biol*, 7, 758–765.
- Cheng, A.G., McAdow, M., Kim, H.K., Bae, T., Missiakas, D.M., Schneewind, O., 2010. Contribution of coagulases towards *Staphylococcus aureus* disease and protective immunity. *PLoS Pathog*, 6, e1001036.
- Chi, C.-Y., Lin, C.-C., Liao, I.-C., Yao, Y.-C., Shen, F.-C., Liu, C.-C., Lin, C.-F., 2014. Panton-Valentine leukocidin facilitates the escape of *Staphylococcus aureus* from human keratinocyte endosomes and induces apoptosis. *J Infect Dis*, 209, 224–235.
- Choi, E.Y., Santoso, S., Chavakis, T., 2009. Mechanisms of neutrophil transendothelial migration. *Front Biosci*, 14, 1596–1605.

- Chow, J.C., Young, D.W., Golenbock, D.T., Christ, W.J., Gusovsky, F., 1999. Toll-like receptor-4 mediates lipopolysaccharide-induced signal transduction. *J Biol Chem*, 274, 10689–10692.
- Chu, M., Zhou, M., Jiang, C., Chen, X., Guo, L., Zhang, M., Chu, Z., Wang, Y., 2018. *Staphylococcus aureus* Phenol-Soluble Modulins  $\alpha 1$ – $\alpha 3$  Act as novel Toll-like receptor (TLR) 4 antagonists to inhibit HMGB1/TLR4/NF- $\kappa$ B signaling pathway. *Front Immunol*, 9, 596–13.
- Chuang, T.-H., Ulevitch, R.J., 2004. Triad3A, an E3 ubiquitin-protein ligase regulating Toll-like receptors. *Nat Immunol*, 5, 495–502.
- Chun, J., Prince, A., 2009. TLR2-induced calpain cleavage of epithelial junctional proteins facilitates leukocyte transmigration. *Cell Host Microbe*, 5, 47–58.
- Chung, M.C., Wines, B.D., Baker, H., Langley, R.J., Baker, E.N., Fraser, J.D., 2007. The crystal structure of Staphylococcal superantigen-like protein 11 in complex with sialyl Lewis X reveals the mechanism for cell binding and immune inhibition. *Mol Microbiol*, 66, 1342–1355.
- Clarke, D.L., Davis, N.H.E., Majithiya, J.B., Piper, S.C., Lewis, A., Sleeman, M.A., Corkill, D.J., May, R.D., 2014. Development of a mouse model mimicking key aspects of a viral asthma exacerbation. *Clin Sci (Lond)*, 126, 567–580.
- Clauditz, A., Resch, A., Wieland, K.-P., Peschel, A., Götz, F., 2006. Staphyloxanthin plays a role in the fitness of *Staphylococcus aureus* and its ability to cope with oxidative stress. *Infect Immun*, 74, 4950–4953.
- Coburn, B., Wang, P.W., Caballero, J.D., Clark, S. T., Brahma, V., Donaldson, S., Zhang, Y., Surendra, A., Gong, Y., Elizabeth Tullis, D., Yau, Y.C., Waters, V.J., Hwang, D.M., Guttman, D.S., 2015. Lung microbiota across age and disease stage in cystic fibrosis. *Sci Rep*, 5, 10241.
- Coffey, B.M., Anderson, G.G., 2014. Biofilm Formation in the 96-Well Microtiter Plate. *Methods Mol Biol*, 1149, 631–641.
- Cohen, P., 2014. The TLR and IL-1 signalling network at a glance. *J Cell Sci*, 127, 2383–2390.
- Cohen T.S., Boland M.L., Boland B.B., Takahashi V., Tovchigrechko A., Lee Y., Wilde A.D., Mazaitis M.J., Jones-Nelson O., Tkaczyk C., Raja R., Stover C.K., Sellman B.R., 2018. *S. aureus* evades macrophage killing through NLRP3-dependent effects on mitochondrial trafficking. *Cell Rep*, 22(9), 2431–2441.
- Collins, J., Rudkin, J., Recker, M., Pozzi, C., Gara, J.P.O.A., Massey, R.C., 2010. Offsetting virulence and antibiotic resistance costs by MRSA. *ISME J*, 4, 577–584.
- Colonne, P.M., Winchell, C.G., Voth, D.E., 2016. Hijacking host cell highways: manipulation of the host actin cytoskeleton by obligate intracellular bacterial pathogens. *Front Cell Infect Microbiol*, 6, 107.
- Conway, B.-A.D., Chu, K.K., Bylund, J., Altman, E., Speert, D.P., 2004. Production of exopolysaccharide by *Burkholderia cenocepacia* results in altered cell - surface interactions and altered bacterial clearance in mice. *J Infect Dis*, 190, 957–966.
- Cosseau, C., Devine, D.A., Dullaghan, E., Gardy, J.L., Chikatamarla, A., Gellatly, S., Yu, L.L., Pistolic, J., Falsafi, R., Tagg, J., Hancock, R.E.W., 2008. The commensal *Streptococcus salivarius* K12 downregulates the innate immune responses of human epithelial cells and promotes host-microbe homeostasis. *Infect Immun*, 76, 4163–4175.

- Cozens, A.L., Yezzi, M.J., Kunzelmann, K., Ohrui, T., Chin, L., Eng, K., Finkbeiner, W.E., Widdicombe, J.H., Gruenert, D.C., 1994. CFTR expression and chloride secretion in polarized immortal human bronchial epithelial cells. *Am J Respir Cell Mol Biol*, 10, 38–47.
- Crapo, J.D., Barry, B.E., Gehr, P., Bachofen, M., Weibel, E.R., 1982. Cell number and cell characteristics of the normal human lung. *Am Rev Respir Dis*, 126, 332–337.
- Cremer, T.J., Shah, P., Cormet-Boyaka, E., Valvano, M.A., Butchar, J.P., Tridandapani, S., 2011. Akt-mediated proinflammatory response of mononuclear phagocytes infected with *Burkholderia cenocepacia* occurs by a novel GSK-3-dependent, I $\kappa$ B Kinase-independent mechanism. *J Immunol*, 187, 635–643.
- Crystal, R.G., Randell, S.H., Engelhardt, J.F., Voynow, J., Sunday, M.E., 2008. Airway epithelial cells: current concepts and challenges. *Proc Am Thorac Soc*, 5, 772–777.
- Cunha, M.V., Sousa, S.A., Leitão, J.H., Moreira, L.M., Videira, P.A., Sá-Correia, I., 2004. Studies on the involvement of the exopolysaccharide produced by cystic fibrosis-associated isolates of the *Burkholderia cepacia* complex in biofilm formation and in persistence of respiratory infections. *J Clin Microbiol*, 42, 3052–3058.
- Cystic Fibrosis Foundation, 2017. Cystic Fibrosis Foundation Patient Registry 2016 Annual Data Report. Bethesda, Maryland.
- D'Souza, W.N., Chang, C.F., Fischer, A.M., Li, M., Hedrick, S.M., 2008. The Erk2 MAPK regulates CD8 T cell proliferation and survival. *J Immunol*, 181, 7617–7629.
- Dai, W., Wang, H., Zhou, Q., Feng, X., Lu, Z., Li, D., Yang, Z., Liu, Y., Li, Y., Xie, G., Shen, K., Yang, Y., Zheng, Y., Li, S., 2018. The concordance between upper and lower respiratory microbiota in children with *Mycoplasma pneumoniae* pneumonia. *Emerg Microbes Infect*, 7, 207.
- Dalton, T., Dowd, S.E., Wolcott, R.D., Sun, Y., Watters, C., Griswold, J.A., Rumbaugh, K.P., 2011. An *in vivo* polymicrobial biofilm wound infection model to study interspecies interactions. *PLoS ONE*, 6, e27317–10.
- Dasu, M.R., Devaraj, S., Park, S., Jialal, I., 2010. Increased Toll-like receptor (TLR) activation and TLR ligands in recently diagnosed type 2 diabetic subjects. *Diabetes Care*, 33, 861–868.
- Davidson, B.A., Vethanayagam, R.R., Grimm, M.J., Mullan, B.A., Raghavendran, K., Blackwell, T.S., Freeman, M.L., Ayyasamy, V., Singh, K.K., Sporn, M.B., Itagaki, K., Hauser, C.J., Knight, P.R., Segal, B.H., 2013. NADPH oxidase and Nrf2 regulate gastric aspiration-induced inflammation and acute lung injury. *J Immunol*, 190, 1714–1724.
- Davies, S.P., Reddy, H., Caivano, M., Cohen, P., 2000. Specificity and mechanism of action of some commonly used protein kinase inhibitors. *Biochem J*, 351, 95–105.
- de C Ventura, G.M., Le Goffic, R., Balloy, V., Plotkowski, M.-C., Chignard, M., Si-Tahar, M., 2008. TLR 5, but neither TLR2 nor TLR4, is involved in lung epithelial cell response to *Burkholderia cenocepacia*. *FEMS Immunol Med Microbiol*, 54, 37–44.
- de Haas, C.J.C., Veldkamp, K.E., Peschel, A., Weerkamp, F., van Wamel, W.J.B., Heezius, E.C.J.M., Poppelier, M.J.J.G., van Kessel, K.P.M., van Strijp, J.A.G., 2004. Chemotaxis inhibitory protein of *Staphylococcus aureus*, a bacterial anti-inflammatory agent. *J Exp Med*, 199, 687–695.
- de Jong, N.W.M., Ramyar, K.X., Guerra, F.E., Nijland, R., Fevre, C., Voyich, J.M., McCarthy, A.J., Garcia, B.L., van Kessel, K.P.M., van Strijp, J.A.G., Geisbrecht, B.V., Haas, P.-J.A., 2017. Immune evasion by a staphylococcal inhibitor of myeloperoxidase. *Proc Natl Acad Sci*, 114, 9439–9444.

- de Oliveira, A., Cataneli Pereira, V., Pinheiro, L., Moraes Riboli, D.F., Benini Martins, K., Ribeiro de Souza da Cunha, M. de L., 2016. Antimicrobial resistance profile of planktonic and biofilm cells of *Staphylococcus aureus* and coagulase-negative Staphylococci. *Int J Mol Sci*, 17, 1423.
- De Soya, A., Ellis, C.D., Khan, C.M.A., Corris, P.A., Demarco de Hormaeche, R., 2004. *Burkholderia cenocepacia* lipopolysaccharide, lipid A, and proinflammatory activity. *Am J Respir Crit Care Med*, 170, 70–77.
- De Soya, A., Meachery, G., Hester, K.L.M., Nicholson, A., Parry, G., Tocewicz, K., Pillay, T., Clark, S., Lordan, J.L., Schueler, S., Fisher, A.J., Dark, J.H., Gould, F.K., Corris, P.A., 2010. Lung transplantation for patients with cystic fibrosis and *Burkholderia cepacia* complex infection: a single-center experience. *J Heart Lung Transplant*, 29, 1395–1404.
- De Soya, A., Silipo, A., Lanzetta, R., Govan, J.R., Molinaro, A., 2008. Chemical and biological features of *Burkholderia cepacia* complex lipopolysaccharides. *Innate Immun*, 14, 127–144.
- Deisenhofer, J., 1981. Crystallographic refinement and atomic models of a human Fc fragment and its complex with fragment B of protein A from *Staphylococcus aureus* at 2.9- and 2.8-Å resolution. *Biochemistry*, 20, 2361–2370.
- DeLeon, S., Clinton, A., Fowler, H., Everett, J., Horswill, A.R., Rumbaugh, K.P., 2014. Synergistic interactions of *Pseudomonas aeruginosa* and *Staphylococcus aureus* in an *in vitro* wound model. *Infect Immun*, 82, 4718–4728.
- Dennehy, R., Romano, M., Ruggiero, A., Mohamed, Y.F., Dignam, S.L., Mujica Troncoso, C., Callaghan, M., Valvano, M.A., Berisio, R., McClean, S., 2017. The *Burkholderia cenocepacia* peptidoglycan-associated lipoprotein is involved in epithelial cell attachment and elicitation of inflammation. *Cell Microbiol*, 19, e12691.
- Devaraj, S., Dasu, M.R., Rockwood, J., Winter, W., Griffen, S.C., Jialal, I., 2008. Increased toll-like receptor (TLR) 2 and TLR4 expression in monocytes from patients with type 1 diabetes: further evidence of a proinflammatory state. *J Clin Endocrinol Metab*, 93, 578–583.
- Dhanasekaran, D.N., Kashef, K., Lee, C.M., Xu, H., Reddy, E.P., 2007. Scaffold proteins of MAP-kinase modules. *Oncogene*, 26, 3185–3202.
- Di Domenico, E.G., Cavallo, I., Bordignon, V., Prignano, G., Sperduti, I., Gurtner, A., Trento, E., Toma, L., Pimpinelli, F., Capitanio, B., Ensoli, F., 2018. Inflammatory cytokines and biofilm production sustain *Staphylococcus aureus* outgrowth and persistence: a pivotal interplay in the pathogenesis of Atopic Dermatitis. *Sci Rep*, 8, 1483.
- Diebold, S.S., Kaisho, T., Hemmi, H., Akira, S., Reis e Sousa, C., 2004. Innate antiviral responses by means of TLR7-mediated recognition of single-stranded RNA. *Science*, 303, 1529–1531.
- Divertie, M.B., Cassan, S.M., Brown, A.L., 1975. Application of ultrastructural morphometry to lung biopsy specimens in pulmonary histiocytosis X. *Thorax*, 30, 326–332.
- Donlan, R.M., Costerton, J.W., 2002. Biofilms: survival mechanisms of clinically relevant microorganisms. *Clin Microbiol Rev*, 15, 167–193.
- Dossett, J.H., Kronvall, G., Williams, R.C., Quie, P.G., 1969. Antiphagocytic effects of Staphylococcal protein A. *J Immunol*, 103, 1405.
- Downey, D.G., Martin, S.L., Dempster, M., Moore, J.E., Keogan, M.T., Starcher, B., Edgar, J., Bilton, D., Elborn, J.S., 2007. The relationship of clinical and inflammatory markers to outcome in stable patients with cystic fibrosis. *Pediatr Pulmonol*, 42, 216–220.

- Dudek, M., Puttur, F., Arnold-Schrauf, C., Kühl, A.A., Holzmann, B., Henriques-Normark, B., Berod, L., Sparwasser, T., 2016. Lung epithelium and myeloid cells cooperate to clear acute pneumococcal infection. *Mucosal Immunol*, 9, 1288–1302.
- Duff, A.J.A., 2002. Psychological consequences of segregation resulting from chronic *Burkholderia cepacia* infection in adults with CF. *Thorax*, 57, 756–758.
- Duff, C., Murphy, P.G., Callaghan, M., McClean, S., 2006. Differences in invasion and translocation of *Burkholderia cepacia* complex species in polarised lung epithelial cells *in vitro*. *Microb Pathog*, 41, 183–192.
- DuMont, A.L., Yoong, P., Day, C.J., Alonzo, F.3., McDonald, W.H., Jennings, M.P., Torres, V.J., 2013. *Staphylococcus aureus* LukAB cytotoxin kills human neutrophils by targeting the CD11b subunit of the integrin Mac-1. *Proc Natl Acad Sci*, 110, 10794–10799.
- D’Elia, R.V., Saint, R.J., Newstead, S.L., Clark, G.C., Atkins, H.S., 2017. Mitogen-activated protein kinases (MAPKs) are modulated during *in vitro* and *in vivo* infection with the intracellular bacterium *Burkholderia pseudomallei*. *Eur J Clin Microbiol Infect Dis*, 36, 2147–2154.
- Edinger, A.L., Thompson, C.B., 2004. Death by design: apoptosis, necrosis and autophagy. *Curr Opin Cell Biol*, 16, 663–669.
- Eisele, N.A., Anderson, D.M., 2011. Host defense and the airway epithelium: frontline responses that protect against bacterial invasion and pneumonia. *J Pathog*, 2011, 249802.
- El-Halfawy, O.M., Naguib, M.M., Valvano, M.A., 2017. Novel antibiotic combinations proposed for treatment of *Burkholderia cepacia* complex infections. *Antimicrob Resist Infect Control*, 6, 120.
- Elias, S., Banin, E., 2012. Multi-species biofilms: living with friendly neighbors. *FEMS Microbiol Rev*, 36, 990–1004.
- Elkhatib, W.F., Hair, P.S., Nyalwidhe, J.O., Cunnion, K.M., 2015. New potential role of serum apolipoprotein E mediated by its binding to clumping factor A during *Staphylococcus aureus* invasive infections to humans. *J Med Microbiol*, 64, 335–343.
- Elliott, C.L., Allport, V.C., Loudon, J.A., Wu, G.D., Bennett, P.R., 2001. Nuclear factor- $\kappa$ B is essential for up-regulation of interleukin-8 expression in human amnion and cervical epithelial cells. *Mol Hum Reprod*, 7, 787–790.
- Essex-Lopresti, A.E., Boddey, J.A., Thomas, R., Smith, M.P., Hartley, M.G., Atkins, T., Brown, N.F., Tsang, C.H., Peak, I.R.A., Hill, J., Beacham, I.R., Titball, R.W., 2005. A type IV pilin, PilA, contributes to adherence of *Burkholderia pseudomallei* and virulence *in vivo*. *Infect Immun*, 73, 1260–1264.
- Evans, C.M., Kim, K., Tuvim, M.J., Dickey, B.F., 2009. Mucus hypersecretion in asthma: causes and effects. *Curr Opin Pulm Med*, 15, 4–11.
- Everaert, A., Coenye, T., 2016. Effect of  $\beta$ -Lactamase inhibitors on *in vitro* activity of  $\beta$ -lactam antibiotics against *Burkholderia cepacia* complex species. *Antimicrob Resist Infect Control*, 5, 44.
- Fahy, J.V., Dickey, B.F., 2010. Airway Mucus Function and Dysfunction. *N Engl J Med*, 363, 2233–2247.
- Falugi, F., Kim, H.K., Missiakas, D.M., Schneewind, O., 2013. Role of protein A in the evasion of host adaptive immune responses by *Staphylococcus aureus*. *MBio*, 4, e00575–13.



- Fan, W., Morinaga, H., Kim, J.J., Bae, E., Spann, N.J., Heinz, S., Glass, C.K., Olefsky, J.M., 2010a. FoxO1 regulates TLR4 inflammatory pathway signalling in macrophages. *EMBO J*, 29, 4223–4236.
- Fang, F.C., 2004. Antimicrobial reactive oxygen and nitrogen species: concepts and controversies. *Nat Rev Micro*, 2, 820–832.
- Fazli, M., Rybtke, M., Steiner, E., Weidel, E., Berthelsen, J., Groizeleau, J., Bin Wu, Zhi, B.Z., Yaming, Z., Kaefer, V., Givskov, M., Hartmann, R.W., Eberl, L., Tolker-Nielsen, T., 2017. Regulation of *Burkholderia cenocepacia* biofilm formation by RpoN and the c-di-GMP effector BerB. *MicrobiologyOpen*, 6, e00480.
- Feil, E.J., Cooper, J.E., Grundmann, H., Robinson, D.A., Enright, M.C., Berendt, T., Peacock, S.J., Smith, J.M., Murphy, M., Spratt, B.G., Moore, C.E., Day, N.P.J., 2003. How Clonal Is *Staphylococcus aureus*? *J Bacterio*, 185, 3307–3316.
- Fels, U., Gevaert, K., Van Damme, P., 2017. Proteogenomics in aid of host–pathogen interaction studies: a bacterial perspective. *Proteomes*, 5, 26–19.
- Feuillet, V., Medjane, S., Mondor, I., Demaria, O., Pagni, P.P., Galan, J.E., Flavell, R.A., Alexopoulou, L., 2006. Involvement of Toll-like receptor 5 in the recognition of flagellated bacteria. *Proc Natl Acad Sci*, 103, 12487–12492.
- Fitzgerald, K.A., Rowe, D.C., Barnes, B.J., Caffrey, D.R., Visintin, A., Latz, E., Monks, B., Pitha, P.M., Golenbock, D.T., 2003. LPS-TLR4 signaling to IRF-3/7 and NF- $\kappa$ B involves the toll adapters TRAM and TRIF. *J Exp Med*, 198, 1043–1055.
- Flannagan, R.S., Jaumouillé, V., Huynh, K.K., Plumb, J.D., Downey, G.P., Valvano, M.A., Grinstein, S., 2012. *Burkholderia cenocepacia* disrupts host cell actin cytoskeleton by inactivating Rac and Cdc42. *Cell Microbiol*, 14, 239–254.
- Flannagan, R.S., Valvano, M.A., 2008. *Burkholderia cenocepacia* requires RpoE for growth under stress conditions and delay of phagolysosomal fusion in macrophages. *Microbiology*, 154, 643–653.
- Flight, W.G., Smith, A., Paisey, C., Marchesi, J.R., Bull, M.J., Norville, P.J., Mutton, K.J., Webb, A.K., Bright-Thomas, R.J., Jones, A.M., Mahenthiralingam, E., 2015. Rapid detection of emerging pathogens and loss of microbial diversity associated with severe lung disease in cystic fibrosis. *J Clin Microbiol*, 53, 2022–2029.
- Fogh, J., Fogh, J.M. and Orfeo, T., 1977. One hundred and twenty-seven cultured human tumor cell lines producing tumors in nude mice. *J Natl Cancer Inst*, 59(1), 221–226.
- Forsgren, A., Sjoquist, J., 1966. “Protein A” from *S. aureus*. *J Immunol*, 97, 822–827.
- Foster, T.J., 2016. The remarkably multifunctional fibronectin binding proteins of *Staphylococcus aureus*. *Eur J Clin Microbiol Infect Dis*, 35, 1923–1931.
- Foster, T.J., 2017. Antibiotic resistance in *Staphylococcus aureus*. Current status and future prospects. *FEMS Microbiol Rev*, 41, 430–449.
- Foster, T.J., Geoghegan, J.A., Ganesh, V.K., Höök, M., 2014. Adhesion, invasion and evasion: the many functions of the surface proteins of *Staphylococcus aureus*. *Nat Rev Microbiol*, 12, 49–62.
- Fournier, B., Philpott, D.J., 2005. Recognition of *Staphylococcus aureus* by the Innate Immune System. *Clin Microbiol Rev*, 18, 521–540.
- Fox, S., Leitch, A.E., Duffin, R., Haslett, C., Rossi, A.G., 2010. Neutrophil apoptosis: relevance to the innate immune response and inflammatory disease. *J Innate Immun*, 2, 216–227.

- Fraser-Pitt, D., Mercer, D., Lovie, E., Robertson, J., O'Neil, D., 2016. Activity of cysteamine against the cystic fibrosis pathogen *Burkholderia cepacia* complex. *Antimicrob Agents Chemother*, 60, 6200–6206.
- Freiberg, J.A., Le Breton, Y., Tran, B.Q., Scott, A.J., Harro, J.M., Ernst, R.K., Goo, Y.A., Mongodin, E.F., Goodlett, D.R., McIver, K.S., Shirtliff, M.E., 2016. Global analysis and comparison of the transcriptomes and proteomes of group a *Streptococcus* biofilms. *mSystems*, 1, e00149–16.
- Fuchs, K., Cardona Gloria, Y., Wolz, O.O., Herster F., Sharma L., Dillen C.A., Täumer C., Dickhöfer S., Bittner Z., Dang T.M., Singh A., Haischer D., Schlöffel M.A., Koymans K.J., Sanmuganantham T., Krach M., Roger T., Le Roy D., Schilling N.A., Frauhammer F., Miller L.S., Nürnberger T., LeibundGut-Landmann S., Gust A.A., Macek B., Frank M., Gouttefangeas C., Dela Cruz C.S., Hartl D., Weber A.N., 2018. The fungal ligand chitin directly binds TLR2 and triggers inflammation dependent on oligomer size. *EMBO rep*, 19, p.e46065.
- Fugère, A., Lalonde Séguin, D., Mitchell, G., Déziel, E., Dekimpe, V., Cantin, A.M., Frost, E., Malouin, F., 2014. Interspecific small molecule interactions between clinical isolates of *Pseudomonas aeruginosa* and *Staphylococcus aureus* from adult cystic fibrosis patients. *PLoS ONE*, 9, e86705.
- Fukao, T., Koyasu, S., 2003. PI3K and negative regulation of TLR signaling. *Trends Immunol*, 24, 358–363.
- Fukao, T., Tanabe, M., Terauchi, Y., Ota, T., Matsuda, S., Asano, T., Kadowaki, T., Takeuchi, T., Koyasu, S., 2002. PI3K-mediated negative feedback regulation of IL-12 production in DCs. *Nat Immunol*, 3, 875–881.
- Fung, C., Naughton, S., Turnbull, L., Tingpej, P., Rose, B., Arthur, J., Hu, H., Harmer, C., Harbour, C., Hassett, D.J., Whitchurch, C.B., Manos, J., 2010. Gene expression of *Pseudomonas aeruginosa* in a mucin-containing synthetic growth medium mimicking cystic fibrosis lung sputum. *J Med Microbiol*, 59, 1089–1100.
- Ganesan, S., Sajjan, U.S., 2011. Host evasion by *Burkholderia cenocepacia*. *Front Cell Infect Microbiol*, 1, 25.
- Ganeshnarayan, K., Shah, S.M., Libera, M.R., Santostefano, A., Kaplan, J.B., 2009. Poly-N-acetylglucosamine matrix polysaccharide impedes fluid convection and transport of the cationic surfactant cetylpyridinium chloride through bacterial biofilms. *Appl Environ Microbiol*, 75, 1308–1314.
- Gao, W., Xiong, Y., Li, Q., Yang, H., 2017. Inhibition of Toll-Like receptor signaling as a promising therapy for inflammatory diseases: a journey from molecular to nano therapeutics. *Front Physiol*, 8, 508.
- García, J.R., Jaumann, F., Schulz, S., Krause, A., Rodríguez-Jiménez, J., Forssmann, U., Adermann, K., Klüver, E., Vogelmeier, C., Becker, D., Hedrich, R., Forssmann, W.G., Bals, R., 2001. Identification of a novel, multifunctional  $\beta$ -defensin (human  $\beta$ -defensin 3) with specific antimicrobial activity. Its interaction with plasma membranes of *Xenopus* oocytes and the induction of macrophage chemoattraction. *Cell Tissue Res*, 306, 257–264.
- García-del Portillo, F., Calvo, E., D'Orazio, V., Pucciarelli, M.G., 2011. Association of ActA to peptidoglycan revealed by cell wall proteomics of intracellular *Listeria monocytogenes*. *J Biol Chem*, 286, 34675–34689.
- Gardete, S., Tomasz, A., 2014. Mechanisms of vancomycin resistance in *Staphylococcus aureus*. *J Clin Invest*, 124, 2836–2840.
- Garzoni, C., Kelley, W.L., 2009. *Staphylococcus aureus*: new evidence for intracellular persistence. *Trends Microbiol*, 17, 59–65.

- Gaupp, R., Ledala, N., Somerville, G., 2012. Staphylococcal response to oxidative stress. *Front Cell Infect Microbiol*, 2, 33.
- Geiger, T., François, P., Liebeke, M., Fraunholz, M., Goerke, C., Krismer, B., Schrenzel, J., Lalk, M., Wolz, C., 2012. The stringent response of *Staphylococcus aureus* and its impact on survival after phagocytosis through the induction of intracellular PSMs expression. *PLoS Pathog*, 8, e1003016.
- Genestier, A.-L., Michallet, M.-C., Prévost, G., Bellot, G., Chalabreysse, L., Peyrol, S., Thivolet, F., Etienne, J., Lina, G., Vallette, F.M., Vandenesch, F., Genestier, L., 2005. *Staphylococcus aureus* Pantón-Valentine leukocidin directly targets mitochondria and induces Bax-independent apoptosis of human neutrophils. *J Clin Invest*, 115, 3117–3127.
- Georas, S.N., Rezaee, F., 2014. Epithelial barrier function: At the front line of asthma immunology and allergic airway inflammation. *J Allergy Clin Immunol*, 134, 509–520.
- Gessain, G., Tsai, Y.-H., Travier, L., Bonazzi, M., Grayo, S., Cossart, P., Charlier, C., Disson, O., Lecuit, M., 2015. PI3-kinase activation is critical for host barrier permissiveness to *Listeria monocytogenes*. *J Exp Med*, 212, 165–183.
- Gillette, D.D., Shah, P.A., Cremer, T., Gavrilin, M.A., Besecker, B.Y., Sarkar, A., Knoell, D.L., Cormet-Boyaka, E., Wewers, M.D., Butchar, J.P., Tridandapani, S., 2013. Analysis of human bronchial epithelial cell proinflammatory response to *Burkholderia cenocepacia* infection. *J Biol Chem*, 288, 3691–3695.
- Goldberg, J.B., Ganesan, S., Comstock, A.T., Zhao, Y., Sajjan, U.S., 2011. Cable pili and the associated 22 kDa adhesin contribute to *Burkholderia cenocepacia* persistence *in vivo*. *PLoS ONE*, 6, e22435.
- Gomez, M.I., Lee, A., Reddy, B., Muir, A., Soong, G., Pitt, A., Cheung, A., Prince, A., 2004. *Staphylococcus aureus* protein A induces airway epithelial inflammatory responses by activating TNFR1. *Nat Med*, 10, 842–848.
- Gomez, M.I., Seaghdha, M.O., Prince, A.S., 2007. *Staphylococcus aureus* protein A activates TACE through EGFR-dependent signaling. *EMBO J*, 26, 701–709.
- Gomez, M.I. and Prince, A., 2007. Airway epithelial cell signaling in response to bacterial pathogens. *Pediatr Pulmonol*, 43, 11–19.
- Gottschalk, R.A., Martins, A.J., Angermann, B.R., Dutta, B., Ng, C.E., Uderhardt, S., Tsang, J.S., Fraser, I.D.C., Meier-Schellersheim, M., Germain, R.N., 2016. Distinct NF- $\kappa$ B and MAPK activation thresholds uncouple steady-state microbe sensing from anti-pathogen inflammatory responses. *Cell Syst*, 2, 378–390.
- Govan, J.R.W., Brown, P.H., Maddinson, J., Doherty, C.J., Nelson, J.W., Dodd, M., Greening, A.P., Webb, A.K., 1993. Evidence for transmission of *Pseudomonas cepacia* by social contact in cystic fibrosis. *Lancet*, 342, 15–19.
- Grabiec, A.M., Hussell, T., 2016. The role of airway macrophages in apoptotic cell clearance following acute and chronic lung inflammation. *Semin Immunopathol*, 38, 409–423.
- Graille, M., Stura, E.A., Corper, A.L., Sutton, B.J., Taussig, M.J., Charbonnier, J.B., Silverman, G.J., 2000. Crystal structure of a *Staphylococcus aureus* protein A domain complexed with the Fab fragment of a human IgM antibody: structural basis for recognition of B-cell receptors and superantigen activity. *Proc Natl Acad Sci*, 97, 5399–5404.
- Grainger, C.I., Greenwell, L.L., Lockley, D.J., Martin, G.P., Forbes, B., 2006. Culture of Calu-3 cells at the air interface provides a representative model of the airway epithelial barrier. *Pharm Res*, 23, 1482–1490.

- Granchelli, A.M., Adler, F.R., Keogh, R.H., Kartsonaki, C., Cox, D.R., Liou, T.G., 2018. Microbial Interactions in the Cystic Fibrosis Airway. *J Clin Microbiol*, 56, e00354-18.
- Greenberg, J.A., Hrusch, C.L., Jaffery, M.R., David, M.Z., Daum, R.S., Hall, J.B., Kress, J.P., Sperling, A.I., Verhoef, P.A., 2018. Distinct T-helper cell responses to *Staphylococcus aureus* bacteremia reflect immunologic comorbidities and correlate with mortality. *Crit Care*, 22, 107.
- Gruenert, D.C., Willems, M., Cassiman, J.J., Frizzell, R.A., 2004. Established cell lines used in cystic fibrosis research. *J Cyst Fibros*, 3 Suppl 2, 191–196.
- Gu, B., Kelesidis, T., Tsiodras, S., Hindler, J., Humphries, R.M., 2012. The emerging problem of linezolid-resistant *Staphylococcus*. *J Antimicrob Chemother*, 68, 4–11.
- Guilhen, C., Forestier, C., Balestrino, D., 2017. Biofilm dispersal: multiple elaborate strategies for dissemination of bacteria with unique properties. *Mol Microbiol*, 105, 188–210.
- Guyen-Maiorov, E., Keskin, O., Gursoy, A., Nussinov, R., 2015. A structural view of negative regulation of the Toll-like receptor-mediated inflammatory pathway. *Biophys J*, 109, 1214–1226.
- Haaber, J., Penadés, J.R., Ingmer, H., 2017. Transfer of Antibiotic Resistance in *Staphylococcus aureus*. *Trend Microbiol*, 25, 893–905.
- Hair, P.S., Echague, C.G., Sholl, A.M., Watkins, J.A., Geoghegan, J.A., Foster, T.J., Cunnion, K.M., 2010. Clumping factor A interaction with complement factor i increases C3b cleavage on the bacterial surface of *Staphylococcus aureus* and decreases complement-mediated phagocytosis. *Infect Immun*, 78, 1717–1727.
- Hajishengallis, G., Reis, E.S., Mastellos, D.C., Ricklin, D., Lambris, J.D., 2017. Novel mechanisms and functions of complement. *Nat Immunol*, 18, 1288–1298.
- Hall, C.W., Mah, T.-F., 2017. Molecular mechanisms of biofilm-based antibiotic resistance and tolerance in pathogenic bacteria. *FEMS Microbiol Rev*, 41, 276–301.
- Ham, H., Sreelatha, A., Orth, K., 2011. Manipulation of host membranes by bacterial effectors. *Nat Rev Micro*, 9, 635–646.
- Hamad, M.A., Di Lorenzo, F., Molinaro, A., Valvano, M.A., 2012. Aminoarabinose is essential for lipopolysaccharide export and intrinsic antimicrobial peptide resistance in *Burkholderia cenocepacia*. *Mol Microbiol*, 85, 962–974.
- Hamaguchi, R., Haginaka, J., Tanimoto, T., Kuroda, Y., 2014. Maintenance of luminal pH and protease activity in lysosomes/late endosomes by vacuolar ATPase in chlorpromazine-treated RAW264 cells accumulating phospholipids. *Cell Biol Toxicol*, 30, 67–77.
- Hammel, M., Sfyroera, G., Pyrpasopoulos, S., Ricklin, D., Ramyar, K.X., Pop, M., Jin, Z., Lambris, J.D., Geisbrecht, B.V., 2007a. Characterization of Ehp, a secreted complement inhibitory protein from *Staphylococcus aureus*. *J Biol Chem*, 282, 30051–30061.
- Hammel, M., Sfyroera, G., Ricklin, D., Magotti, P., Lambris, J.D., Geisbrecht, B.V., 2007b. A structural basis for complement inhibition by *Staphylococcus aureus*. *Nat Immunol*, 8, 430–437.
- Han, J., Tschernutter, V., Yang, J., Eckle, T., Borchers, C.H., 2013. Analysis of selected sugars and sugar phosphates in mouse heart tissue by reductive amination and liquid chromatography-electrospray ionization mass spectrometry. *Anal chem*, 85, 5965–5973.
- Hanke, M.L., Kielian, T., 2012. Deciphering mechanisms of staphylococcal biofilm evasion of host immunity. *Front Cell Infect Microbiol*, 2, 62.

- Hardison, S.E., Brown, G.D., 2012. C-type lectin receptors orchestrate antifungal immunity. *Nat Immunol*, 13, 817–822.
- Hartleib, J., Kohler, N., Dickinson, R.B., Chhatwal, G.S., Sixma, J.J., Hartford, O.M., Foster, T.J., Peters, G., Kehrel, B.E., Herrmann, M., 2000. Protein A is the von Willebrand factor binding protein on *Staphylococcus aureus*. *Blood*, 96, 2149–2156.
- Hartsock, A., Nelson, W.J., 2008. Adherens and tight junctions: structure, function and connections to the actin cytoskeleton. *Biochim Biophys Acta*, 1778, 660–669.
- Hassett, D.J., Borchers, M.T., Panos, R.J., 2014. Chronic obstructive pulmonary disease: evaluation from clinical, immunological and bacterial pathogenesis perspectives. *J Microbiol*, 52, 211–226.
- Haupt, K., Reuter, M., van den Elsen, J., Burman, J., Hälbig, S., Richter, J., Skerka, C., Zipfel, P.F., 2008. The *Staphylococcus aureus* protein Sbi acts as a complement inhibitor and forms a tripartite complex with host complement Factor H and C3b. *PLoS Pathog*, 4, e1000250.
- Hauser, A.R., Jain, M., Bar-Meir, M., McColley, S.A., 2011. Clinical significance of microbial infection and adaptation in cystic fibrosis. *Clin Microbiol Rev*, 24, 29–70.
- Hauser, N., Orsini, J., 2015. Cepacia syndrome in a non-cystic fibrosis patient. *Case Rep Infect Dis*, 2015, 537627–4.
- Hawkins, P.T., Stephens, L.R., 2015. PI3K signalling in inflammation. *Biochim Biophys Acta*, 1851, 882–897.
- Hayashi, F., Smith, K.D., Ozinsky, A., Hawn, T.R., Yi, E.C., Goodlett, D.R., Eng, J.K., Akira, S., Underhill, D.M., Aderem, A., 2001. The innate immune response to bacterial flagellin is mediated by Toll-like receptor 5. *Nature*, 410, 1099–1103.
- Häcker, H., Redecke, V., Blagoev, B., Kratchmarova, I., Hsu, L.-C., Wang, G.G., Kamps, M.P., Raz, E., Wagner, H., Häcker, G., Mann, M., Karin, M., 2006. Specificity in Toll-like receptor signalling through distinct effector functions of TRAF3 and TRAF6. *Nature*, 439, 204–207.
- Häcker, H., Tseng, P.-H., Karin, M., 2011. Expanding TRAF function: TRAF3 as a tri-faced immune regulator. *Nat Rev Immunol*, 11, 457–468.
- He, H., Genovese, K.J., Nisbet, D.J., Kogut, M.H., 2006. Profile of Toll-like receptor expressions and induction of nitric oxide synthesis by Toll-like receptor agonists in chicken monocytes. *Mol Immunol*, 43, 783–789.
- Heil, F., Hemmi, H., Hochrein, H., Ampenberger, F., Kirschning, C., Akira, S., Lipford, G., Wagner, H., Bauer, S., 2004. Species-specific recognition of single-stranded RNA via Toll-like receptor 7 and 8. *Science*, 303, 1526–1529.
- Henrick BM, Yao X-D, Taha AY, German JB, Rosenthal KL. Insights into soluble Toll-like receptor 2 as a downregulator of virally induced inflammation. *Front Immunol*, 7, 291.
- Herasimenka, Y., Cescutti, P., Impallomeni, G., Campana, S., Taccetti, G., Ravenni, N., Zanetti, F., Rizzo, R., 2007. Exopolysaccharides produced by clinical strains belonging to the *Burkholderia cepacia* complex. *J Cyst Fibros*, 6, 145–152.
- Hernandez, L.D., Pypaert, M., Flavell, R.A., Galán, J.E., 2003. A *Salmonella* protein causes macrophage cell death by inducing autophagy. *J Cell Biol*, 163, 1123–1131.
- Hiemstra, P.S., 2002. Novel roles of protease inhibitors in infection and inflammation. *Biochem Soc Trans*, 30, 116–120.
- Hilmi, D., Parcina, M., Stollewerk, D., Ostrop, J., Josten, M., Meilaender, A., Zaehring, U., Wichelhaus, T.A., Bierbaum, G., Heeg, K., Wolz, C., Bekeredjian-Ding, I., 2014.

- Heterogeneity of Host TLR2 Stimulation by *Staphylococcus aureus* Isolates. *PLoS ONE*, 9, e96416.
- Hodge, S., Hodge, G., Scicchitano, R., Reynolds, P.N., Holmes, M., 2003. Alveolar macrophages from subjects with chronic obstructive pulmonary disease are deficient in their ability to phagocytose apoptotic airway epithelial cells. *Immunol Cell Biol*, 81, 289–296.
- Hoffman, L.R., Déziel, E., D'Argenio, D.A., Lépine, F., Emerson, J., McNamara, S., Gibson, R.L., Ramsey, B.W., Miller, S.I., 2006. Selection for *Staphylococcus aureus* small-colony variants due to growth in the presence of *Pseudomonas aeruginosa*. *Proc Natl Acad Sci*, 103, 19890–19895.
- Holden, M.T.G., Feil, E.J., Lindsay, J.A., Peacock, S.J., Day, N.P.J., Enright, M.C., Foster, T.J., Moore, C.E., Hurst, L., Atkin, R., Barron, A., Bason, N., Bentley, S.D., Chillingworth, C., Chillingworth, T., Churcher, C., Clark, L., Corton, C., Cronin, A., Doggett, J., Dowd, L., Feltwell, T., Hance, Z., Harris, B., Hauser, H., Holroyd, S., Jagels, K., James, K.D., Lennard, N., Line, A., Mayes, R., Moule, S., Mungall, K., Ormond, D., Quail, M.A., Rabinowitsch, E., Rutherford, K., Sanders, M., Sharp, S., Simmonds, M., Stevens, K., Whitehead, S., Barrell, B.G., Spratt, B.G., Parkhill, J., 2004. Complete genomes of two clinical *Staphylococcus aureus* strains: evidence for the rapid evolution of virulence and drug resistance. *Proc Natl Acad Sci*, 101, 9786–9791.
- Holden, M.T.G., Seth-Smith, H.M.B., Crossman, L.C., Sebaihia, M., Bentley, S.D., Cerdeno-Tarraga, A.M., Thomson, N.R., Bason, N., Quail, M.A., Sharp, S., Cherevach, I., Churcher, C., Goodhead, I., Hauser, H., Holroyd, N., Mungall, K., Scott, P., Walker, D., White, B., Rose, H., Iversen, P., Mil-Homens, D., Rocha, E.P.C., Fialho, A.M., Baldwin, A., Dowson, C., Barrell, B.G., Govan, J.R., Vandamme, P., Hart, C.A., Mahenthiralingam, E., Parkhill, J., 2009. The genome of *Burkholderia cenocepacia* J2315, an epidemic pathogen of cystic fibrosis patients. *J Bacterio*, 191, 261–277.
- Holmes, A., Nolan, R., Taylor, R., Finley, R., Riley, M., Jiang, R.Z., Steinbach, S., Goldstein, R., 1999. An epidemic of *Burkholderia cepacia* transmitted between patients with and without cystic fibrosis. *J Infect Dis*, 179, 1197–1205.
- Holt, P.G., Oliver, J., Bilyk, N., McMenamin, C., McMenamin, P.G., Kraal, G., Thepen, T., 1993. Downregulation of the antigen presenting cell function(s) of pulmonary dendritic cells *in vivo* by resident alveolar macrophages. *J Exp Med*, 177, 397–407.
- Hoving, J.C., Wilson, G.J., Brown, G.D., 2014. Signalling C-Type lectin receptors, microbial recognition and immunity. *Cell Microbiol*, 16, 185–194.
- Hovsepian, E., Penas, F., Siffo, S., Mirkin, G.A., Goren, N.B., 2013. IL-10 inhibits the NF- $\kappa$ B and Erk/MAPK-mediated production of pro-inflammatory mediators by up-regulation of SOCS-3 in *Trypanosoma cruzi*-infected cardiomyocytes. *PLoS ONE*, 8, e79445.
- Hu, W., Jain, A., Gao, Y., Dozmorov, I.M., Mandraju, R., Wakeland, E.K., Pasare, C., 2015. Differential outcome of TRIF-mediated signaling in TLR4 and TLR3 induced DC maturation. *Proc Natl Acad Sci*, 112, 13994–13999.
- Huang, Q.-Q., Pope, R.M., 2009. The role of toll-like receptors in rheumatoid arthritis. *Curr Rheumatol Rep*, 11, 357–364.
- Huber, B., Riedel, K., Hentzer, M., Heydorn, A., Gotschlich, A., Givskov, M., Molin, S., Eberl, L., 2001. The cep quorum-sensing system of *Burkholderia cepacia* H111 controls biofilm formation and swarming motility. *Microbiology*, 147, 2517–2528.
- Humlicek, A.L., Manzel, L.J., Chin, C.L., Shi, L., Excoffon, K.J.D.A., Winter, M.C., Shasby, D.M., Look, D.C., 2007. Paracellular permeability restricts airway epithelial responses to selectively allow activation by mediators at the basolateral surface. *J Immunol*, 178, 6395–6403.

- Hussell, T., Bell, T.J., 2014. Alveolar macrophages: plasticity in a tissue-specific context. *Nat Rev Immunol*, 14, 81–93.
- Høiby, N., Bjarnsholt, T., Givskov, M., Molin, S.R., Ciofu, O., 2010. Antibiotic resistance of bacterial biofilms. *Int J Antimicrob Agents*, 35, 322–332.
- Illek, B., Fu, Z., Schwarzer, C., Banzon, T., Jalickee, S., Miller, S.S., Machen, T.E., 2008. Flagellin-stimulated Cl<sup>-</sup> secretion and innate immune responses in airway epithelia: role for p38. *Am J Physiol Lung Cell Mol Physiol*, 295, L531–L542.
- Inagaki-Ohara K., Kondo T., Ito M., Yoshimura A., 2013. SOCS, inflammation, and cancer. *JAK-STAT*, 2, e24053.
- Into, T., Kiura, K., Yasuda, M., Kataoka, H., Inoue, N., Hasebe, A., Takeda, K., Akira, S., Shibata, K.-I., 2004. Stimulation of human Toll-like receptor (TLR) 2 and TLR6 with membrane lipoproteins of *Mycoplasma fermentans* induces apoptotic cell death after NF- $\kappa$ B activation. *Cell Microbiol*, 6, 187–199.
- Ioannidis, I., Ye, F., McNally, B., Willette, M., Flano, E., 2013. Toll-like receptor expression and induction of type I and type III interferons in primary airway epithelial cells. *J Virol*, 87, 3261–3270.
- Ip, W.K.E., Sokolovska, A., Charriere, G.M., Boyer, L., Dejardin, S., Cappillino, M.P., Yantosca, L.M., Takahashi, K., Moore, K.J., Lacy-Hulbert, A., Stuart, L.M., 2010. Phagocytosis and phagosome acidification are required for pathogen processing and MyD88-dependent responses to *Staphylococcus aureus*. *J Immunol*, 184, 7071–7081.
- Isles, A., Maclusky, I., Corey, M., Gold, R., Prober, C., Fleming, P., Levison, H., 1984. *Pseudomonas cepacia* infection in cystic fibrosis: an emerging problem. *J Pediatr*, 104, 206–210.
- Itoh, S., Hamada, E., Kamoshida, G., Yokoyama, R., Takii, T., Onozaki, K., Tsuji, T., 2010. Staphylococcal superantigen-like protein 10 (SSL10) binds to human immunoglobulin G (IgG) and inhibits complement activation via the classical pathway. *Mol Immunol*, 47, 932–938.
- Itoh, S., Yokoyama, R., Kamoshida, G., Fujiwara, T., Okada, H., Takii, T., Tsuji, T., Fujii, S., Hashizume, H., Onozaki, K., 2013. Staphylococcal superantigen-like protein 10 (SSL10) inhibits blood coagulation by binding to prothrombin and factor Xa via their gamma-carboxyglutamic acid (Gla) domain. *J Biol Chem*, 288, 21569–21580.
- Iwaki, D., Mitsuzawa, H., Murakami, S., Sano, H., Konishi, M., Akino, T., Kuroki, Y., 2002. The extracellular Toll-like receptor 2 domain directly binds peptidoglycan derived from *Staphylococcus aureus*. *J Biol Chem*, 277, 24315–24320.
- Iwami, K.I., Matsuguchi, T., Masuda, A., Kikuchi, T., Musikacharoen, T., Yoshikai, Y., 2000. Cutting edge: naturally occurring soluble form of mouse Toll-like receptor 4 inhibits lipopolysaccharide signaling. *J Immunol*, 165, 6682–6686.
- Jahani, S., Shakiba, A., Jahani, L., 2015. The antimicrobial effect of lactoferrin on Gram-negative and Gram-positive bacteria. *Int J Infect*, 2, e27954.
- Jakubzick, C., Tacke, F., Llodra, J., Van Rooijen, N., Randolph, G.J., 2006. Modulation of dendritic cell trafficking to and from the airways. *J Immunol*, 176, 3578–3584.
- Jevons, M.P., 1961. “Celbenin” resistant staphylococci. *Br Med J*, 1, 124–125.
- Jia, H.P., Kline, J.N., Penisten, A., Apicella, M.A., Gioannini, T.L., Weiss, J., McCray, P.B., 2004. Endotoxin responsiveness of human airway epithelia is limited by low expression of MD-2. *Am J Physiol Lung Cell Mol Physiol*, 287, L428–37.

- Jin, T., Bokarewa, M., Foster, T., Mitchell, J., Higgins, J., Tarkowski, A., 2004. *Staphylococcus aureus* resists human defensins by production of staphylokinase, a novel bacterial evasion mechanism. *J Immunol*, 172, 1169–1176.
- Joiner, K.A., Brown, E.J., Frank, M.M., 1984. Complement and bacteria: chemistry and biology in host defense. *Annu Rev Immunol*, 2, 461–491.
- Jones, A.M., Dodd, M.E., Govan, J.R.W., Barcus, V., Doherty, C.J., Morris, J., Webb, A.K., 2004. *Burkholderia cenocepacia* and *Burkholderia multivorans*: influence on survival in cystic fibrosis. *Thorax*, 59, 948–951.
- Jones, R.N., 2010. Microbial etiologies of hospital-acquired bacterial pneumonia and ventilator-associated bacterial pneumonia. *Clin Infect Dis*, 51, S81–S87.
- Jongerijs, I., Garcia, B.L., Geisbrecht, B.V., van Strijp, J.A.G., Rooijackers, S.H.M., 2010a. Convertase inhibitory properties of Staphylococcal extracellular complement-binding protein. *J Biol Chem*, 285, 14973–14979.
- Jongerijs, I., Kohl, J., Pandey, M.K., Ruyken, M., van Kessel, K.P.M., van Strijp, J.A.G., Rooijackers, S.H.M., 2007. Staphylococcal complement evasion by various convertase-blocking molecules. *J Exp Med*, 204, 2461–2471.
- Jongerijs, I., Puister, M., Wu, J., Ruyken, M., van Strijp, J.A.G., Rooijackers, S.H.M., 2010b. Staphylococcal complement inhibitor modulates phagocyte responses by dimerization of convertases. *J Immunol*, 184, 420–425.
- Jonsson, K., Signas, C., Muller, H.-P., Lindberg, M., 1991. Two different genes encode fibronectin binding proteins in *Staphylococcus aureus*. The complete nucleotide sequence and characterization of the second gene. *Eur J Biochem*, 202, 1041–1048.
- Jouault, T., Ibata-Ombetta, S., Takeuchi, O., Trinel, P.-A., Sacchetti, P., Lefebvre, P., Akira, S., Poulain, D., 2003. *Candida albicans* phospholipomannan is sensed through Toll-like receptors. *J Infect Dis*, 188, 165–172.
- Jovanovic, M., Rooney, M.S., Mertins, P., Przybylski, D., Chevrier, N., Satija, R., Rodriguez, E.H., Fields, A.P., Schwartz, S., Raychowdhury, R., Mumbach, M.R., Eisenhaure, T., Rabani, M., Gennert, D., Lu, D., Delorey, T., Weissman, J.S., Carr, S.A., Hacohen, N., Regev, A., 2015. Dynamic profiling of the protein life cycle in response to pathogens. *Science*, 347, 1259038.
- Jung, K.Y., Cha, J.D., Lee, S.H., Woo, W.H., Lim, D.S., Choi, B.K., Kim, K.J., 2001. Involvement of staphylococcal protein A and cytoskeletal actin in *Staphylococcus aureus* invasion of cultured human oral epithelial cells. *J Med Microbiol*, 50, 35–41.
- Jusko, M., Potempa, J., Kantyka, T., Bielecka, E., Miller, H.K., Kalinska, M., Dubin, G., Garred, P., Shaw, L.N., Blom, A.M., 2014. Staphylococcal proteases aid in evasion of the human complement system. *J Innate Immun*, 6, 31–46.
- Kaci, G., Goudercourt, D., Dennin, V., Pot, B., Doré, J., Ehrlich, S.D., Renault, P., Blottière, H.M., Daniel, C., Delorme, C., 2014. Anti-inflammatory properties of *Streptococcus salivarius*, a commensal bacterium of the oral cavity and digestive tract. *Appl Environ Microbiol*, 80, 928–934.
- Kaci, G., Lakhdari, O., Doré, J., Ehrlich, S.D., Renault, P., Blottière, H.M., Delorme, C., 2011. Inhibition of the NF- $\kappa$ B pathway in human intestinal epithelial cells by commensal *Streptococcus salivarius*. *Appl Environ Microbiol*, 77, 4681–4684.
- Kagan, J.C., Su, T., Horng, T., Chow, A., Akira, S., Medzhitov, R., 2008. TRAM couples endocytosis of Toll-like receptor 4 to the induction of interferon- $\beta$ . *Nat Immunol*, 9, 361–368.



- Kao, C.-Y., Chen, Y., Thai, P., Wachi, S., Huang, F., Kim, C., Harper, R.W., Wu, R., 2004. IL-17 markedly up-regulates  $\beta$ -defensin-2 expression in human airway epithelium via JAK and NF- $\kappa$ B signaling pathways. *J Immunol*, 173, 3482–3491.
- Kaplan, J.B., 2010. Biofilm dispersal: mechanisms, clinical implications, and potential therapeutic uses. *J Dent Res*, 89, 205–218.
- Kato K, Lillehoj E P, Kim K C. *Pseudomonas aeruginosa* stimulates tyrosine phosphorylation of and TLR5 association with the MUC1 cytoplasmic tail through EGFR activation. *Inflamm Res*, 2016, 65(3): 225-233.
- Kawai, T., Akira, S., 2007. Signaling to NF- $\kappa$ B by Toll-like receptors. *Trends Mol Med*, 13, 460–469.
- Kawai, T., Akira, S., 2010. The role of pattern-recognition receptors in innate immunity: update on Toll-like receptors. *Nat Immunol*, 11, 373–384.
- Kawasaki, T. & Kawai, T., 2014. Toll-like receptor signaling pathways. *Front Immunol*, 5, 461.
- Kayagaki, N., Phung, Q., Chan, S., Chaudhari, R., Quan, C., O'Rourke, K.M., Eby, M., Pietras, E., Cheng, G., Bazan, J.F., Zhang, Z., Arnott, D., Dixit, V.M., 2007. DUBA: a deubiquitinase that regulates type I interferon production. *Science*, 318, 1628–1632.
- Kaza, S.K., McClean, S., Callaghan, M., 2010. IL-8 released from human lung epithelial cells induced by cystic fibrosis pathogens *Burkholderia cepacia* complex affects the growth and intracellular survival of bacteria. *Int J Med Microbiol*, 301, 26–33.
- Keith, K.E., Hynes, D.W., Sholdice, J.E., Valvano, M.A., 2009. Delayed association of the NADPH oxidase complex with macrophage vacuoles containing the opportunistic pathogen *Burkholderia cenocepacia*. *Microbiology*, 155, 1004–1015.
- Kenna, D.T.D., Lilley, D., Coward, A., Martin, K., Perry, C., Pike, R., Hill, R., Turton, J.F., 2017. Prevalence of *Burkholderia* species, including members of *Burkholderia cepacia* complex, among UK cystic and non-cystic fibrosis patients. *J Med Microbiol*, 66, 490–501.
- Khiem, D., Cyster, J.G., Schwarz, J.J., Black, B.L., 2008. A p38 MAPK-MEF2C pathway regulates B-cell proliferation. *Proc Natl Acad Sci*, 105, 17067–17072.
- Kiedrowski, M.R., Paharik, A.E., Ackermann, L.W., Shelton, A.U., Singh, S.B., Starnes, T.D., Horswill, A.R., 2016. Development of an in vitro colonization model to investigate *Staphylococcus aureus* interactions with airway epithelia. *Cell Microbiol*, 18, 720–732.
- Kierbel, A., Gassama-Diagne, A., Mostov, K., Engel, J.N., 2005. The phosphoinositol-3-kinase–protein kinase B/Akt pathway is critical for *Pseudomonas aeruginosa* strain PAK internalization. *Mol Biol Cell*, 16, 2577–2585.
- Kilburn, K.H., 1968. A hypothesis for pulmonary clearance and its implications. *Am Rev Respir Dis*, 98, 449–463.
- Kim, J.Y., Sajjan, U.S., Krasan, G.P., LiPuma, J.J., 2005. Disruption of tight junctions during traversal of the respiratory epithelium by *Burkholderia cenocepacia*. *Infect Immun*, 73, 7107–7112.
- Kim, M., Ashida, H., Ogawa, M., Yoshikawa, Y., Mimuro, H., Sasakawa, C., 2010. Bacterial interactions with the host epithelium. *Cell Host Microbe*, 8, 20–35.
- Kim, Y.K., Shin, J.-S., Nahm, M.H., 2016. NOD-like receptors in infection, immunity, and diseases. *Yonsei Med J*, 57, 5–14.
- Knodler, L.A., Finlay, B.B., Steele-Mortimer, O., 2005. The *Salmonella* effector protein SopB protects epithelial cells from apoptosis by sustained activation of Akt. *J Biol Chem*, 280, 9058–9064.

- Kobayashi, K., Hernandez, L.D., Galán, J.E., Janeway, C.A.J., Medzhitov, R., Flavell, R.A., 2002. IRAK-M is a negative regulator of Toll-like receptor signaling. *Cell*, 110, 191–202.
- Kojima, T., Go, M., Takano, K.-I., Kurose, M., Ohkuni, T., Koizumi, J.-I., Kamekura, R., Ogasawara, N., Masaki, T., Fuchimoto, J., Obata, K., Hirakawa, S., Nomura, K., Keira, T., Miyata, R., Fujii, N., Tsutsumi, H., Himi, T., Sawada, N., 2013. Regulation of tight junctions in upper airway epithelium. *Biomed Res Int*, 2013, 947072–11.
- Kondo, T., Kawai, T., Akira, S., 2012a. Dissecting negative regulation of Toll-like receptor signaling. *Trends Immunol*, 33, 449–458.
- Konrad, S., Ali, S.R., Wiege, K., Syed, S.N., Engling, L., Piekorz, R.P., Hirsch, E., Nürnberg, B., Schmidt, R.E., Gessner, J., 2008. Phosphoinositide 3-kinases  $\gamma$  and  $\delta$ , linkers of coordinate C5a receptor-Fc $\gamma$  receptor activation and immune complex-induced inflammation. *J Biol Chem*, 283, 33296–33303.
- Kooi, C., Corbett, C.R., Sokol, P.A., 2005. Functional analysis of the *Burkholderia cenocepacia* ZmpA metalloprotease. *J Bacterio*, 187, 4421–4429.
- Kooi, C., Sokol, P.A., 2009. *Burkholderia cenocepacia* zinc metalloproteases influence resistance to antimicrobial peptides. *Microbiology*, 155, 2818–2825.
- Kooi, C., Subsin, B., Chen, R., Pohorelic, B., Sokol, P.A., 2006. *Burkholderia cenocepacia* ZmpB is a broad-specificity zinc metalloprotease involved in virulence. *Infect Immun*, 74, 4083–4093.
- Kopf, M., Schneider, C., Nobs, S.P., 2015. The development and function of lung-resident macrophages and dendritic cells. *Nat Immunol*, 16, 36–44.
- Korea, C.G., Balsamo, G., Pezzicoli, A., Merakou, C., Tavarini, S., Bagnoli, F., Serruto, D., Unnikrishnan, M., 2014. Staphylococcal Esx proteins modulate apoptosis and release of intracellular *Staphylococcus aureus* during infection in epithelial cells. *Infect Immun*, 82, 4144–4153.
- Korneev, K.V., Atretkhany, K.-S.N., Drutskaya, M.S., Grivennikov, S.I., Kuprash, D.V., Nedospasov, S.A., 2017. TLR-signaling and proinflammatory cytokines as drivers of tumorigenesis. *Cytokine*, 89, 127–135.
- Kotrange, S., Kopp, B., Akhter, A., Abdelaziz, D., Abu Khweek, A., Caution, K., Abdulrahman, B., Wewers, M.D., McCoy, K., Marsh, C., Loutet, S.A., Ortega, X., Valvano, M.A., Amer, A.O., 2011. *Burkholderia cenocepacia* O polysaccharide chain contributes to caspase-1-dependent IL-1 $\beta$  production in macrophages. *J Leukoc Biol*, 89, 481–488.
- Koymans, K.J., Goldmann, O., Karlsson, C.A.Q., Sital, W., Thänert, R., Bisschop, A., Vrieling, M., Malmström, J., van Kessel, K.P.M., de Haas, C.J.C., van Strijp, J.A.G., Medina, E., 2017. The TLR2 antagonist Staphylococcal superantigen-like protein 3 acts as a virulence factor to promote bacterial pathogenicity *in vivo*. *J Innate Immun*, 9, 561–573.
- Krachler, A.M., Woolery, A.R., Orth, K., 2011. Manipulation of kinase signaling by bacterial pathogens. *J Cell Biol*, 195, 1083–1092.
- Kraft, R., Herndon, D.N., Finnerty, C.C., Cox, R.A., Song, J., Jeschke, M.G., 2015. Predictive value of IL-8 for sepsis and severe infections after burn injury: a clinical study. *Shock*, 43, 222–227.
- Kreft, M.E., Jerman, U.D., Lasiča, E., Hevir-Kene, N., Rižner, T.L., Peternel, L., Kristan, K., 2015. The characterization of the human cell line Calu-3 under different culture conditions and its use as an optimized *in vitro* model to investigate bronchial epithelial function. *Eur J Pharm Sci*, 69, 1–9.

- Kretschmer, D., Gleske, A.-K., Rautenberg, M., Wang, R., Köberle, M., Bohn, E., Schöneberg, T., Rabiet, M.-J., Boulay, F., Klebanoff, S.J., van Kessel, K.A., van Strijp, J.A., Otto, M., Peschel, A., 2010. Human formyl peptide receptor 2 senses highly pathogenic *Staphylococcus aureus*. *Cell Host Microbe*, 7, 463–473.
- Krismer, B., Peschel, A., 2011. Does *Staphylococcus aureus* nasal colonization involve biofilm formation? *Future Microbiol*, 6, 489–493.
- Kruger, P., Saffarzadeh, M., Weber, A.N.R., Rieber, N., Radsak, M., Bernuth, von, H., Benarafa, C., Roos, D., Skokowa, J., Hartl, D., 2015. Neutrophils: between host defence, immune modulation, and tissue injury. *PLoS Pathog*, 11, e1004651.
- Kubica, M., Guzik, K., Koziel, J., Zarebski, M., Richter, W., Gajkowska, B., Golda, A., Maciag-Gudowska, A., Brix, K., Shaw, L., Foster, T., Potempa, J., 2008. A potential new pathway for *Staphylococcus aureus* dissemination: the silent survival of *S. aureus* phagocytosed by human monocyte-derived macrophages. *PLoS ONE*, 3, e1409.
- Kukavica-Ibrulj, I., Facchini, M., Cigana, C., Levesque, R.C., Bragonzi, A., 2014. Assessing *Pseudomonas aeruginosa* virulence and the host response using murine models of acute and chronic lung infection. *Methods Mol Biol*, 1149, 757–771.
- Kulkarni, H.S., Liszewski, M.K., Brody, S.L., Atkinson, J.P., 2018. The complement system in the airway epithelium: an overlooked host defense mechanism and therapeutic target? *J Allergy Clin Immunol*, 141, 1582–1586.e1.
- Kumar, Brijesh, Cardona, S.T., 2016. Synthetic cystic fibrosis sputum medium regulates flagellar biosynthesis through the *flhF* gene in *Burkholderia cenocepacia*. *Front Cell Infect Microbiol*, 6, 5024.
- Kumar, Neha, David, M.Z., Boyle-Vavra, S., Sieth, J., Daum, R.S., 2015. High *Staphylococcus aureus* colonization prevalence among patients with skin and soft tissue infections and controls in an urban emergency department. *J Clin Microbiol*, 53, 810–815.
- Kumar, Nitin, Su, C.-C., Chou, T.-H., Radhakrishnan, A., Delmar, J.A., Rajashankar, K.R., Yu, E.W., 2017. Crystal structures of the *Burkholderia multivorans* hopanoid transporter HpnN. *Proc Natl Acad Sci*, 114, 6557–6562.
- Laarman, A.J., Mijnheer, G., Mootz, J.M., van Rooijen, W.J.M., Ruyken, M., Malone, C.L., Heezius, E.C., Ward, R., Milligan, G., van Strijp, J.A.G., de Haas, C.J.C., Horswill, A.R., van Kessel, K.P.M., Rooijackers, S.H.M., 2012. *Staphylococcus aureus* Staphopain A inhibits CXCR2-dependent neutrophil activation and chemotaxis. *EMBO J*, 31, 3607–3619.
- Laarman, A.J., Ruyken, M., Malone, C.L., van Strijp, J.A.G., Horswill, A.R., Rooijackers, S.H.M., 2011. *Staphylococcus aureus* metalloprotease aureolysin cleaves complement C3 to mediate immune evasion. *J Immunol*, 186, 6445–6453.
- Lamothe, J., Huynh, K.K., Grinstein, S., Valvano, M.A., 2007. Intracellular survival of *Burkholderia cenocepacia* in macrophages is associated with a delay in the maturation of bacteria-containing vacuoles. *Cell Microbiol*, 9, 40–53.
- Langjahr, P., Diaz-Jimenez, D., la Fuente, De, M., Rubio, E., Golenbock, D., Bronfman, F.C., Quera, R., Gonzalez, M.-J., Hermoso, M.A., 2014. Metalloproteinase-dependent TLR2 ectodomain shedding is involved in soluble Toll-like receptor 2 production. *PLoS ONE* 9, e104624.
- Lardi, M., Aguilar, C., Pedrioli, A., Omasits, U., Suppiger, A., Cárcamo-Oyarce, G., Schmid, N., Ahrens, C.H., Eberl, L., Pessi, G., 2015.  $\sigma$ 54-dependent response to nitrogen limitation and virulence in *Burkholderia cenocepacia* Strain H111. *Appl Environ Microbiol*, 81, 4077–4089.

- Le Bon, A., Etchart, N., Rossmann, C., Ashton, M., Hou, S., Gewert, D., Borrow, P., Tough, D.F., 2003. Cross-priming of CD8<sup>+</sup> T cells stimulated by virus-induced type I interferon. *Nat Immunol*, 4, 1009–1015.
- Leadbetter, E.A., Rifkin, I.R., Hohlbaum, A.M., Beaudette, B.C., Shlomchik, M.J., Marshak-Rothstein, A., 2002. Chromatin-IgG complexes activate B cells by dual engagement of IgM and Toll-like receptors. *Nature* 416, 603–607.
- LeBouder, E., Rey-Nores, J.E., Rushmere, N.K., Grigorov, M., Lawn, S.D., Affolter, M., Griffin, G.E., Ferrara, P., Schiffrin, E.J., Morgan, B.P., Labéta, M.O., 2003. Soluble forms of Toll-like receptor (TLR)2 capable of modulating TLR2 signaling are present in human plasma and breast milk. *J Immunol*, 171, 6680–6689.
- Lee, S.M.-Y., Yip, T.-F., Yan, S., Jin, D.-Y., Wei, H.-L., Guo, R.-T., Peiris, J.S.M., 2018. Recognition of double-stranded RNA and regulation of interferon pathway by Toll-like receptor 10. *Front Immunol*, 9, 516.
- Leech, J.M., Lacey, K.A., Mulcahy, M.E., Medina, E., McLoughlin, R.M., 2017. IL-10 plays opposing roles during *Staphylococcus aureus* systemic and localized infections. *J Immunol*, 198, 2352–2365.
- Lehnert, B.E., Valdez, Y.E., Holland, L.M., 1985. Pulmonary macrophages: alveolar and interstitial populations. *Exp Lung Res*, 9, 177–190.
- Leid, J.G., Willson, C.J., Shirliff, M.E., Hassett, D.J., Parsek, M.R., Jeffers, A.K., 2005. The exopolysaccharide alginate protects *Pseudomonas aeruginosa* biofilm bacteria from IFN- $\gamma$ -mediated macrophage killing. *J Immunol*, 175, 7512–7518.
- Lewenza, S., Conway, B., Greenberg, E.P., Sokol, P.A., 1999. Quorum sensing in *Burkholderia cepacia*: identification of the LuxRI homologs CepRI. *J Bacterio*, 181, 748–756.
- Lewis, R.S., Kolesnik, T.B., Kuang, Z., D'Cruz, A.A., Blewitt, M.E., Masters, S.L., Low, A., Willson, T., Norton, R.S., Nicholson, S.E., 2011. TLR regulation of SPSB1 controls inducible nitric oxide synthase induction. *J Immunol*, 187, 3798–3805.
- Li Jeon, N., Baskaran, H., Dertinger, S.K.W., Whitesides, G.M., Van De Water, L., Toner, M., 2002. Neutrophil chemotaxis in linear and complex gradients of interleukin-8 formed in a microfabricated device. *Nat Biotechnol*, 20, 826–830.
- Li, Xiaotong, Udagawa, N., Takami, M., Sato, N., Kobayashi, Y., Takahashi, N., 2003. p38 mitogen-activated protein kinase is crucially involved in osteoclast differentiation but not in cytokine production, phagocytosis, or dendritic cell differentiation of bone marrow macrophages. *Endocrinology*, 144, 4999–5005.
- Li, Xinyan, Jiang, S., Tapping, R.I., 2010. Toll-like receptor signaling in cell proliferation and survival. *Cytokine*, 49, 1–9.
- Liew, F.Y., Xu, D., Brint, E.K., O'Neill, L.A.J., 2005. Negative regulation of Toll-like receptor-mediated immune responses. *Nat Rev Immunol*, 5, 446–458.
- Lim, K.H., Staudt, L.M., 2013. Toll-Like Receptor Signaling. *Cold Spring Harb Perspect Biol*, 5, a011247.
- Lin, C.H., Cheng, H.W., Ma, H.P., Wu, C.H., Hong, C.Y., Chen, B.C., 2011. Thrombin induces NF- $\kappa$ B activation and IL-8/CXCL8 expression in lung epithelial cells by a Rac1-dependent PI3K/Akt pathway. *J Biol Chem*, 286, 10483–10494.
- Lin, S.C., Lo, Y.C., Wu, H., 2010. Helical assembly in the MyD88–IRAK4–IRAK2 complex in TLR/IL-1R signalling. *Nature*, 465, 885–890.
- Liou T.G., Adler F.R., Fitzsimmons S.C., Cahill B.C., Hibbs J.R., Marshall B.C., 2001. Predictive 5-year survivorship model of cystic fibrosis. *Am J Epidemiol*, 153, 345–52.

- LiPuma, J.J., 2010. The changing microbial epidemiology in cystic fibrosis. *Clin Microbiol Rev*, 23, 299–323.
- LiPuma, J.J., Mortensen, J.E., Dasen, S.E., Edlind, T.D., Schidlow, D.V., Burns, J.L., Stull, T.L., 1988. Ribotype analysis of *Pseudomonas cepacia* from cystic fibrosis treatment centers. *J Pediatr*, 113, 859–862.
- LiPuma, J.J., Spilker, T., Coenye, T., Gonzalez, C.F., 2002. An epidemic *Burkholderia cepacia* complex strain identified in soil. *Lancet*, 359, 2002–2003.
- Liu, Y.-S., Beyer, A., Aebersold, R., 2016. On the dependency of cellular protein levels on mRNA abundance. *Cell*, 165, 535–550.
- Liu, Yilei, Lardi, M., Pedrioli, A., Eberl, L., Pessi, G., 2017. NtrC-dependent control of exopolysaccharide synthesis and motility in *Burkholderia cenocepacia* H111. *PLoS ONE*, 12, e0180362.
- Loutet, S.A., Flannagan, R.S., Kooi, C., Sokol, P.A., Valvano, M.A., 2006. A complete lipopolysaccharide inner core oligosaccharide is required for resistance of *Burkholderia cenocepacia* to antimicrobial peptides and bacterial survival *in vivo*. *J Bacterio*, 188, 2073–2080.
- Loutet, S.A., Mussen, L.E., Flannagan, R.S., Valvano, M.A., 2010. A two-tier model of polymyxin B resistance in *Burkholderia cenocepacia*. *Environ Microbiol Rep*, 3, 278–285.
- Love, W., Dobbs, N., Tabor, L., Simecka, J.W., 2010. Toll-like receptor 2 plays a major role in innate resistance in the lung against murine *Mycoplasma*. *PLoS ONE*, 5, e10739.
- Luo, H.R., Mondal, S., 2015. Molecular control of PtdIns(3,4,5)P3 signaling in neutrophils. *EMBO rep*, 16, 149–163.
- Luong, T.T., Lee, C.Y., 2002. Overproduction of type 8 capsular polysaccharide augments *Staphylococcus aureus* virulence. *Infect Immun*, 70, 3389–3395.
- Luyendyk, J.P., Schabbauer, G.A., Tencati, M., Holscher, T., Pawlinski, R., Mackman, N., 2008. Genetic analysis of the role of the PI3K-Akt pathway in lipopolysaccharide-induced cytokine and tissue factor gene expression in monocytes/macrophages. *J Immunol*, 180, 4218–4226.
- Lynch, S.V., Bruce, K.D., 2013. The cystic fibrosis airway microbiome. *Cold Spring Harb Perspect Med*, 3, a009738.
- Ma, J., Rubin, B.K., Voynow, J.A., 2018. Mucins, mucus, and goblet cells. *Chest*, 154, 169–176.
- Malik, Z., Roscioli, E., Murphy, J., Ou, J., Bassiouni, A., Wormald, P.-J., Vreugde, S., 2015. *Staphylococcus aureus* impairs the airway epithelial barrier in vitro. *Int Forum Allergy Rhinol*, 5, 551–556.
- Malott, R.J., Steen-Kinnaird, B.R., Lee, T.D., Speert, D.P., 2012. Identification of hopanoid biosynthesis genes involved in polymyxin resistance in *Burkholderia multivorans*. *Antimicrob Agents Chemother*, 56, 464–471.
- Mandell, G.L., 1975. Catalase, superoxide dismutase, and virulence of *Staphylococcus aureus*. *In vitro* and *in vivo* studies with emphasis on staphylococcal-leukocyte interaction. *J Clin Invest*, 55, 561–566.
- Manicassamy, S., Pulendran, B., 2009. Modulation of adaptive immunity with Toll-like receptors. *Semin Immunol*, 21, 185–193.
- Mann, E.E., Wozniak, D.J., 2012. *Pseudomonas* biofilm matrix composition and niche biology. *FEMS Microbiol Rev*, 36, 893–916.

- Mansell, A., Smith, R., Doyle, S.L., Gray, P., Fenner, J.E., Crack, P.J., Nicholson, S.E., Hilton, D.J., O'Neill, L.A.J., Hertzog, P.J., 2006. Suppressor of cytokine signaling 1 negatively regulates Toll-like receptor signaling by mediating Mal degradation. *Nat Immunol*, 7, 148–155.
- Marsh, R.L., Thornton, R.B., Smith-Vaughan, H.C., Richmond, P., Pizzutto, S.J., Chang, A.B., 2014. Detection of biofilm in bronchoalveolar lavage from children with non-cystic fibrosis bronchiectasis. *Pediatr Pulmonol*, 50, 284–29.
- Martin, C., Burdon, P.C.E., Bridger, G., Gutierrez-Ramos, J.C., Williams, T.J., Rankin, S.M., 2003. Chemokines acting via CXCR2 and CXCR4 control the release of neutrophils from the bone marrow and their return following senescence. *Immunity*, 19, 583–593.
- Martin, K.J.S., Muessel, M.J., Pullar, C.E., Willars, G.B., Wardlaw, A.J., 2015. The role of phosphoinositide 3-kinases in neutrophil migration in 3D collagen gels. *PLoS ONE*, 10, e0116250.
- Massengale, A.R., Quinn, F.J., Yankaskas, J., Weissman, D., McClellan, W.T., Cuff, C., Aronoff, S.C., 1999. Reduced interleukin-8 production by cystic fibrosis airway epithelial cells. *Am J Respir Cell Mol Biol*, 20, 1073–1080.
- Mathew, A., Eberl, L., Carlier, A.L., 2014. A novel siderophore-independent strategy of iron uptake in the genus *Burkholderia*. *Mol Microbiol*, 91, 805–820.
- Matsumiya, T., Stafforini, D.M., 2010. Function and regulation of retinoic acid-inducible gene-I. *Crit Rev Immunol*, 30, 489–513.
- Mayer, A.K., Muehmer, M., Mages, J., Gueinzus, K., Hess, C., Heeg, K., Bals, R., Lang, R., Dalpke, A.H., 2007. Differential recognition of TLR-dependent microbial ligands in human bronchial epithelial cells. *J Immunol*, 178, 3134–3142.
- Mayer-Barber, K.D., Andrade, B.B., Barber, D.L., Hieny, S., Feng, C.G., Caspar, P., Oland, S., Gordon, S., Sher, A., 2011. Innate and adaptive interferons suppress IL-1 $\alpha$  and IL-1 $\beta$  production by distinct pulmonary myeloid subsets during *Mycobacterium tuberculosis* infection. *Immunity*, 35, 1023–1034.
- Mazer, D.M., Young, C., Kalikin, L.M., Spilker, T., LiPuma, J.J., 2017. *In vitro* activity of ceftolozane-tazobactam and other antimicrobial agents against *Burkholderia cepacia* complex and *Burkholderia gladioli*. *Antimicrob Agents Chemother*, 61, e00766–17.
- McArdle, A.J., Turkova, A., Cunningham, A.J., 2018. When do co-infections matter? *Curr Opin Infect Dis*, 31, 209–215.
- McClean, S., Callaghan, M., 2009. *Burkholderia cepacia* complex: epithelial cell-pathogen confrontations and potential for therapeutic intervention. *J Med Microbiol*, 58, 1–12.
- McClure, R., Massari, P., 2014. TLR-dependent human mucosal epithelial cell responses to microbial pathogens. *Front Immunol*, 5, 386.
- McGuire, V.A., Arthur, J.S.C., 2015. Subverting Toll-like receptor signaling by bacterial pathogens. *Front Immunol*, 6, 373.
- McLaws, F.B., Larsen, A.R., Skov, R.L., Chopra, I., O'Neill, A.J., 2011. Distribution of fusidic acid resistance determinants in methicillin-resistant *Staphylococcus aureus*. *Antimicrob Agents Chemother*, 55, 1173–1176.
- McMillan, S.J., Sharma, R.S., McKenzie, E.J., Richards, H.E., Zhang, J., Prescott, A., Crocker, P.R., 2013. Siglec-E is a negative regulator of acute pulmonary neutrophil inflammation and suppresses CD11b  $\beta$ 2-integrin-dependent signaling. *Blood*, 121, 2084–2094.

- Medzhitov, R., 2001. Toll-like receptors and innate immunity. *Nat Rev Immunol*, 1, 135–145.
- Mittal, S.K., Cho, K.-J., Ishido, S., Roche, P.A., 2015. Interleukin 10-mediated Immunosuppression: March-I induction regulates antigen presentation by macrophages but not dendritic cells. *J Biol Chem*, 290, 27158–27167.
- Mohammad Hosseini, A., Majidi, J., Baradaran, B., Yousefi, M., 2015. Toll-like receptors in the pathogenesis of autoimmune diseases. *Adv Pharm Bull*, 5, 605–614.
- Monick, M.M., Yarovinsky, T.O., Powers, L.S., Butler, N.S., Carter, A.B., Gudmundsson, G., Hunninghake, G.W., 2003. Respiratory syncytial virus up-regulates TLR4 and sensitizes airway epithelial cells to endotoxin. *J Biol Chem*, 278, 53035–53044.
- Montgomery, C.P., Boyle-Vavra, S., Adem, P.V., Lee, J.C., Husain, A.N., Clasen, J., Daum, R.S., 2008. Comparison of virulence in community-associated methicillin-resistant *Staphylococcus aureus* pulsotypes USA300 and USA400 in a rat model of pneumonia. *J Infect Dis*, 198:561–570.
- Montgomery, C.P., Daniels, M., Zhao, F., Alegre, M.-L., Chong, A.S., Daum, R.S., 2014. Protective immunity against recurrent *Staphylococcus aureus* skin infection requires antibody and interleukin-17A. *Infect Immun*, 82, 2125–2134.
- Moreilhon, C., Gras, D., Hologne, C., Bajolet, O., Cottrez, F., Magnone, V., Merten, M., Groux, H., Puchelle, E., Barbry, P., 2005. Live *Staphylococcus aureus* and bacterial soluble factors induce different transcriptional responses in human airway cells. *Physiol Genomics*, 20, 244–255.
- Morimoto, K., Janssen, W.J., Terada, M., 2012. Defective efferocytosis by alveolar macrophages in IPF patients. *Respir Med*, 106, 1800–1803.
- Morris, D.G., Huang, X., Kaminski, N., Wang, Y., Shapiro, S.D., Dolganov, G., Glick, A., Sheppard, D., 2003. Loss of integrin  $\alpha\beta6$ -mediated TGF- $\beta$  activation causes Mmp12-dependent emphysema. *Nature*, 422, 169–173.
- Muir, A., Soong, G., Sokol, S., Reddy, B., Gomez, M.I., van Heeckeren, A., Prince, A., 2004. Toll-like receptors in normal and cystic fibrosis airway epithelial cells. *Am J Respir Cell Mol Biol*, 30, 777–783.
- Muller-Anstett, M.A., Muller, P., Albrecht, T., Nega, M., Wagener, J., Gao, Q., Kaesler, S., Schaller, M., Biedermann, T., Götz, F., 2010. Staphylococcal peptidoglycan co-localizes with Nod2 and TLR2 and activates innate immune response via both receptors in primary murine keratinocytes. *PLoS ONE*, 5, e13153.
- Murray J L, Connell J L, Stacy A, et al. Mechanisms of synergy in polymicrobial infections. *J Microbiol*, 2014, 52(3): 188-199.
- Nagl, M., Kacani, L., Müllauer, B., Lemberger, E.-M., Stoiber, H., Sprinzl, G.M., Schennach, H., Dierich, M.P., 2002. Phagocytosis and killing of bacteria by professional phagocytes and dendritic cells. *Clin Diagn Lab Immunol*, 9, 1165–1168.
- Nair, D., Memmi, G., Hernandez, D., Bard, J., Beaume, M., Gill, S., Francois, P., Cheung, A.L., 2011. Whole-genome sequencing of *Staphylococcus aureus* strain RN4220, a key laboratory strain used in virulence research, identifies mutations that affect not only virulence factors but also the fitness of the strain. *J Bacterio*, 193, 2332–2335.
- Nakayama K, Okugawa S, Yanagimoto S, et al. Involvement of IRAK-M in peptidoglycan-induced tolerance in macrophages. *J of Biol Chem*, 2004, 279(8): 6629-6634.
- Nanra, J.S., Buitrago, S.M., Crawford, S., Ng, J., Fink, P.S., Hawkins, J., Scully, I.L., McNeil, L.K., Aste-Amézaga, J.M., Cooper, D., Jansen, K.U., Anderson, A.S., 2013. Capsular polysaccharides are an important immune evasion mechanism for *Staphylococcus aureus*. *Hum Vaccin Immunother*, 9, 480–487.

- Nash, J.A., Ballard, T.N.S., Weaver, T.E., Akinbi, H.T., 2006. The peptidoglycan-degrading property of lysozyme is not required for bactericidal activity *in vivo*. *J Immunol*, 177, 519–526.
- Nguyen, A.T., Oglesby-Sherrouse, A.G., 2016. Interactions between *Pseudomonas aeruginosa* and *Staphylococcus aureus* during co-cultivations and polymicrobial infections. *Appl Microbiol Biotechnol*, 100, 6141–6148.
- Ni, M., MacFarlane, A.W., Toft, M., Lowell, C.A., Campbell, K.S., Hamerman, J.A., 2012. B-cell adaptor for PI3K (BCAP) negatively regulates Toll-like receptor signaling through activation of PI3K. *Proc Natl Acad Sci*, 109, 267–272.
- Nicholas, D.A., Zhang, K., Hung, C., Glasgow, S., Aruni AW, Unternaehrer J., Payne, K.J., Langridge, W.H.R., De Leon, M., 2017. Palmitic acid is a toll-like receptor 4 ligand that induces human dendritic cell secretion of IL-1 $\beta$ . *PLoS ONE*, 12, p.e0176793.
- Nixon, G.M., Armstrong, D.S., Carzino, R., Carlin, J.B., Olinsky, A., Robertson, C.F., Grimwood, K., 2001. Clinical outcome after early *Pseudomonas aeruginosa* infection in cystic fibrosis. *J Pediatr*, 138, 699–704.
- Nold-Petry, C.A., Lo, C.Y., Rudloff, I., Elgass, K.D., Li, S., Gantier, M.P., Lotz-Havla, A.S., Gersting, S.W., Cho, S.X., Lao, J.C., Ellisdon, A.M., Rotter, B., Azam, T., Mangan, N.E., Rossello, F.J., Whisstock, J.C., Bufler, P., Garlanda, C., Mantovani, A., Dinarello, C.A., Nold, M.F., 2015. IL-37 requires the receptors IL-18R $\alpha$  and IL-1R8 (SIGIRR) to carry out its multifaceted anti-inflammatory program upon innate signal transduction. *Nat Immunol*, 16, 354–365.
- Nomura, H., Sato, E., Koyama, S., Haniuda, M., Kubo, K., Nagai, S., Izumi, T., 2001. Histamine stimulates alveolar macrophages to release neutrophil and monocyte chemotactic activity. *J Lab Clin Med*, 138, 226–235.
- O'Gorman, W.E., Hsieh, E.W.Y., Savig, E.S., Gherardini, P.F., Hernandez, J.D., Hansmann, L., Balboni, I.M., Utz, P.J., Bendall, S.C., Fantl, W.J., Lewis, D.B., Nolan, G.P., Davis, M.M., 2015. Single-cell systems-level analysis of human Toll-like receptor activation defines a chemokine signature in patients with systemic lupus erythematosus. *J Allergy Clin Immunol*, 136, 1326–1336.
- O'Halloran, D.P., Wynne, K., Geoghegan, J.A., 2015. Protein A is released into the *Staphylococcus aureus* culture supernatant with an unprocessed sorting signal. *Infect Immun*, 83, 1598–1609.
- Oever, ten, J., Kox, M., van de Veerdonk, F.L., Mothapo, K.M., Slavcovic, A., Jansen, T.L., Tweehuysen, L., Giamarellos-Bourboulis, E.J., Schneeberger, P.M., Wever, P.C., Stoffels, M., Simon, A., van der Meer, J.W., Johnson, M.D., Kullberg, B.-J., Pickkers, P., Pachot, A., Joosten, Leo AB, Netea, M.G., 2014. The discriminative capacity of soluble Toll-like receptor (sTLR)2 and sTLR4 in inflammatory diseases. *BMC Immunol*, 15, 55.
- Okshevsky, M., Meyer, R.L., 2015. The role of extracellular DNA in the establishment, maintenance and perpetuation of bacterial biofilms. *Crit Rev Microbiol*, 41, 341–352.
- Orazi, G., O'Toole, G.A., 2017. *Pseudomonas aeruginosa* alters *Staphylococcus aureus* sensitivity to vancomycin in a biofilm model of cystic fibrosis infection. *MBio*, 8, e00873-17.
- Oviedo-Boyso, J., Cortes-Vieyra, R., Huante-Mendoza, A., Yu, H.B., Valdez-Alarcon, J.J., Bravo-Patino, A., Cajero-Juarez, M., Finlay, B.B., Baizabal-Aguirre, V.M., 2011. The Phosphoinositide-3-kinase-Akt signaling pathway is important for *Staphylococcus aureus* internalization by endothelial cells. *Infect Immun*, 79, 4569–4577.
- O'Neill, A.J., 2010. *Staphylococcus aureus* SH1000 and 8325-4: comparative genome sequences of key laboratory strains in Staphylococcal research. *Lett Appl Microbiol*, 51, 358–361.



- Pace, E., Ferraro, M., Siena, L., Melis, M., Montalbano, A.M., Johnson, M., Bonsignore, M.R., Bonsignore, G., Gjomarkaj, M., 2008. Cigarette smoke increases Toll-like receptor 4 and modifies lipopolysaccharide-mediated responses in airway epithelial cells. *Immunology*, 124, 401–411.
- Palfreyman, R.W., Watson, M.L., Eden, C., Smith, A.W., 1997. Induction of biologically active interleukin-8 from lung epithelial cells by *Burkholderia (Pseudomonas) cepacia* Products 65, 617–622. *Infect Immun*, 65, 617–622.
- Palmqvist, N., Patti, J.M., Tarkowski, A., Josefsson, E., 2004. Expression of Staphylococcal clumping factor A impedes macrophage phagocytosis. *Microbes Infect*, 6, 188–195.
- Parker, D., Prince, A., 2013. Epithelial uptake of flagella initiates proinflammatory signaling. *PLoS ONE*, 8, e59932.
- Patel, D., Wines, B.D., Langley, R.J., Fraser, J.D., 2010. Specificity of Staphylococcal superantigen-like protein 10 toward the human IgG1 Fc domain. *J Immunol*, 184, 6283–6292.
- Patwa, A., Shah, A., 2015. Anatomy and physiology of respiratory system relevant to anaesthesia. *Indian J Anaesth*, 59, 533–541.
- Pauli, N.T., Kim, H.K., Falugi, F., Huang, M., Dulac, J., Henry Dunand, C., Zheng, N.-Y., Kaur, K., Andrews, S.F., Huang, Y., DeDent, A., Frank, K.M., Charnot-Katsikas, A., Schneewind, O., Wilson, P.C., 2014. *Staphylococcus aureus* infection induces protein A-mediated immune evasion in humans. *J Exp Med*, 211, 2331–2339.
- Peddayelachagiri, B.V., Paul, S., Nagaraj, S., Gogoi, M., Sripathy, M.H., Batra, H.V., 2016. Prevalence and identification of *Burkholderia pseudomallei* and near-neighbor species in the malabar coastal region of India. *PLOS Negl Trop Dis*, 10, e0004956.
- Peres, A.G., Stegen, C., Li, J., Xu, A.Q., Levast, B., Surette, M.G., Cousineau, B., Desrosiers, M., Madrenas, J., 2015. Uncoupling of pro- and anti-inflammatory properties of *Staphylococcus aureus*. *Infect Immun*, 83, 1587–1597.
- Perez, A.C., Pang, B., King, L.B., Tan, L., Murrah, K.A., Reimche, J.L., Wren, J.T., Richardson, S.H., Ghandi, U., Swords, W.E., 2014. Residence of *Streptococcus pneumoniae* and *Moraxella catarrhalis* within polymicrobial biofilm promotes antibiotic resistance and bacterial persistence *in vivo*. *Pathog Dis*, 70, 280–288.
- Peschel, A., Jack, R.W., Otto, M., Collins, L.V., Staubitz, P., Nicholson, G., Kalbacher, H., Nieuwenhuizen, W.F., Jung, G., Tarkowski, A., 2001. *Staphylococcus aureus* resistance to human defensins and evasion of neutrophil killing via the novel virulence factor MprF is based on modification of membrane lipids with l-lysine. *J Exp Med*, 193, 1067–1076.
- Peschel, A., Otto, M., Jack, R.W., Kalbacher, H., Jung, G., Götz, F., 1999. Inactivation of the *dlt* operon in *Staphylococcus aureus* confers sensitivity to defensins, protegrins, and other antimicrobial peptides. *J Biol Chem*, 274, 8405–8410.
- Peterson, B.W., Sharma, P.K., van der Mei, H.C., Busscher, H.J., 2011. Bacterial cell surface damage due to centrifugal compaction. *Appl Environ Microbiol*, 78, 120–125.
- Pica, F., Franzese, O., D'Onofrio, C., Bonmassar, E., Favalli, C., Garaci, E., 1996. Prostaglandin E2 induces apoptosis in resting immature and mature human lymphocytes: a c-Myc-dependent and Bcl-2-independent associated pathway. *J Pharmacol Exp Ther*, 277, 1793–1800.
- Pillay, C.S., Elliott, E., Dennison, C., 2002. Endolysosomal proteolysis and its regulation. *Biochem J*, 363, 417–429.
- Podnecky, N., Rhodes, K., Schweizer, H., 2015. efflux pump-mediated drug resistance in *Burkholderia*. *Front Microbio*, 6, 305.

- Pragman, A.A., Berger, J.P., Williams, B.J., 2016. Understanding persistent bacterial lung infections: clinical implications informed by the biology of the microbiota and biofilms. *Clin Pulm Med*, 23, 57–66.
- Prat, C., Bestebroer, J., de Haas, C.J.C., van Strijp, J.A.G., van Kessel, K.P.M., 2006. A new Staphylococcal anti-inflammatory protein that antagonizes the formyl peptide receptor-like 1. *J Immunol*, 177, 8017–8026.
- Prat, C., Haas, P.-J., Bestebroer, J., de Haas, C.J.C., van Strijp, J.A.G., van Kessel, K.P.M., 2009. A homolog of formyl peptide receptor-like 1 (FPRL1) inhibitor from *Staphylococcus aureus* (FPRL1 inhibitory protein) that inhibits FPRL1 and FPR. *J Immunol*, 183, 6569–6578.
- Proctor, R.A., Eiff, von, C., Kahl, B.C., Becker, K., McNamara, P., Herrmann, M., Peters, G., 2006. Small colony variants: a pathogenic form of bacteria that facilitates persistent and recurrent infections. *Nat Rev Micro*, 4, 295–305.
- Ragupathy, S., Esmaili, F., Paschoud, S., Sublet, E., Citi, S., Borchard, G., 2014. Toll-like receptor 2 regulates the barrier function of human bronchial epithelial monolayers through atypical protein kinase C  $\zeta$ , and an increase in expression of claudin-1. *Tissue Barriers*, 2, e29166–8.
- Ramos, F.L., Krahne, J.S., Kim, V., 2014. Clinical issues of mucus accumulation in COPD. *Int J Chron Obstruct Pulmon Dis*, 9, 139–150.
- Ramsay, K.A., Butler, C.A., Paynter, S., Ware, R.S., Kidd, T.J., Wainwright, C.E., Bell, S.C., 2013. Factors influencing acquisition of *Burkholderia cepacia* complex organisms in patients with cystic fibrosis. *J Clin Microbiol*, 51, 3975–3980.
- Ramsey, M.M., Whiteley, M., 2009. Polymicrobial interactions stimulate resistance to host innate immunity through metabolite perception. *Proc Natl Acad Sci*, 106, 1578–1583.
- Resch, A., Leicht, S., Saric, M., Pasztor, L., Jakob, A., Götz, F., Nordheim, A., 2006. Comparative proteome analysis of *Staphylococcus aureus* biofilm and planktonic cells and correlation with transcriptome profiling. *Proteomics*, 6, 1867–1877.
- Reyes-Robles, T., Alonzo, F.3., Kozhaya, L., Lacy, D.B., Unutmaz, D., Torres, V.J., 2013. *Staphylococcus aureus* leukotoxin ED targets the chemokine receptors CXCR1 and CXCR2 to kill leukocytes and promote infection. *Cell Host Microbe*, 14, 453–459.
- Reynolds, J.M., Pappu, B.P., Peng, J., Martinez, G.J., Zhang, Y., Chung, Y., Ma, L., Yang, X.O., Nurieva, R.I., Tian, Q., Dong, C., 2010. Toll-like receptor 2 signaling in CD4<sup>+</sup> T lymphocytes promotes T helper 17 responses and regulates the pathogenesis of autoimmune disease. *Immunity*, 32, 692–702.
- Rhodes, K.A., Schweizer, H.P., 2016. Antibiotic resistance in *Burkholderia* species. *Drug Resist Updat*, 28, 82–90.
- Rice A.B., Ingram J.L., Bonner J.C., 2002. p38 mitogen-activated protein kinase regulates growth factor-induced mitogenesis of rat pulmonary myofibroblasts. *Am J Respir Cell Mol Biol*, 27, 759–765.
- Richardson, A.R., Libby, S.J., Fang, F.C., 2008. A nitric oxide-inducible lactate dehydrogenase enables *Staphylococcus aureus* to resist innate immunity. *Science*, 319, 1672–1676.
- Ricklin, D., Ricklin-Lichtsteiner, S.K., Markiewski, M.M., Geisbrecht, B.V., Lambris, J.D., 2008. Cutting Edge: Members of the *Staphylococcus aureus* extracellular fibrinogen-binding protein family inhibit the interaction of C3d with complement receptor 2. *J Immunol*, 181, 7463–7467.
- Rivas-Santiago, B., Hernandez-Pando, R., Carranza, C., Juarez, E., Contreras, J.L., Aguilar-Leon, D., Torres, M., Sada, E., 2008. Expression of cathelicidin LL-37 during

- Mycobacterium tuberculosis* infection in human alveolar macrophages, monocytes, neutrophils, and epithelial cells. *Infect Immun*, 76, 935–941.
- Robinson, D.A., Monk, A.B., Cooper, J.E., Feil, E.J., Enright, M.C., 2005. Evolutionary genetics of the accessory gene regulator (*agr*) locus in *Staphylococcus aureus*. *J Bacterio*, 187, 8312–8321.
- Roche, F.M., Downer, R., Keane, F., Speziale, P., Park, P.W., Foster, T.J., 2004. The N-terminal a domain of fibronectin-binding proteins A and B promotes adhesion of *Staphylococcus aureus* to elastin. *J Biol Chem*, 279, 38433–38440.
- Rooijackers, S.H.M., Ruyken, M., Roos, A., Daha, M.R., Presanis, J.S., Sim, R.B., van Wamel, W.J.B., van Kessel, K.P.M., van Strijp, J.A.G., 2005. Immune evasion by a *Staphylococcal* complement inhibitor that acts on C3 convertases. *Nat Immunol*, 6, 920–927.
- Rosadini, C.V., Kagan, J.C., 2015. Microbial strategies for antagonizing Toll-like-receptor signal transduction. *Innate Immun*, 32, 61–70.
- Rosadini, C.V., Kagan, J.C., 2017. Early innate immune responses to bacterial LPS. *Innate Immun*, 44, 14–19.
- Rosadini, C.V., Zanoni, I., Odendall, C., Green, E.R., Paczosa, M.K., Philip, N.H., Brodsky, I.E., Meccas, J., Kagan, J.C., 2015. A single bacterial immune evasion strategy dismantles both MyD88 and TRIF signalling pathways downstream of TLR4. *Cell Host Microbe*, 18, 682–693.
- Rosales-Reyes, R., Aubert, D.F., Tolman, J.S., Amer, A.O., Valvano, M.A., 2012a. *Burkholderia cenocepacia* type VI secretion system mediates escape of type II secreted proteins into the cytoplasm of infected macrophages. *PLoS ONE*, 7, e41726.
- Rosales-Reyes, R., Skeldon, A.M., Aubert, D.F., Valvano, M.A., 2012b. The Type VI secretion system of *Burkholderia cenocepacia* affects multiple Rho family GTPases disrupting the actin cytoskeleton and the assembly of NADPH oxidase complex in macrophages. *Cell Microbiol*, 14, 255–273.
- Rosenberger, C.M., Gallo, R.L., Finlay, B.B., 2004. Interplay between antibacterial effectors: a macrophage antimicrobial peptide impairs intracellular *Salmonella* replication. *Proc Natl Acad Sci*, 101, 2422–2427.
- Rothlin, C.V., Ghosh, S., Zuniga, E.I., Oldstone, M.B.A., Lemke, G., 2007. TAM receptors are pleiotropic inhibitors of the innate immune response. *Cell*, 131, 1124–1136.
- Rubins, J.B., 2003. Alveolar macrophages: wielding the double-edged sword of inflammation. *Am J Respir Crit Care Med*, 167, 103–104.
- Ruggeri, P., Nicocia, G., Venza, I., Venza, M., Valenti, A., Teti, D., 2000. Polyamine metabolism in prostaglandin E2-treated human T lymphocytes. *Immunopharmacol Immunotoxicol*, 22, 117–129.
- Rutz, M., Metzger, J., Gellert, T., Lippa, P., Lipford, G.B., Wagner, H., Bauer, S., 2004. Toll-like receptor 9 binds single-stranded CpG-DNA in a sequence- and pH-dependent manner. *Eur J Immunol*, 34, 2541–2550.
- Ryu, J.-H., Yoo, J.-Y., Kim, M.-J., Hwang, S.-G., Ahn, K.C., Ryu, J.-C., Choi, M.-K., Joo, J.H., Kim, C.-H., Lee, S.-N., Lee, W.-J., Kim, J., Shin, D.M., Kweon, M.-N., Bae, Y.S., Yoon, J.-H., 2013. Distinct TLR-mediated pathways regulate house dust mite-induced allergic disease in the upper and lower airways. *J Allergy Clin Immunol*, 131, 549–561.
- Sajjan, S.U., Carmody, L.A., Gonzalez, C.F., LiPuma, J.J., 2008. A type IV secretion system contributes to intracellular survival and replication of *Burkholderia cenocepacia*. *Infect Immun*, 76, 5447–5455.

- Sajjan, U.S., Hershenson, M.B., Forstner, J.F., LiPuma, J.J., 2008. *Burkholderia cenocepacia* ET12 strain activates TNFR1 signalling in cystic fibrosis airway epithelial cells. *Cell Microbiol*, 10, 188–201.
- Saldias, M.S., Lamothe, J., Wu, R., Valvano, M.A., 2008. *Burkholderia cenocepacia* requires the RpoN  $\sigma$  factor for biofilm formation and intracellular trafficking within macrophages. *Infect Immun*, 76, 1059–1067.
- Saldias, M.S., Valvano, M.A., 2009. Interactions of *Burkholderia cenocepacia* and other *Burkholderia cepacia* complex bacteria with epithelial and phagocytic cells. *Microbiology*, 155, 2809–2817.
- Saldías, M.S., Ortega, X., Valvano, M.A., 2009. *Burkholderia cenocepacia* O antigen lipopolysaccharide prevents phagocytosis by macrophages and adhesion to epithelial cells. *J Med Microbiol*, 58, 1542–1548.
- Salta, E., Kanata, E., Ouzounis, C.A., Gilch, S., Schätzl, H., Sklaviadis, T., 2014. Assessing proteinase K resistance of fish prion proteins in a scrapie-infected mouse neuroblastoma cell line. *Viruses*, 6, 4398–4421.
- Sang, H.R., Kim, H., Moyer, M.P., Pothoulakis, C., 2006. Role of MyD88 in phosphatidylinositol 3-kinase activation by flagellin/Toll-like receptor 5 engagement in colonic epithelial cells. *J Biol Chem*, 281, 18560–18568.
- Sass, L.A., Hair, P.S., Perkins, A.M., Shah, T.A., Krishna, N.K., Cunnion, K.M., 2015. Complement effectors of inflammation in cystic fibrosis lung fluid correlate with clinical measures of disease. *PLoS ONE*, 10, e0144723.
- Sasso, E.H., Silverman, G.J., Mannik, M., 1989. Human IgM molecules that bind Staphylococcal protein A contain VHIII H chains. *J Immunol*, 142, 2778–2783.
- Savage, V.J., Chopra, I., O'Neill, A.J., 2013. Staphylococcus aureus biofilms promote horizontal transfer of antibiotic resistance. *Antimicrob Agents Chemother*, 57, 1968–1970.
- Schilcher, K., Andreoni, F., Dengler Haunreiter, V., Seidl, K., Hasse, B., Zinkernagel, A.S., 2016. Modulation of *Staphylococcus aureus* biofilm matrix by subinhibitory concentrations of clindamycin. *Antimicrob Agents Chemother*, 60, 5957–5967.
- Schmitz, F.J., 2001. Resistance to tetracycline and distribution of tetracycline resistance genes in European *Staphylococcus aureus* isolates. *J Antimicrob Chemother*, 47, 239–240.
- Schommer, N.N., Christner, M., Hentschke, M., Ruckdeschel, K., Aepfelbacher, M., Rohde, H., 2011. *Staphylococcus epidermidis* uses distinct mechanisms of biofilm formation to interfere with phagocytosis and activation of mouse macrophage-like cells 774A.1. *Infect Immun*, 79, 2267–2276.
- Schröder, N.W., Pfeil, D., Opitz, B., Michelsen, K.S., Amberger, J., Zähringer, U., Göbel, U.B., Schumann, R.R., 2001. Activation of mitogen-activated protein kinases p42/44, p38, and stress-activated protein kinases in myelo-monocytic cells by *Treponema lipoteichoic acid*. *J Biol Chem*, 276, 9713–9719.
- Schwab, U., Abdullah, L.H., Perlmutter, O.S., Albert, D., Davis, C.W., Arnold, R.R., Yankaskas, J.R., Gilligan, P., Neubauer, H., Randell, S.H., Boucher, R.C., 2014. Localization of *Burkholderia cepacia* complex bacteria in cystic fibrosis lungs and interactions with *Pseudomonas aeruginosa* in hypoxic mucus. *Infect Immun*, 82, 4729–4745.
- Schwab, U., Leigh, M., Ribeiro, C., Yankaskas, J., Burns, K., Gilligan, P., Sokol, P., Boucher, R., 2002. Patterns of epithelial cell invasion by different species of the

- Burkholderia cepacia* complex in well-differentiated human airway epithelia. *Infect Immun*, 70, 4547–4555.
- Schwandner, R., Dziarski, R., Wesche, H., Rothe, M., Kirschning, C.J., 1999. Peptidoglycan- and lipoteichoic acid-induced cell activation is mediated by Toll-like receptor 2. *J Biol Chem*, 274, 17406–17409.
- Schweighoffer, E., Nys, J., Vanes, L., Smithers, N., Tybulewicz, V.L.J., 2017. TLR4 signals in B lymphocytes are transduced via the B cell antigen receptor and SYK. *J Exp Med*, 214, 1269–1280.
- Schweppe, D.K., Harding, C., Chavez, J.D., Wu, X., Ramage, E., Singh, P.K., Manoil, C., Bruce, J.E., 2015. Host-microbe protein interactions during bacterial infection. *Chem Biol*, 22, 1521–1530.
- Scoffone, V.C., Chiarelli, L.R., Makarov, V., Brackman, G., Israyilova, A., Azzalin, A., Forneris, F., Riabova, O., Savina, S., Coenye, T., Riccardi, G., Buroni, S., 2016. Discovery of new diketopiperazines inhibiting *Burkholderia cenocepacia* quorum sensing *in vitro* and *in vivo*. *Sci Rep*, 6, 964.
- Scoffone, V.C., Chiarelli, L.R., Trespidi, G., Mentasti, M., Riccardi, G., Buroni, S., 2017. *Burkholderia cenocepacia* infections in cystic fibrosis patients: drug resistance and therapeutic approaches. *Front Microbio*, 8, 1592.
- Scott, M.G., Yan, H., Hancock, R.E., 1999. Biological properties of structurally related  $\alpha$ -helical cationic antimicrobial peptides. *Infect Immun*, 67, 2005–2009.
- Sears, P.R., Yin, W.-N., Ostrowski, L.E., 2015. Continuous mucociliary transport by primary human airway epithelial cells *in vitro*. *Am J Physiol Lung Cell Mol Physiol*, 309, L99–108.
- Secor, P.R., James, G.A., Fleckman, P., Olerud, J.E., McInnerney, K., Stewart, P.S., 2011. *Staphylococcus aureus* Biofilm and Planktonic cultures differentially impact gene expression, MAPK phosphorylation, and cytokine production in human keratinocytes. *BMC Microbiol*, 11, 143.
- Seiler, F., Hellberg, J., Lepper, P.M., Kamyschnikow, A., Herr, C., Bischoff, M., Langer, F., Schafers, H.J., Lammert, F., Menger, M.D., Bals, R., Beisswenger, C., 2013. FOXO transcription factors regulate innate immune mechanisms in respiratory epithelial cells. *J Immunol*, 190, 1603–1613.
- Selsted, M.E., Ouellette, A.J., 2005. Mammalian defensins in the antimicrobial immune response. *Nat Immunol*, 6, 551–557.
- Shembade, N., Ma, A., Harhaj, E.W., 2010. Inhibition of NF- $\kappa$ B signaling by A20 through disruption of ubiquitin enzyme complexes. *Science*, 327, 1135–1139.
- Shen, B.Q., Finkbeiner, W.E., Wine, J.J., Mrnsy, R.J., Widdicombe, J.H., 1994. Calu-3: a human airway epithelial cell line that shows cAMP-dependent Cl<sup>-</sup> secretion. *Am J Physiol*, 266, L493–L501.
- Sheppard, D., 2006. Transforming growth factor  $\beta$ : a central modulator of pulmonary and airway inflammation and fibrosis. *Proc Am Thorac Soc*, 3, 413–417.
- Shikhagaie, M.M., Andersson, C.K., Mori, M., Kortekaas Krohn, I., Bergqvist, A., Dahl, R., Ekblad, E., Hoffmann, H.J., Björner, L., Erjefält, J.S., 2014. Mapping of TLR5 and TLR7 in central and distal human airways and identification of reduced TLR expression in severe asthma. *Clin Exp Allergy*, 44, 184–196.
- Shimizu, T., Kida, Y., Kuwano, K., 2007. Triacylated lipoproteins derived from *Mycoplasma pneumoniae* activate nuclear factor- $\kappa$ B through Toll-like receptors 1 and 2. *Immunology*, 121, 473–483.

- Shinji, H., Yosizawa, Y., Tajima, A., Iwase, T., Sugimoto, S., Seki, K., Mizunoe, Y., 2011. Role of fibronectin-binding proteins A and B in *in vitro* cellular infections and *in vivo* septic infections by *Staphylococcus aureus*. *Infect Immun*, 79, 2215–2223.
- Shinjyo, N., Ståhlberg, A., Dragunow, M., Pekny, M., Pekna, M., 2009. Complement-derived anaphylatoxin C3a regulates *in vitro* differentiation and migration of neural progenitor cells. *Stem Cells*, 27, 2824–2832.
- Sibley, C.D., Duan, K., Fischer, C., Parkins, M.D., Storey, D.G., Rabin, H.R., Surette, M.G., 2008. Discerning the complexity of community interactions using a drosophila model of polymicrobial infections. *PLoS Pathog*, 4, e1000184.
- Sieprawska-Lupa, M., Mydel, P., Krawczyk, K., Wójcik, K., Puklo, M., Lupa, B., Suder, P., Silberring, J., Reed, M., Pohl, J., Shafer, W., McAleese, F., Foster, T., Travis, J., Potempa, J., 2004. Degradation of human antimicrobial peptide LL-37 by *Staphylococcus aureus*-derived proteinases. *Antimicrob Agents Chemother*, 48, 4673–4679.
- Silipo, A., Molinaro, A., Ieranò, T., De Soyza, A., Sturiale, L., Garozzo, D., Aldridge, C., Corris, P.A., Khan, C.M.A., Lanzetta, R., Parrilli, M., 2007. The complete structure and pro-inflammatory activity of the lipooligosaccharide of the highly epidemic and virulent Gram-negative bacterium *Burkholderia cenocepacia* ET-12 (Strain J2315). *Chemistry*, 13, 3501–3511.
- Silva, M.T., 2012. Classical labeling of bacterial pathogens according to their lifestyle in the host: inconsistencies and alternatives. *Front Microbio*, 3, 71.
- Singh, P.K., 2004. Iron sequestration by human lactoferrin stimulates *P. aeruginosa* surface motility and blocks biofilm formation. *BioMetals*, 17, 267–270.
- Singh, P.K., Jia, H.P., Wiles, K., Hesselberth, J., Liu, L., Conway, B.-A.D., Greenberg, E.P., Valore, E.V., Welsh, M.J., Ganz, T., Tack, B.F., McCray, P.B., 1998. Production of  $\beta$ -defensins by human airway epithelia. *Proc Natl Acad Sci*, 95, 14961–14966.
- Singh, P.K., Tack, B.F., McCray, P.B., Jr., Welsh, M.J., 2000. Synergistic and additive killing by antimicrobial factors found in human airway surface liquid. *Am J Physiol Lung Cell Mol Physiol*, 279, L799–L805.
- Singh, R., Sahore, S., Kaur, P., Rani, A., Ray, P., 2016. Penetration barrier contributes to bacterial biofilm-associated resistance against only select antibiotics, and exhibits genus-, strain- and antibiotic-specific differences. *Pathog Dis*, 74, ftw056.
- Singla, S., Harjai, K., Chhibber, S., 2012. Susceptibility of different phases of biofilm of *Klebsiella pneumoniae* to three different antibiotics. *J Antibiot*, 66, 61–66.
- Sinha, B., Francois, P.P., Nusse, O., Foti, M., Hartford, O.M., Vaudaux, P., Foster, T.J., Lew, D.P., Herrmann, M., Krause, K.H., 1999. Fibronectin-binding protein acts as *Staphylococcus aureus* invasin via fibronectin bridging to integrin  $\alpha 5\beta 1$ . *Cell Microbiol*, 1, 101–117.
- Siqueiros-Cendon, T., Arevalo-Gallegos, S., Iglesias-Figueroa, B.F., Garcia-Montoya, I.A., Salazar-Martinez, J., Rascon-Cruz, Q., 2014. Immunomodulatory effects of lactoferrin. *Acta Pharmacol Sin*, 35, 557–566.
- Snelgrove, R.J., Goulding, J., Didierlaurent, A.M., Lyonga, D., Vekaria, S., Edwards, L., Gwyer, E., Sedgwick, J.D., Barclay, A.N., Hussell, T., 2008. A critical function for CD200 in lung immune homeostasis and the severity of influenza infection. *Nat Immunol*, 9, 1074–1083.
- Soong, G., Martin, F.J., Chun, J., Cohen, T.S., Ahn, D.S., Prince, A., 2011. *Staphylococcus aureus* protein A mediates invasion across airway epithelial cells

- through activation of RhoA GTPase signaling and proteolytic activity. *J Biol Chem*, 286, 35891–35898.
- Spaan, A.N., Henry, T., van Rooijen, W.J.M., Perret, M., Badiou, C., Aerts, P.C., Kemmink, J., de Haas, C.J.C., van Kessel, K.P.M., Vandenesch, F., Lina, G., van Strijp, J.A.G., 2013a. The Staphylococcal toxin Pantón-Valentine Leukocidin targets human C5a receptors. *Cell Host Microbe*, 13, 584–594.
- Spaan, A.N., Reyes-Robles, T., Badiou, C., Cochet, S., Boguslawski, K.M., Yoong, P., Day, C.J., de Haas, C.J.C., van Kessel, K.P.M., Vandenesch, F., Jennings, M.P., Le Van Kim, C., Colin, Y., van Strijp, J.A.G., Henry, T., Torres, V.J., 2015. *Staphylococcus aureus* targets the duffy antigen receptor for chemokines (DARC) to lyse erythrocytes. *Cell Host Microbe*, 18, 363–370.
- Spaan, A.N., Surewaard, B.G.J., Nijland, R., van Strijp, J.A.G., 2013b. Neutrophils versus *Staphylococcus aureus*: a biological tug of war. *Annu Rev Microbiol*, 67, 629–650.
- Spaan, A.N., van Strijp, J.A.G., Torres, V.J., 2017. Leukocidins: Staphylococcal bi-component pore-forming toxins find their receptors. *Nat Rev Micro*, 15, 435–447.
- Spaan, A.N., Vrieling, M., Wallet, P., Badiou, C., Reyes-Robles, T., Ohneck, E.A., Benito, Y., de Haas, C.J.C., Day, C.J., Jennings, M.P., Lina, G., Vandenesch, F., van Kessel, K.P.M., Torres, V.J., van Strijp, J.A.G., Henry, T., 2014. The Staphylococcal toxins  $\gamma$ -haemolysin AB and CB differentially target phagocytes by employing specific chemokine receptors. *Nat Commun*, 5, 5438.
- Stapels, D.A.C., Ramyar, K.X., Bischoff, M., Köckritz-Blickwede, von, M., Milder, F.J., Ruyken, M., Eisenbeis, J., McWhorter, W.J., Herrmann, M., van Kessel, K.P.M., Geisbrecht, B.V., Rooijackers, S.H.M., 2014. *Staphylococcus aureus* secretes a unique class of neutrophil serine protease inhibitors. *Proc Natl Acad Sci*, 111, 13187–13192.
- Stewart, C.E., Torr, E.E., Mohd Jamili, N.H., Bosquillon, C., Sayers, I., 2012. Evaluation of differentiated human bronchial epithelial cell culture systems for asthma research. *J Allergy*, 2012, 1–11.
- Stout, S.L., Wyatt, T.A., Adams, J.J., Sisson, J.H., 2007. Nitric oxide-dependent cilia regulatory enzyme localization in bovine bronchial epithelial cells. *J Histochem Cytochem*, 55, 433–442.
- Strassheim, D., Asehnoune, K., Park, J.-S., Kim, J.-Y., He, Q., Richter, D., Kuhn, K., Mitra, S., Abraham, E., 2004. Phosphoinositide 3-kinase and Akt occupy central roles in inflammatory responses of Toll-like receptor 2-stimulated neutrophils. *J Immunol*, 172, 5727–5733.
- Strebovsky, J., Walker, P., Lang, R., Dalpke, A.H., 2011. Suppressor of cytokine signaling 1 limits NF- $\kappa$ B signaling by decreasing p65 stability within the cell nucleus. *FASEB J*, 25, 863–874.
- Strobel, M., Pförtner, H., Tuchscher, L., Völker, U., Schmidt, F., Kramko, N., Schnittler, H.J., Fraunholz, M.J., Löffler, B., Peters, G., Niemann, S., 2016. Post-invasion events after infection with *Staphylococcus aureus* are strongly dependent on both the host cell type and the infecting *S. aureus* strain. *Clin Microbiol Infect*, 22, 799–809.
- Stutman, H.R., Lieberman, J.M., Nussbaum, E., Marks, M.I., 2002. Antibiotic prophylaxis in infants and young children with cystic fibrosis: A randomized controlled trial. *J Pediatr*, 140, 299–305.
- Sun, S.-C., 2010. CYLD: a tumor suppressor deubiquitinase regulating NF- $\kappa$ B activation and diverse biological processes. *Cell Death Differ*, 17, 25–34.
- Surette, M.G., 2014. The cystic fibrosis lung microbiome. *Ann Am Thorac Soc*, 1, S61–65.

- Surewaard, B.G.J., de Haas, C.J.C., Vervoort, F., Rigby, K.M., DeLeo, F.R., Otto, M., van Strijp, J.A.G., Nijland, R., 2013. Staphylococcal  $\alpha$ -phenol soluble modulins contribute to neutrophil lysis after phagocytosis. *Cell Microbiol*, 15, 1427–1437.
- Szondy, Z., Sarang, Z., Kiss, B., Garabuczi, É., Köröskényi, K., 2017. Anti-inflammatory mechanisms triggered by apoptotic cells during their clearance. *Front Immunol*, 8, 909.
- Tajima, A., Iwase, T., Shinji, H., Seki, K., Mizunoe, Y., 2008. Inhibition of endothelial interleukin-8 production and neutrophil transmigration by *Staphylococcus aureus*  $\beta$ -hemolysin. *Infect Immun*, 77, 327–334.
- Tajima, A., Seki, K., Shinji, H., Masuda, S., 2006. Inhibition of interleukin-8 production in human endothelial cells by *Staphylococcus aureus* supernatant. *Clin Exp Immunol*, 147, 148–154.
- Takeuchi, O., Akira, S., 2010. Pattern recognition receptors and inflammation. *Cell*, 140, 805–820.
- Takeuchi, O., Hoshino, K., Akira, S., 2000. Cutting edge: TLR2-deficient and MyD88-deficient mice are highly susceptible to *Staphylococcus aureus* infection. *J Immunol*, 165, 5392–5396.
- Tanimura, N., Saitoh, S., Matsumoto, F., Akashi-Takamura, S., Miyake, K., 2008. Roles for LPS-dependent interaction and relocation of TLR4 and TRAM in TRIF-signaling. *Biochem Biophys Res Commun*, 368, 94–99.
- Tauszig, S., Jouanguy, E., Hoffmann, J.A., Imler, J.L., 2000. Toll-related receptors and the control of antimicrobial peptide expression in *Drosophila*. *Proc Natl Acad Sci*, 97, 10520–10525.
- Thakker, M., Park, J.S., Carey, V., Lee, J.C., 1998. *Staphylococcus aureus* serotype 5 capsular polysaccharide is antiphagocytic and enhances bacterial virulence in a murine bacteremia model. *Infect Immun*, 66, 5183–5189.
- Thammavongsa, V., Kim, H.K., Missiakas, D., Schneewind, O., 2015. Staphylococcal manipulation of host immune responses. *Nat Rev Micro*, 13, 529–543.
- Thepen, T., Van Rooijen, N., Kraal, G., 1989. Alveolar macrophage elimination *in vivo* is associated with an increase in pulmonary immune response in mice. *J Exp Med*, 170, 499–509.
- Thomas, M.S., 2007. Iron acquisition mechanisms of the *Burkholderia cepacia* complex. *BioMetals*, 20, 431–452.
- Thurlow, L.R., Hanke, M.L., Fritz, T., Angle, A., Aldrich, A., Williams, S.H., Engebretsen, I.L., Bayles, K.W., Horswill, A.R., Kielian, T., 2011. *Staphylococcus aureus* biofilms prevent macrophage phagocytosis and attenuate inflammation *in vivo*. *J Immunol*, 186, 6585–6596.
- Tomich, M., Herfst, C.A., Golden, J.W., Mohr, C.D., 2002. Role of flagella in host cell invasion by *Burkholderia cepacia*. *Infect Immun*, 70, 1799–1806.
- Tong, S.Y.C., Davis, J.S., Eichenberger, E., Holland, T.L., Fowler, V.G., 2015. *Staphylococcus aureus* infections: epidemiology, pathophysiology, clinical manifestations, and management. *Clin Microbiol Rev*, 28, 603–661.
- Traber, K. and Novick, R., 2006. A slipped-mispairing mutation in AgrA of laboratory strains and clinical isolates results in delayed activation of *agr* and failure to translate  $\delta$ - and  $\alpha$ -haemolysins. *Mol Microbiol*, 59, 1519–1530.
- Tranchemontagne, Z.R., Camire, R.B., O'Donnell, V.J., Baugh, J., Burkholder, K.M., 2016. *Staphylococcus aureus* strain USA300 perturbs acquisition of lysosomal enzymes and



- requires phagosomal acidification for survival inside macrophages. *Infect Immun*, 84, 241–253.
- Trifilieff, A., Fujitani, Y., Mentz, F., Dugas, B., Fuentes, M., Bertrand, C., 2000. Inducible nitric oxide synthase inhibitors suppress airway inflammation in mice through down-regulation of chemokine expression. *J Immunol*, 165, 1526–1533.
- Trinchieri, G., 2010. Type I interferon: friend or foe? *J Exp Med*, 207, 2053–2063.
- Troeger, C., Forouzanfar, M., Rao, P.C., Khalil, I., Brown, A., Swartz, S., Fullman, N., Mosser, J., Thompson, R.L., Reiner, R.C., Abajobir, A., Alam, N., Alemayohu, M.A., Amare, A.T., Antonio, C.A., Asayesh, H., Avokpaho, E., Barac, A., Beshir, M.A., Boneya, D.J., Brauer, M., Dandona, L., Dandona, R., Fitchett, J.R.A., Gebrehiwot, T.T., Hailu, G.B., Hotez, P.J., Kasaeian, A., Khoja, T., Kissoon, N., Knibbs, L., Kumar, G.A., Rai, R.K., Razek, El, H.M.A., Mohammed, M.S.K., Nielson, K., Oren, E., Osman, A., Patton, G., Qorbani, M., Roba, H.S., Sartorius, B., Savic, M., Shigematsu, M., Sykes, B., Swaminathan, S., Topor-Madry, R., Ukwaja, K., Werdecker, A., Yonemoto, N., Sayed Zaki, El, M., Lim, S.S., Naghavi, M., Vos, T., Hay, S.I., Murray, C.J.L., Mokdad, A.H., 2017. Estimates of the global, regional, and national morbidity, mortality, and aetiologies of lower respiratory tract infections in 195 countries: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet Infect Dis*, 17, 1133–1161.
- Troutman, T.D., Bazan, J.F., Pasare, C., 2014. Toll-like receptors, signaling adapters and regulation of the pro-inflammatory response by PI3K. *Cell Cycle*, 11, 3559–3567.
- Trzciński, K., van Leeuwen, W., van Belkum, A., Grzesiowski, P., Kluytmans, J., Sijmons, M., Verbrugh, H., Witte, W., Hryniewicz, W., 1997. Two clones of methicillin-resistant *Staphylococcus aureus* in Poland. *Clin Microbiol Infect*, 3, 198–207.
- Udo, E.E., 2013. Community-acquired methicillin-resistant *Staphylococcus aureus*: the new face of an old foe? *Med Princ Pract*, 22 suppl 1, 20–29.
- Uematsu, S., Akira, S., 2007. Toll-like receptors and type I interferons. *J Biol Chem*, 282, 15319–15323.
- Ullah, M.O., Sweet, M.J., Mansell, A., Kellie, S., Kobe, B., 2016. TRIF-dependent TLR signaling, its functions in host defense and inflammation, and its potential as a therapeutic target. *J Leukoc Biol*, 100, 27–45.
- Urban, T.A., Goldberg, J.B., Forstner, J.F., Sajjan, U.S., 2005. Cable pili and the 22-kDa adhesin are required for *Burkholderia cenocepacia* binding to and transmigration across the squamous epithelium. *Infect Immun*, 73, 5426–5437.
- Urban, T.A., Griffith, A., Torok, A.M., Smolkin, M.E., Burns, J.L., Goldberg, J.B., 2004. Contribution of *Burkholderia cenocepacia* flagella to infectivity and inflammation. *Infect Immun*, 72, 5126–5134.
- Uzoamaka, M., Ngozi, O., Johnbull, O.S., Martin, O., 2017. Bacterial etiology of lower respiratory tract infections and their antimicrobial susceptibility. *Am J Med Sci*, 354, 471–475.
- Valkov, E., Stamp, A., Dimaio, F., Baker, D., Verstak, B., Roversi, P., Kellie, S., Sweet, M.J., Mansell, A., Gay, N.J., Martin, J.L., Kobe, B., 2011. Crystal structure of Toll-like receptor adaptor MAL/TIRAP reveals the molecular basis for signal transduction and disease protection. *Proc Natl Acad Sci*, 108, 14879–14884.
- van Belkum, A., 2016. Hidden *Staphylococcus aureus* carriage: overrated or underappreciated? *MBio*, 7, e00079–16.
- van Kessel, K.P.M., Bestebroer, J., van Strijp, J.A.G., 2014. Neutrophil-mediated phagocytosis of *Staphylococcus aureus*. *Front Immunol*, 5, 467.

- van Krüchten, A., Wilden, J.J., Niemann, S., Peters, G., Löffler, B., Ludwig, S., Ehrhardt, C., 2018. *Staphylococcus aureus* triggers a shift from influenza virus-induced apoptosis to necrotic cell death. *FASEB J*, 32, 2779–2793.
- Vandecandelaere, I., Van Nieuwerburgh, F., Deforce, D., Coenye, T., 2017. Metabolic activity, urease production, antibiotic resistance and virulence in dual species biofilms of *Staphylococcus epidermidis* and *Staphylococcus aureus*. *PLoS ONE*, 12, e0172700.
- Vandivier, R.W., Fadok, V.A., Ogden, C.A., Hoffmann, P.R., Brain, J.D., Accurso, F.J., Fisher, J.H., Greene, K.E., Henson, P.M., 2002. Impaired clearance of apoptotic cells from cystic fibrosis airways. *Chest*, 121, 89S.
- Venugopalan, C.S., Krautmann, M.J., Holmes, E.P., Maher, T.J., 1998. Involvement of nitric oxide in the mediation of NANC inhibitory neurotransmission of guinea-pig trachea. *J Auton Pharmacol*, 18, 281–286.
- Vergadi, E., Vaporidi, K., Theodorakis, E.E., Doxaki, C., Lagoudaki, E., Ieronymaki, E., Alexaki, V.I., Helms, M., Kondili, E., Soennichsen, B., Stathopoulos, E.N., Margioris, A.N., Georgopoulos, D., Tsatsanis, C., 2014. Akt2 deficiency protects from acute lung injury via alternative macrophage activation and miR-146a induction in mice. *J Immunol*, 192, 394–406.
- Vinion-Dubiel, A.D., Goldberg, J.B., 2003. Lipopolysaccharide of *Burkholderia cepacia* complex. *J Endotoxin Res*, 9, 201–213.
- Vlahos, C.J., Matter, W.F., Hui, K.Y., Brown, R.F., 1994. A specific inhibitor of phosphatidylinositol 3-kinase, 2-(4-morpholinyl)-8-phenyl-4H-1-benzopyran-4-one (LY294002). *J Biol Chem*, 269, 5241–5248.
- Voyich, J.M., Otto, M., Mathema, B., Braughton, K.R., Whitney, A.R., Welty, D., Long, R.D., Dorward, D.W., Gardener, D.J., Lina, G., Kreiswirth, B.N., DeLeo, F.R., 2006. Is panton-valentine leukocidin the major virulence determinant in community-associated methicillin-resistant *Staphylococcus aureus* disease? *J Infect Dis*, 194, 1761–1770.
- Vroiling, A.B., Konijn, T., Samsom, J.N., Kraal, G., 2011. The production of secretory leukocyte protease inhibitor by dendritic cells. *Mol Immunol*, 48, 630–636.
- Wald, D., Qin, J., Zhao, Z., Qian, Y., Naramura, M., Tian, L., Towne, J., Sims, J.E., Stark, G.R., Li, X., 2003. SIGIRR, a negative regulator of Toll-like receptor-interleukin 1 receptor signaling. *Nat Immunol*, 4, 920–927.
- Walenkamp, A.M., Boer, I.G., Bestebroer, J., Rozeveld, D., Timmer-Bosscha, H., Hemrika, W., van Strijp, J.A., de Haas, C.J., 2009. Staphylococcal Superantigen-like 10 Inhibits CXCL12-Induced Human Tumor Cell Migration. *Neoplasia*, 11, 333–344.
- Walker, E.H., Pacold, M.E., Perisic, O., Stephens, L., Hawkins, P.T., Wymann, M.P., Williams, R.L., 2000. Structural determinants of phosphoinositide 3-kinase inhibition by wortmannin, LY294002, quercetin, myricetin, and staurosporine. *Mol Cell*, 6, 909–919.
- Walters, M.C.3., Roe, F., Bugnicourt, A., Franklin, M.J., Stewart, P.S., 2003. Contributions of antibiotic penetration, oxygen limitation, and low metabolic activity to tolerance of *Pseudomonas aeruginosa* biofilms to ciprofloxacin and tobramycin. *Antimicrob Agents Chemother*, 47, 317–323.
- Wang, C., Deng, L., Hong, M., Akkaraju, G.R., Inoue, J., Chen, Z.J., 2001. TAK1 is a ubiquitin-dependent kinase of MKK and IKK. *Nature*, 412, 346–351.
- Wang, J., Roderiquez, G., Norcross, M.A., 2012. Control of adaptive immune responses by *Staphylococcus aureus* through IL-10, PD-L1, and TLR2. *Sci Rep*, 2, 606.
- Wang, R., Braughton, K.R., Kretschmer, D., Bach, T.H., Queck, S.Y., Li, M., Kennedy, A.D., Dorward, D.W., Klebanoff, S.J., Peschel, A., DeLeo, F.R., Otto, M., 2007.

- Identification of novel cytolytic peptides as key virulence determinants for community-associated MRSA. *Nat Med*, 13:1510–1514.
- Wann, E.R., Gurusiddappa, S., Höök, M., 2000. The fibronectin-binding MSCRAMM FnbpA of *Staphylococcus aureus* is a bifunctional protein that also binds to fibrinogen. *J Biol Chem*, 275, 13863–13871.
- Waters, V., Yau, Y., Beaudoin, T., Wettlaufer, J., Tom, S.K., McDonald, N., Rizvi, L., Klingel, M., Ratjen, F., Tullis, E., 2017. Pilot trial of tobramycin inhalation powder in cystic fibrosis patients with chronic *Burkholderia cepacia* complex infection. *J Cyst Fibros*, 16, 492–495.
- Watford, W.T., Wright, J.R., Hester, C.G., Jiang, H., Frank, M.M., 2001. Surfactant protein A regulates complement activation. *J Immunol*, 167, 6593–6600.
- Weibel, E.R., 2015. On the tricks alveolar epithelial cells play to make a good lung. *Am J Respir Crit Care Med*, 191, 504–513.
- Weichhart, T., Säemann, M.D., 2008. The PI3K/Akt/mTOR pathway in innate immune cells: emerging therapeutic applications. *Ann Rheum Dis*, 67 Suppl 3, iii70–4.
- Weigel, L.M., Donlan, R.M., Shin, D.H., Jensen, B., Clark, N.C., McDougal, L.K., Zhu, W., Musser, K.A., Thompson, J., Kohlerschmidt, D., Dumas, N., Limberger, R.J., Patel, J.B., 2007. High-level vancomycin-resistant *Staphylococcus aureus* isolates associated with a polymicrobial biofilm. *Antimicrob Agents Chemother*, 51, 231–238.
- Weinlich R., Green D.R., 2014. The two faces of receptor interacting protein kinase-1. *Mol Cell*, 56(4):469-80.
- Weiss, D.S., Takeda, K., Akira, S., Zychlinsky, A., Moreno, E., 2005. MyD88, but not Toll-like receptors 4 and 2, is required for efficient clearance of *Brucella abortus*. *Infect Immun*, 73, 5137–5143.
- Weitnauer, M., Mijošek, V., Dalpke, A.H., 2016. Control of local immunity by airway epithelial cells. *Mucosal Immunol*, 9, 287–298.
- Wertheim, H.F., Melles, D.C., Vos, M.C., van Leeuwen, W., van Belkum, A., Verbrugh, H.A., Nouwen, J.L., 2005. The role of nasal carriage in *Staphylococcus aureus* infections. *Lancet Infect Dis*, 5, 751–762.
- West, T.E., Ernst, R.K., Jansson-Hutson, M.J., Skerrett, S.J., 2008. Activation of Toll-like receptors by *Burkholderia pseudomallei*. *BMC Immunol*, 9, 46.
- West, T.E., Hawn, T.R., Skerrett, S.J., 2009. Toll-like receptor signaling in airborne *Burkholderia thailandensis* infection. *Infect Immun*, 77, 5612–5622.
- Whitsett, J.A., Alenghat, T., 2015. Respiratory epithelial cells orchestrate pulmonary innate immunity. *Nat Immunol*, 16, 27–35.
- Whitsett, J.A., Wert, S.E., Weaver, T.E., 2010. Alveolar surfactant homeostasis and the pathogenesis of pulmonary disease. *Annu Rev Med*, 61, 105–119.
- Wiersinga, W.J., van der Poll, T., White, N.J., Day, N.P., Peacock, S.J., 2006. Melioidosis: insights into the pathogenicity of *Burkholderia pseudomallei*. *Nat Rev Micro*, 4, 272–282.
- Williams, R.E., 1963. Healthy carriage of *Staphylococcus aureus*: its prevalence and importance. *Bacteriol Rev*, 27, 56–71.
- Winder, A.A., Wohlford-Lenane, C., Scheetz, T.E., Nardy, B.N., Manzel, L.J., Look, D.C., McCray, P.B., 2009. Differential effects of cytokines and corticosteroids on Toll-like receptor 2 expression and activity in human airway epithelia. *Respir Res*, 10, 675–11.

- Wintersdorff, von, C.J.H., Penders, J., van Niekerk, J.M., Mills, N.D., Majumder, S., van Alphen, L.B., Savelkoul, P.H.M., Wolffs, P.F.G., 2016. Dissemination of antimicrobial resistance in microbial ecosystems through horizontal gene transfer. *Front Microbio*, 7, 173.
- Wolcott, R., Costerton, J.W., Raoult, D., Cutler, S.J., 2013. The polymicrobial nature of biofilm infection. *Clin Microbiol Infect*, 19, 107–112.
- Wood, T.K., Knabel, S.J., Kwan, B.W., 2013. Bacterial persister cell formation and dormancy. *J Appl Environ Microbiol*, 79, 7116–7121.
- Woodhead, M., Blasi, F., Ewig, S., Garau, J., Huchon, G., Ieven, M., Ortqvist, A., Schaberg, T., Torres, A., van der Heijden, G., Read, R., Verheij, T.J.M., 2011. Guidelines for the management of adult lower respiratory tract infections. *Clin Microbiol Infect*, 17, Suppl 6, E1–59.
- Workman, A.D., Carey, R.M., Kohanski, M.A., Kennedy, D.W., Palmer, J.N., Adappa, N.D., Cohen, N.A., 2017. Relative susceptibility of airway organisms to antimicrobial effects of nitric oxide. *Int Forum Allergy Rhinol*, 7, 770–776.
- Yokoyama, R., Itoh, S., Kamoshida, G., Takii, T., Fujii, S., Tsuji, T., Onozaki, K., 2012. Staphylococcal superantigen-like protein 3 binds to the Toll-like receptor 2 extracellular domain and inhibits cytokine production induced by *Staphylococcus aureus*, cell wall component, or lipopeptides in murine macrophages. *Infect Immun*, 80, 2816–2825.
- Yoshimura, A., Muto, G., 2011. TGF- $\beta$  function in immune suppression. *Curr Top Microbiol Immunol*, 350, 127–147.
- Yu, D., Marchiando, A.M., Weber, C.R., Raleigh, D.R., Wang, Y., Shen, L., Turner, J.R., 2010. MLCK-dependent exchange and actin binding region-dependent anchoring of ZO-1 regulate tight junction barrier function. *Proc Natl Acad Sci*, 107, 8237–8241.
- Yu, H., Liu, Y., McFarland, B.C., Deshane, J.S., Hurst, D.R., Ponnazhagan, S., Benveniste, E.N., Qin, H., 2015. SOCS3 deficiency in myeloid cells promotes tumor development: involvement of STAT3 activation and myeloid-derived suppressor cells. *Cancer Immunol Res*, 3, 727–740.
- Yu, Y.-R.A., Hotten, D.F., Malakhau, Y., Volker, E., Ghio, A.J., Noble, P.W., Kraft, M., Hollingsworth, J.W., Gunn, M.D., Tighe, R.M., 2016. Flow cytometric analysis of myeloid cells in human blood, bronchoalveolar lavage, and lung tissues. *Am J Respir Cell Mol Biol*, 54, 13–24.
- Yuan, H., Barnes, K.R., Weissleder, R., Cantley, L., Josephson, L., 2007. Covalent reactions of wortmannin under physiological conditions. *Chem Biol*, 14, 321–328.
- Zarembek, K.A., Godowski, P.J., 2002. Tissue expression of human Toll-like receptors and differential regulation of Toll-like receptor mRNAs in leukocytes in response to microbes, their products, and cytokines. *J Immunol*, 168, 554–561.
- Zhang, S., Cherwinski, H., Sedgwick, J.D., Phillips, J.H., 2004. Molecular mechanisms of CD200 inhibition of mast cell activation. *J Immunol*, 173, 6786–6793.
- Zhang, X., Li, N., Shao, H., Meng, Y., Wang, L., Wu, Q., Yao, Y., Li, J., Bian, J., Zhang, Y., Deng, X., 2016. Methane limit LPS-induced NF- $\kappa$ B/MAPKs signal in macrophages and suppress immune response in mice by enhancing PI3K/AKT/GSK-3 $\beta$ -mediated IL-10 expression. *Sci Rep*, 6, 2085.
- Zhang, Z., Ohto, U., Shibata, T., Krayukhina, E., Taoka, M., Yamauchi, Y., Tanji, H., Isobe, T., Uchiyama, S., Miyake, K., Shimizu, T., 2016. Structural analysis reveals that Toll-like receptor 7 is a dual receptor for guanosine and single-stranded RNA. *Immunity*, 45, 737–748.

- Zhou, X., Liu, Z., Cheng, X., Zheng, Y., Zeng, F., He, Y., 2015. Socs1 and Socs3 degrades Traf6 via polyubiquitination in LPS-induced acute necrotizing pancreatitis. *Cell Death Dis*, 6, e2012.
- Zlosnik, J.E., Costa, P.S., Brant, R., Mori, P.Y., Hird, T.J., Fraenkel, M.C., Wilcox, P.G., Davidson, A.G.F., Speert, D.P., 2011. Mucoïd and nonmucoïd *Burkholderia cepacia* complex bacteria in cystic fibrosis infections. *Am J Respir Crit Care Med*, 183, 67–72.
- Zlosnik, J.E.A., Zhou, G., Brant, R., Henry, D.A., Hird, T.J., Mahenthiralingam, E., Chilvers, M.A., Wilcox, P., Speert, D.P., 2015. *Burkholderia* species infections in patients with cystic fibrosis in British Columbia, Canada. 30 years' experience. *Ann Am Thorac Soc*, 12, 70–78.
- Zong, Y., Xu, Y., Liang, X., Keene, D.R., Höök, A., Gurusiddappa, S., Höök, M., Narayana, S.V.L., 2005. A 'Collagen Hug' Model for *Staphylococcus aureus* CNA binding to collagen. *EMBO J*, 24, 4224–4236.
- Zughaier, S.M., Ryley, H.C., Jackson, S.K., 1999. Lipopolysaccharide (LPS) from *Burkholderia cepacia* is more active than LPS from *Pseudomonas aeruginosa* and *Stenotrophomonas maltophilia* in stimulating tumor necrosis factor  $\alpha$  from human monocytes. *Infect Immun*, 67, 1505–1507.